



**KAPLAN**  
MEDICAL



**USMLE™ Step 1**  
**Lecture Notes**

**Pharmacology**



# **USMLE Step 1**

**Lecture Notes**

**Pharmacology**

©2009 Kaplan, Inc.

All rights reserved. No part of this book may be reproduced in any form, by photostat, microfilm, xerography or any other means, or incorporated into any information retrieval system, electronic or mechanical, without the written permission of Kaplan, Inc.

Not for resale.

## **Author**

Lionel P. Raymon, Pharm.D., Ph.D.

*Department of Pathology*

*Forensic Toxicology Laboratory*

*University of Miami School of Medicine*

*Miami, FL*

---

## **Coauthors**

Craig Davis, Ph.D.

*Associate Professor*

*University of South Carolina School of Medicine*

*Department of Pharmacology, Physiology, and Neuroscience*

*Columbia, SC*

Steven R. Harris, Ph.D.

*Associate Dean for Academic Affairs*

*Associate Professor of Pharmacology*

*Pikeville College School of Osteopathic Medicine*

*Pikeville, KY*

## **Contributors**

Maris Victor Nora, Pharm.D., Ph.D.

*Associate Professor*

*Rush Medical College*

*Chicago, IL*

Anthony Trevor, Ph.D.

*Professor Emeritus*

*Department of Cellular and Molecular Pharmacology*

*University of California*

*San Francisco, CA*

---

## **Director of Medical Curriculum**

Mark Tyler-Lloyd, M.D.

## **Directors of Step 1 Curriculum**

Michael S. Manley, M.D.

Leslie D. Manley, Ph.D.

## **Associate Director of Medical Curriculum**

Shefali Vyas, M.D.

## **Editorial Director**

Ruth Baygell

## **Production Manager**

Michael Wolff

## **Medical Illustrators**

Rich LaRocco

Christine Schaar



# Contents

---

Preface .....ix

## **Section I: General Principles**

**Chapter 1:** Pharmacokinetics ..... 3

**Chapter 2:** Pharmacodynamics ..... 19

**Chapter 3:** Practice Questions ..... 29

## **Section II: Autonomic Pharmacology**

**Chapter 1:** The Autonomic Nervous System (ANS) ..... 39

**Chapter 2:** Cholinergic Pharmacology ..... 45

**Chapter 3:** Adrenergic Pharmacology ..... 53

**Chapter 4:** Autonomic Drugs: Glaucoma Treatment and  
ANS Practice Problems ..... 61

**Chapter 5:** Autonomic Drug List and Practice Questions ..... 69

## **Section III: Cardiac and Renal Pharmacology**

**Chapter 1:** Fundamental Concepts ..... 83

**Chapter 2:** Antiarrhythmic Drugs ..... 89

**Chapter 3:** Antihypertensive Drugs ..... 93

**Chapter 4:** Drugs for Heart Failure ..... 99

**Chapter 5:** Antianginal Drugs ..... 103

<b>Chapter 6: Diuretics</b> .....	107
<b>Chapter 7: Antihyperlipidemics</b> .....	113
<b>Chapter 8: Cardiac and Renal Drug List and Practice Questions</b> .....	117

## **Section IV: CNS Pharmacology**

<b>Chapter 1: Sedative-Hypnotic-Anxiolytic Drugs</b> .....	129
<b>Chapter 2: Alcohols</b> .....	133
<b>Chapter 3: Anticonvulsants</b> .....	135
<b>Chapter 4: Drugs Used in Anesthesia</b> .....	139
<b>Chapter 5: Opioid Analgesics</b> .....	145
<b>Chapter 6: Drugs Used in Parkinson Disease and Psychosis</b> .....	149
<b>Chapter 7: Drugs Used for Depression, Bipolar Disorders, and Attention Deficit Hyperactivity Disorder (ADHD)</b> .....	155
<b>Chapter 8: Drugs of Abuse</b> .....	159
<b>Chapter 9: CNS Drug List and Practice Questions</b> .....	163

## **Section V. Antimicrobial Agents**

<b>Chapter 1: Antibacterial Agents</b> .....	175
<b>Chapter 2: Antifungal Agents</b> .....	191
<b>Chapter 3: Antiviral Agents</b> .....	195
<b>Chapter 4: Antiprotozoal Agents</b> .....	203
<b>Chapter 5: Antimicrobial Drug List and Practice Questions</b> .....	205

## **Section VI. Drugs for Inflammatory and Related Disorders**

<b>Chapter 1: Histamine and Antihistamines</b> .....	219
<b>Chapter 2: Drugs Used in Gastrointestinal Dysfunction</b> .....	221

<b>Chapter 3: Drugs Acting on Serotonergic Systems</b> .....	225
<b>Chapter 4: Eicosanoid Pharmacology</b> .....	227
<b>Chapter 5: Drugs Used for Treatment of Rheumatoid Arthritis</b> .....	233
<b>Chapter 6: Drugs Used for Treatment of Gout</b> .....	235
<b>Chapter 7: Glucocorticoids</b> .....	237
<b>Chapter 8: Drugs Used for Treatment of Asthma</b> .....	239
<b>Chapter 9: List of Drugs for Inflammatory Disorders and Practice Questions</b> ..	243

## **Section VII: Drugs Used in Blood Disorders**

<b>Chapter 1: Anticoagulants</b> .....	257
<b>Chapter 2: Thrombolytics</b> .....	261
<b>Chapter 3: Antiplatelet Drugs</b> .....	263
<b>Chapter 4: List of Drugs Used in Blood Disorders and Practice Questions</b> .....	265

## **Section VIII: Endocrine Pharmacology**

<b>Chapter 1: Drugs Used in Diabetes</b> .....	271
<b>Chapter 2: Steroid Hormones</b> .....	275
<b>Chapter 3: Antithyroid Agents</b> .....	281
<b>Chapter 4: Drugs Related to Hypothalamic and Pituitary Hormones</b> .....	283
<b>Chapter 5: Drugs Used for Bone and Mineral Disorders</b> .....	285
<b>Chapter 6: Endocrine Drug List and Practice Questions</b> .....	287

## **Section IX: Anticancer Drugs**

<b>Chapter 1: Anticancer Drugs</b> .....	295
<b>Chapter 2: Anticancer Drug Practice Questions</b> .....	299



**Section X: Immunopharmacology**

**Chapter 1: Immunopharmacology** ..... 303

**Chapter 2: Immunopharmacology Practice Questions** ..... 305

**Section XI: Toxicology**

**Chapter 1: Toxicology**..... 309

**Chapter 2: Toxicology Practice Questions**..... 315

**Index** ..... 317

# Preface

---

**These** seven volumes of Lecture Notes represent the most-likely-to-be-tested material on the **current** USMLE Step 1 exam. Please note that these are Lecture Notes, not review books. The **Notes** were designed to be accompanied by faculty lectures—live, on DVD, or on the web. **Reading** these Notes without accessing the accompanying lectures is not an effective way to **review** for the USMLE.

To maximize the effectiveness of these Notes, annotate them as you listen to lectures. To facilitate this process, we've created wide, blank margins. While these margins are occasionally punctuated by faculty high-yield "margin notes," they are, for the most part, left blank for your notations.

Many students find that previewing the Notes prior to the lecture is a very effective way to prepare for class. This allows you to anticipate the areas where you'll need to pay particular attention. It also affords you the opportunity to map out how the information is going to be presented and what sorts of study aids (charts, diagrams, etc.) you might want to add. This strategy works regardless of whether you're attending a live lecture or watching one on video or the web.

Finally, we want to hear what you think. What do you like about the notes? What do you think could be improved? Please share your feedback by E-mailing us at [medfeedback@kaplan.com](mailto:medfeedback@kaplan.com).

Thank you for joining Kaplan Medical, and best of luck on your Step 1 exam!

**Kaplan Medical**



**SECTION I**

**General Principles**



# Pharmacokinetics

Pharmacokinetic characteristics of drug molecules concern the processes of absorption, distribution, metabolism, and excretion. The biodisposition of a drug involves its permeation across cellular membrane barriers.




Figure I-1-1. Drug Biodisposition

**In A Nutshell**

For Weak Acids and Weak Bases

Ionized = Water soluble

Nonionized = Lipid soluble

**PERMEATION**

- Drug permeation is dependent on:
  - **Solubility.** Ability to diffuse through lipid bilayers (lipid solubility) is important for most drugs; however, water solubility can influence permeation through aqueous phases.
  - **Concentration gradient.** Diffusion down a concentration gradient—only free, unionized drug forms contribute to the concentration gradient.
  - **Surface area and vascularity.** Important with regard to absorption of drugs into the systemic circulation. The larger the surface area and the greater the vascularity, the better is the absorption of the drug.
- Ionization
  - Many drugs are weak acids or weak bases and can exist in either nonionized or ionized forms in an equilibrium, depending on the pH of the environment and the pKa (the pH at which the molecule is 50% ionized and 50% nonionized)
  - Only the nonionized (uncharged) form of a drug crosses biomembranes.
  - The ionized form is better renally excreted because it is water soluble.





Figure I-1-2. Degree of Ionization and Clearance Versus pH Deviation from pKa

## Ionization Increases Renal Clearance of Drugs

- Only free, unbound drug is filtered.
- Both ionized and nonionized forms of a drug are filtered.
- Only nonionized forms undergo active secretion and active or passive reabsorption.
- Ionized forms of drugs are “trapped” in the filtrate.
- Acidification of urine → increases ionization of weak bases → increases renal elimination.
- Alkalinization of urine → increases ionization of weak acids → increases renal elimination.




Figure I-1-3. Renal Clearance of Drug

## Modes of Drug Transport Across a Membrane

Table I-1-1. The Three Basic Modes of Drug Transport Across a Membrane

Mechanism	Direction	Energy Required	Carrier	Saturable
Passive diffusion	Down gradient	No	No	No
Facilitated diffusion	Down gradient	No	Yes	Yes
Active transport	Against gradient (concentration/electrical)	Yes	Yes	Yes

### Clinical Correlate

#### To Change Urinary pH

- Acidify:  $\text{NH}_4\text{Cl}$ , vitamin C, cranberry juice
- Alkalinize:  $\text{NaHCO}_3$ , acetazolamide (historically)
- See Aspirin Overdose and Management in Section VI.

### Bridge to Physiology


Ion and molecular transport mechanisms are discussed in greater detail in Section I of Physiology.



### ABSORPTION

- Concerns the processes of entry of a drug into the systemic circulation from the site of its administration.
- The determinants of absorption are those described for drug permeation.
- Intravascular administration (e.g., IV) does not involve absorption, and there is no loss of drug. Bioavailability = 100%
- With extravascular administration (e.g., per os [PO; oral], intramuscular [IM], subcutaneous [SC], inhalation), less than 100% of a dose may reach the systemic circulation because of variations in bioavailability.

### Plasma Level Curves



$C_{max}$  = maximal drug level obtained with the dose.  
 $t_{max}$  = time at which  $C_{max}$  occurs.  
 Lag time = time from administration to appearance in blood.  
 Onset of activity = time from administration to blood level reaching minimal effective concentration (MEC).  
 Duration of action = time plasma concentration remains greater than MEC.  
 Time to peak = time from administration to  $C_{max}$ .

Figure I-1-4. Plot of Plasma Concentration Versus Time

### Bioavailability (f)

Measure of the fraction of a dose that reaches the systemic circulation. By definition, intravenous doses have 100% bioavailability,  $f = 1$ .




Figure I-1-5. Area Under the Curve for an IV Bolus and Extravascular Doses

### Abbreviations

- AUC: area under the curve
- PO: oral
- IV: intravenous bolus
- AUC<sub>IV</sub>: horizontally striped area
- AUC<sub>PO</sub>: vertically striped area

### First-Pass Effect

With oral administration, drugs are absorbed into the portal circulation and initially distributed to the liver. For some drugs, their rapid hepatic metabolism decreases bioavailability—the “first-pass” effect.

Examples:

- Lidocaine (IV vs. PO)
- Nitroglycerin (sublingual)





Figure I-1-6. Bioavailability and First-Pass Metabolism

## DISTRIBUTION

- The processes of distribution of a drug from the systemic circulation to organs and tissue.
- Conditions affecting distribution include:
  - Under normal conditions, protein-binding capacity is much larger than is drug concentration. Consequently, the free fraction is generally constant.
  - Many drugs bind to plasma proteins, including albumin, with an equilibrium between bound and free molecules (recall that only unbound drugs cross biomembranes).



- Competition between drugs for plasma protein-binding sites may increase the “free fraction,” possibly enhancing the effects of the drug displaced. Example: sulfonamides and bilirubin in a neonate

### Special Barriers to Distribution

- Placental—most small molecular weight drugs cross the placental barrier, although fetal blood levels are usually lower than maternal. Example: propylthiouracil (PTU) versus methimazole
- Blood–brain—permeable only to lipid-soluble drugs or those of very low molecular weight. Example: levodopa versus dopamine

### Apparent Volume of Distribution ( $V_d$ )

A kinetic parameter of a drug that correlates dose with plasma level at zero time.

$$V_d = \frac{\text{Dose}}{C^0} \quad \text{where } C^0 = [\text{plasma}] \text{ at zero time}$$

#### Bridge to Physiology

##### Approximate $V_d$ Values (weight 70 kg)

- plasma volume (3 L)
- blood volume (5 L)
- extracellular fluid (ECF 12–14 L)
- total body water (TBW 40–42 L)

- This relationship can be used for calculating  $V_d$  by using the *dose* only if one knows  $C^0$ .
- $V_d$  is low when a high percentage of a drug is bound to plasma proteins.
- $V_d$  is high when a high percentage of a drug is being sequestered in tissues. This raises the possibility of displacement by other agents; examples: verapamil and quinidine can displace digoxin from tissue-binding sites.
- $V_d$  is needed to calculate a loading dose in the clinical setting (*see* Pharmacokinetic Calculation section, Equation 4).

## Redistribution

In addition to crossing the blood–brain barrier (BBB), lipid-soluble drugs redistribute into fat tissues prior to elimination.

In the case of CNS drugs, the duration of action of an initial dose may depend more on the redistribution rate than on the half-life. With a second dose, the blood/fat is less; therefore, the rate of redistribution is less and the second dose has a longer duration of action.




Figure I-1-7. Redistribution

## BIOTRANSFORMATION

- The general principle of biotransformation is the metabolic conversion of drug molecules to more water-soluble metabolites that are more readily excreted.
- In many cases, metabolism of a drug results in its conversion to compounds that have little or no pharmacologic activity.
- In other cases, biotransformation of an active compound may lead to the formation of metabolites that also have pharmacologic actions.
- A few compounds (**prodrugs**) have no activity until they undergo metabolic activation.




Figure I-1-8. Biotransformation of Drugs

### Clinical Correlate

**Active Metabolites**  
 Biotransformation of the benzodiazepine diazepam results in formation of nordiazepam, a metabolite with sedative-hypnotic activity and a long duration of action.

## Biotransformation Classification

There are two broad types of biotransformation, called phase I and phase II.

### Phase I

- Definition: modification of the drug molecule via oxidation, reduction, or hydrolysis.
  - Microsomal metabolism
    - Cytochrome P450 isozymes*
      - These are major enzyme systems involved in phase I reactions. Localized in the smooth endoplasmic reticulum (microsomal fraction) of cells (especially liver, but including GI tract, lungs, and kidney).
      - P450s have an absolute requirement for molecular oxygen and NADPH.
      - Oxidations include hydroxylations and dealkylations.
      - Multiple CYP families differing by amino acid (AA) composition, by substrate specificity, and by sensitivity to inhibitors and to inducing agents.

### Clinical Correlate

#### Grapefruit Juice

Active components in grapefruit juice include furanocoumarins capable of inhibiting the metabolism of many drugs, including alprazolam, midazolam, atorvastatin, and cyclosporine. Such compounds may also enhance oral bioavailability by inhibiting drug transporters in the GI tract responsible for intestinal efflux of drugs.

Table I-1-2. Cytochrome P450 Isozymes

CYP450	Substrate Example	Inducers	Inhibitors	Genetic Polymorphisms
1A2	Theophylline Acetaminophen	Aromatic hydrocarbons (smoke) Cruciferous vegetables	Quinolones Macrolides	No
2C9	Phenytoin Warfarin	General inducers*	—	Yes
2D6	Many cardiovascular and CNS drugs	None known	Haloperidol Quinidine	Yes
3A4	60% of drugs in PDR	General inducers*	General inhibitors† Grapefruit juice	No

\* General inducers: anticonvulsants (barbiturates, phenytoin, carbamazepine), antibiotics (rifampin), chronic alcohol, glucocorticoids.

† General inhibitors: antiulcer medications (cimetidine, omeprazole), antibiotics (chloramphenicol, macrolides, ritonavir, ketoconazole), acute alcohol.

– Nonmicrosomal metabolism

*Hydrolysis*

- Phase I reaction involving addition of a water molecule with subsequent bond breakage
- Includes esterases and amidases
- Genetic polymorphism exists with pseudocholinesterases
- Example: local anesthetics

*Monoamine oxidases*

- Metabolism of endogenous amine neurotransmitters (dopamine, norepinephrine, and serotonin)
- Metabolism of exogenous compounds (tyramine)

*Alcohol metabolism*

- Alcohols are metabolized to aldehydes and then to acids by dehydrogenases (*see* CNS Pharmacology, section IV)
- Genetic polymorphisms exist

## Phase II

- Definition: Conjugation with endogenous compounds via the activity of transferases
- May follow phase I or occur directly
- Types of conjugation:

*Glucuronidation*

- Inducible
- May undergo enterohepatic cycling (Drug: Glucuronide → intestinal bacterial glucuronidases → free drug)
- Reduced activity in neonates
- Examples: morphine and chloramphenicol

*Acetylation*

- Genotypic variations (fast and slow metabolizers)
- Drug-induced SLE by slow acetylators with hydralazine > procainamide > isoniazid (INH)

*Glutathione (GSH) conjugation*

- Depletion of GSH in the liver is associated with acetaminophen hepatotoxicity

## ELIMINATION

Concerns the processes involved in the elimination of drugs from the body (and/or plasma) and their kinetic characteristics. The major modes of drug elimination are:

- Biotransformation to inactive metabolites
- Excretion via the kidney
- Excretion via other modes, including the bile duct, lungs, and sweat
- Definition: Time to eliminate 50% of a given amount (or to decrease plasma level to 50% of a former level) is called the elimination half-life ( $t_{1/2}$ ).

### Zero-Order Elimination Rate

- A constant amount of drug is eliminated per unit time; for example, if 80 mg is administered and 10 mg is eliminated every 4 h, the time course of drug elimination is:

4 h                      4 h                      4 h                      4 h  
 80 mg    →    70 mg    →    60 mg    →    50 mg    →    40 mg

- Rate of elimination is independent of plasma concentration (or amount in the body).
- Drugs with zero-order elimination have no fixed half-life ( $t_{1/2}$  is a variable).
- Drugs with zero-order elimination include ethanol (except low blood levels), phenytoin (high therapeutic doses), and salicylates (toxic doses).




Figure I-1-9a. Plots of Zero-Order Kinetics

### First-Order Elimination Rate

- A constant fraction of the drug is eliminated per unit time ( $t_{1/2}$  is a constant). Graphically, first-order elimination follows an exponential decay versus time.
- For example, if 80 mg of a drug is administered and its elimination half-life = 4 h, the time course of its elimination is:

4 h                      4 h                      4 h                      4 h  
 80 mg    →    40 mg    →    20 mg    →    10 mg    →    5 mg

- Rate of elimination is directly proportional to plasma level (or the amount present)—the higher the amount, the more rapid the elimination.
- Most drugs follow first-order elimination rates.
- $t_{1/2}$  is a constant




Figure I-1-9b. Plots of First-Order Kinetics

**In A Nutshell**

**Elimination Kinetics**

- Most drugs follow first order—rate falls as plasma level falls.
- Zero order is due to saturation of elimination mechanisms; e.g., drug-metabolizing reactions have reached  $V_{max}$
- Zero order—elimination rate is constant;  $t_{1/2}$  is a variable.
- First order—elimination rate is variable;  $t_{1/2}$  is a constant.

**Graphic Analysis**

Example of a graphic analysis of  $t_{1/2}$ :




Figure I-1-10. Plasma Decay Curve—First-Order Elimination

Figure I-1-10 shows a plasma decay curve of a drug with first-order elimination plotted on semilog graph paper. The elimination half-life ( $t_{1/2}$ ) and the theoretical plasma concentration at zero time ( $C^0$ ) can be estimated from the graphic relationship between plasma concentrations and time.  $C^0$  is estimated by extrapolation of the linear plasma decay curve to intercept with the vertical axis.



**Bridge to Renal Physiology**

Inulin clearance is used to estimate GFR because it is not reabsorbed or secreted. A normal GFR is close to 120 mL/min.

**Renal Elimination**

- Rate of elimination = glomerular filtration rate (GFR) + active secretion – reabsorption (active or passive).
- Filtration is a nonsaturable linear function. Ionized and nonionized forms of drugs are filtered, but protein-bound drug molecules are not.
- Clearance (Cl):
  - Definition: volume of blood cleared of drug per unit of time
  - Cl is constant in first-order kinetics
  - $Cl = GFR$  when there is no reabsorption or secretion and no plasma protein binding
  - Protein-bound drug is not cleared;  $Cl = \text{free fraction} \times GFR$

**STEADY STATE**

- Steady state is reached either when **rate in = rate out** or when values associated with a dosing interval are the same as those in the succeeding interval.

**Plateau Principle**

The time to reach steady state is dependent only on the elimination half-life of a drug and is independent of dose size and frequency of administration.

**Note**

$$MD = \frac{C^{ss} \times Cl \times \tau}{f}$$

Figure I-1-11 shows plasma levels (solid lines) achieved following the IV bolus administration of 100 units of a drug at intervals equivalent to every **half-life  $t_{1/2} = 4$  h ( $\tau$ )**. With such intermittent dosing, plasma levels oscillate through peaks and troughs, with averages shown in the diagram by the dashed line.

**Note**

See legend on page 17.

**Classic Clues**


**Time and Steady State**

50% = 1  $\times$  half-life

90% = 3.3  $\times$  half-life

95% = 4–5  $\times$  half-life

“100”% = >7  $\times$  half-life



**Figure I-1-11. Oscillations in Plasma Levels Following IV Bolus Administration at Intervals Equal to Drug Half-Life**

**Note:** Although it takes >7  $t_{1/2}$  to reach **mathematical** steady state, by convention **clinical** steady state is accepted to be reached at 4–5  $t_{1/2}$ .

## Rate of Infusion

Figure I-1-12 shows the increases in plasma levels of the same drug infused at five different rates. Regardless of the rate of infusion, it takes the same amount of time to reach steady state.




Figure I-1-12. Effect of Rate of Infusion on Plasma Level


Rate of infusion ( $k_0$ ) does determine plasma level at steady state. If the rate of infusion is doubled, then the plasma level of the drug at steady state is doubled. A similar relationship can exist for other forms of drug administration (e.g., per oral)—doubling oral doses can double the average plasma levels of a drug. Plotting dose against plasma concentration yields a straight line (linear kinetics).

### Note

- Remember that dose and plasma concentration ( $C^{SS}$ ) are directly proportional.

### Effect of Loading Dose

- It takes 4–5 half-lives to achieve steady state.
- In some situations, it may be necessary to give a higher dose (loading dose) to more rapidly achieve effective blood levels ( $C_p$ ).



**Figure I-1-13. Effect of a Loading Dose on the Time Required to Achieve the Minimal Effective Plasma Concentration**

- Such loading doses are often one time only and (as shown in Figure I-1-13) are estimated to put into the body the amount of drug that should be there at a steady state.
- For the exam, if doses are to be administered at each half-life of the drug and the minimum effective concentration is equivalent to  $C_{min}^{SS}$ , then the loading dose is twice the amount of the dose used for maintenance (assuming normal clearance and same bioavailability for maintenance doses). For any other interval of dosing, Equation 2 (below) is used.

## IMPORTANT PHARMACOKINETICS CALCULATIONS

The following five relationships are important for calculations:

### Single-Dose Equations

- (1) Volume of distribution ( $V_d$ )

$$V_d = \frac{D}{C_0}$$

- (2) Half-life ( $t_{1/2}$ )

$$t_{1/2} = 0.7 \times \frac{V_d}{Cl}$$

### Multiple Doses or Infusion Rate Equations

- (3) Infusion rate ( $k_0$ )

$$k_0 = Cl \times C^{ss}$$

- (4) Loading dose (LD)

$$LD = \frac{V_d \times C_p}{f}$$

- (5) Maintenance dose (MD)

$$MD = \frac{Cl \times C^{ss} \times \tau}{f}$$

### Note

$$LD = \frac{V_d \times C_p}{f}$$

### Legend

$C^0$  = conc. at time zero

Cl = clearance

$C_p$  = conc. in plasma

$C^{ss}$  = steady state conc.

D = dose

f = bioavailability

$k_0$  = infusion rate

LD = loading dose

MD = maintenance dose

$\tau$  = dosing interval

$V_d$  = volume of distribution

## Chapter Summary

- \* The pharmacokinetic characteristics of a drug are dependent upon the processes of absorption, distribution, metabolism, and excretion. An important element concerning drug biodistribution is permeation, which is the ability to cross membranes, cellular and otherwise.
- \* A drug's ability to permeate is dependent on its solubility, the concentration gradient, and the available surface area, which is influenced by the degree of vascularity. Ionization affects permeation because unionized molecules are minimally water soluble but do cross biomembranes, a feat beyond the capacity of ionized molecules. Figure I-1-2 illustrates the principles associated with ionization; Table I-1-1 summarizes the three basic modes of transport across a membrane: passive, facilitated, and active.
- \* Absorption concerns the processes of entry into the systemic circulation. Except for the intravascular route, some absorptive process is always involved. These have the same determinants as those of permeation. Because absorption may not be 100% efficient, less than the entire dose administered may get into the circulation.
- \* Any orally administered hydrophilic drug will be absorbed first into the portal vein and sent directly to the liver, where it may be partially deactivated. This is the first-pass effect.

(Continued)

**Chapter Summary (cont'd)**

- \* The distribution of a drug into the various compartments of the body is dependent upon its permeation properties and its tendency to bind to plasma proteins. The placental and blood-brain barriers are of particular importance in considering distribution. The  $V_d$  is a kinetic parameter that correlates the dose given to the plasma level obtained: the greater the  $V_d$  value, the less the plasma concentration.
- \* As well as having the ability to cross the blood-brain barrier, lipophilic drugs have a tendency to be deposited in fat tissue. As blood concentrations fall, some of this stored drug is released. This is called redistribution. Because with each administration more lipophilic drug is absorbed into the fat, the duration of action of such a drug increases with the number of doses until the lipid stores are saturated.
- \* Biotransformation is the metabolic conversion of drugs, generally to less active compounds but sometimes to iso-active or more active forms. Phase I biotransformation occurs via oxidation, reduction, or hydrolysis. Phase II metabolism occurs via conjugation.
- \* The cytochrome P-450 isozymes are a family of microsomal enzymes that collectively have the capacity to transform thousands of different molecules. The transformations include hydroxylations and dealkylations, as well as the promotion of oxidation/reduction reactions. These enzymes have an absolute requirement for NADPH and  $O_2$ . The various isozymes have different substrate and inhibitor specificities.
- \* Other enzymes involved in phase I reactions are hydrolases (e.g., esterases and amidases) and the nonmicrosomal oxidases (e.g., monoamine oxidase and alcohol and aldehyde dehydrogenase).
- \* Phase II reactions involve conjugation, sometimes after a phase I hydroxylation. The conjugation may be glucuronidation, acetylation, sulfation, or addition of glutathione.
- \* Modes of drug elimination are biotransformation, renal excretion, and excretion by other routes (e.g., bile, sweat, lungs, etc.). Most drugs follow first-order elimination rates. Figures I-1-9a and I-1-9b compare zero- and first-order elimination, and Figure I-1-10 demonstrates how the  $t_{1/2}$  and the theoretical zero time plasma concentration ( $C^0$ ) can be graphically determined. An important relationship is  $\text{dose} = V_d \times C^0$ .
- \* Renal clearance ( $Cl_R$ ) represents the volume of blood cleared by the kidney per unit time and is a constant for drugs with first-order elimination kinetics. Total body clearance equals renal plus nonrenal clearance. An important relationship is  $Cl = k \times V_d$ .
- \* A steady state is achieved when the rate coming in equals the rate going out. The time to reach a steady state is dependent only on the elimination half-life. It is independent of dose and frequency of administration or rate of infusion (see Figures I-1-11, -12, and -13).
- \* Other equations describing relationships important for calculation are those used to determine the loading dose, infusion rate, and maintenance dose.

# Pharmacodynamics

## DEFINITIONS

- Pharmacodynamics relates to drugs binding to receptors and their effects.
- **Agonist:** A drug is called an agonist when binding to the receptor results in a response.
- **Antagonist:** A drug is called an antagonist when binding to the receptor is *not* associated with a response. The drug has an effect only by preventing an agonist from binding to the receptor.
- **Affinity:** ability of drug to bind to receptor, shown by the proximity of the curve to the y axis (if the curves are parallel); the nearer the y axis, the greater the affinity
- **Potency:** shows relative doses of two or more agonists to produce the same magnitude of effect, again shown by the proximity of the respective curves to the y axis (if the curves do not cross)
- **Efficacy:** a measure of how well a drug produces a response (effectiveness), shown by the maximal height reached by the curve

## GRADED (QUANTITATIVE) DOSE-RESPONSE (D-R) CURVES

Plots of dose (or log dose) versus response for drugs (agonists) that activate receptors can reveal information about affinity, potency, and efficacy of these agonists.

### Parallel and Nonparallel D-R Curves




Figure I-2-1. Comparison of D-R Curves for Two Drugs Acting on the Same (*left panel*) and on Different (*right panel*) Receptors

## Bridge to Biochemistry

### Definitions

**Affinity:** how well a drug and a receptor recognize each other. Affinity is inversely related to the  $K_d$  of the drug. Notice the analogy to the  $K_m$  value used in enzyme kinetic studies.

**Potency:** the quantity of drug required to achieve a desired effect. In D-R measurements, the chosen effect is usually 50% of the maximal effect, but clinically, *any* size response can be sought.

**Efficacy:** the maximal effect an agonist can achieve at the highest practical concentration. Notice the analogy with the  $V_{max}$  used in enzyme kinetic studies.

It may be seen from the log dose-response curves in Figure I-2-1 that:

1. When two drugs interact with the same receptor (same pharmacologic mechanism), the D-R curves will have parallel slopes. Drugs A and B have the same mechanism; drugs X and Y do not.
2. Affinity can be compared only when two drugs bind to the same receptor. Drug A has a greater affinity than drug B.
3. In terms of potency, drug A has greater potency than drug B, and X is more potent than Y.
4. In terms of efficacy, drugs A and B are equivalent. Drug X has greater efficacy than drug Y.

### Full and Partial Agonists

- Full agonists produce a maximal response—they have maximal efficacy.
- Partial agonists are incapable of eliciting a maximal response and are less effective than full agonists.
- In Figure I-2-2, drug B is a full agonist, and drugs A and C are partial agonists.




Figure I-2-2. Efficacy and Potency of Full and Partial Agonists

- Drug A is more potent than drug C, and drug B is more potent than drug C. However, no general comparisons can be made between drugs A and B in terms of potency because the former is a partial agonist and the latter is a full agonist. At low responses, A is more potent than B, but at high responses, the reverse is true.

## Duality of Partial Agonists

- In Figure I-2-3, the lower curve represents effects of a partial agonist when used alone—its *ceiling effect* = 50% of maximal in this example.




Figure I-2-3. Duality of Partial Agonists

- The upper curve shows the effect of increasing doses of the partial agonist on the maximal response (100%) achieved in the presence of or by pretreatment with a full agonist.
- As the partial agonist displaces the full agonist from the receptor, the response is reduced—the partial agonist is acting as an **antagonist**.

## Antagonism and Potentiation

- Graded dose-response curves also provide information about antagonists—drugs that interact with receptors to interfere with their activation by agonists.




Figure I-2-4. D-R Curves of Antagonists and Potentiators

- Pharmacologic antagonism (same receptor)
  - Competitive antagonists:
    - Cause a parallel shift to the right in the D-R curve for agonists
    - Can be reversed by  $\uparrow$  the dose of the agonist drug
    - Appears to  $\downarrow$  the potency of the agonist

### Bridge to Biochemistry

#### Parallels between Receptor Antagonists and Enzyme Inhibitors

Competitive antagonists are analogous to competitive inhibitors; they decrease affinity ( $\uparrow K_m$ ) but not maximal response ( $V_{max}$  remains the same).

Noncompetitive antagonists decrease  $V_{max}$  but do not change the  $K_m$ .



- Noncompetitive antagonists:
  - Cause a nonparallel shift to the right
  - Can be only partially reversed by ↑ the dose of the agonist
  - Appear to ↓ the efficacy of the agonist
- Physiologic antagonism (different receptor)
  - Two agonists with opposing action antagonize each other
  - Example: a vasoconstrictor with a vasodilator
- Chemical antagonism:
  - Formation of a complex between effector drug and another compound
  - Example: protamine binds to heparin to reverse its actions
- Potentiation
  - Causes a parallel shift to the left to the D-R curve
  - Appears to ↑ the potency of the agonist

### QUANTAL (CUMULATIVE) D-R CURVES

- These curves plot the percentage of a population responding to a specified drug effect versus dose or log dose. They permit estimations of the median effective dose, or effective dose in 50% of a population—ED50.
- Quantal curves can reveal the range of intersubject variability in drug response. Steep D-R curves reflect little variability; flat D-R curves indicate great variability in patient sensitivity to the effects of a drug.

### Toxicity and the Therapeutic Index (TI)

- Comparisons between ED50 and TD50 values permit evaluation of the relative safety of a drug (the therapeutic index), as would comparison between ED50 and the lethal median dose (LD50) if the latter is known.

$$TI = \frac{TD50}{ED50} \text{ or } \frac{LD50}{ED50}$$




Figure I-2-5. Quantal D-R Curves of Therapeutic and Toxic Effects of a Drug

- As shown in Figure I-2-5, these D-R curves can also be used to show the relationship between dose and toxic effects of a drug. The median toxic dose of a drug (TD<sub>50</sub>) is the dose that causes toxicity in 50% of a population.
- From the data shown,  $TI = 10/2 = 5$
- Such indices are of most value when toxicity represents an extension of the pharmacologic actions of a drug. They do not predict idiosyncratic reactions or drug hypersensitivity.

## SIGNALING MECHANISMS: TYPES OF DRUG-RESPONSIVE SIGNALING MECHANISMS

- Binding of an agonist drug to its receptor activates an effector or signaling mechanism.
- Several different types of drug-responsive signaling mechanisms are known.

### Intracellular Receptors

- These include receptors for steroids. Binding of hormones or drugs to such receptors releases regulatory proteins that permit activation and in some cases dimerization of the hormone-receptor complex. Such complexes translocate to the nucleus, where they interact with response elements in spacer DNA. This interaction leads to changes in gene expression. For example, drugs interacting with glucocorticoid receptors lead to gene expression of proteins that inhibit the production of inflammatory mediators.
- Other examples include intracellular receptors for thyroid hormones, gonadal steroids, and vitamin D.
- Pharmacologic responses elicited via modification of gene expression are usually slower in onset but longer in duration than many other drugs.

### Membrane Receptors Directly Coupled to Ion Channels

- Many drugs act by mimicking or antagonizing the actions of endogenous ligands that regulate flow of ions through excitable membranes via their activation of receptors that are directly coupled (no second messengers) to ion channels.
- For example, the nicotinic receptor for ACh (present in autonomic nervous system [ANS] ganglia, the skeletal myoneural junction, and the central nervous system [CNS]) is coupled to a  $Na^+/K^+$  ion channel. The receptor is a target for many drugs, including nicotine, choline esters, ganglion blockers, and skeletal muscle relaxants.
- Similarly, the GABA<sub>A</sub> receptor in the CNS, which is coupled to a chloride ion channel, can be modulated by anticonvulsants, benzodiazepines, and barbiturates.

### Receptors Linked Via Coupling Proteins to Intracellular Effectors

- Many receptor systems are coupled via GTP-binding proteins (G-proteins) to adenylyl cyclase, the enzyme that converts ATP to cAMP, a second messenger that promotes protein phosphorylation by activating protein kinase A. These receptors are typically “serpentine,” with seven transmembrane spanning domains, the third of which is coupled to the G-protein effector mechanism.
- Protein kinase A serves to phosphorylate a set of tissue-specific substrate enzymes or transcription factors (CREB), thereby affecting their activity.

**In A Nutshell**

**Key ANS Receptors**

$M_1, M_3, \alpha_1$ :  $G_q$  activation of phospholipase C

$M_2, \alpha_2, D_2$ :  $G_i$  inhibition of adenylyl cyclase

$\beta_1, \beta_2, D_1$ :  $G_s$  activation of adenylyl cyclase

**$G_s$  Proteins**

- Binding of agonists to receptors linked to  $G_s$  proteins increases cAMP production.
- Such receptors include those for catecholamines (beta), dopamine ( $D_1$ ), glucagon, histamine ( $H_2$ ), prostacyclin, and some serotonin subtypes.

**$G_i$  Proteins**

- Binding of agonists to receptors linked to  $G_i$  proteins decreases cAMP production.
- Such receptors include adrenoreceptors ( $\alpha_2$ ), ACh ( $M_2$ ), dopamine ( $D_2$  subtypes), and several opioid and serotonin subtypes.

**$G_q$  Proteins**

- Other receptor systems are coupled via GTP-binding proteins ( $G_q$ ), which activate phospholipase C. Activation of this enzyme releases the second messengers inositol triphosphate ( $IP_3$ ) and diacylglycerol (DAG) from the membrane phospholipid phosphatidylinositol biphosphate ( $PIP_2$ ). The  $IP_3$  induces release of  $Ca^{2+}$  from the sarcoplasmic reticulum (SR), which, together with DAG, activates protein kinase C. The protein kinase C serves then to phosphorylate a set of tissue-specific substrate enzymes, usually not phosphorylated by protein kinase A, and thereby affects their activity.
- These signaling mechanisms are invoked following activation of receptors for ACh ( $M_1$  and  $M_3$ ), norepinephrine ( $\alpha_1$ ), angiotensin II, and several serotonin subtypes.




Figure I-2-6. Receptors Using Cyclic-AMP and  $IP_3$ , DAG,  $Ca^{2+}$  as Second Messengers

## Cyclic GMP and Nitric Oxide Signaling

- cGMP is a second messenger in vascular smooth muscle that facilitates dephosphorylation of myosin light chains, preventing their interaction with actin and thus causing vasodilation.
- Nitric oxide (NO) is synthesized in endothelial cells and diffuses into smooth muscle.
- NO activates guanylyl cyclase, thus increasing cGMP in smooth muscle.
- Vasodilators ↑ synthesis of NO by endothelial cells.

## Receptors That Function as Enzymes or Transporters

- There are multiple examples of drug action that depend on enzyme inhibition, including inhibitors of acetylcholinesterase, angiotensin-converting enzyme, aspartate protease, carbonic anhydrase, cyclooxygenases, dihydrofolate reductase, DNA/RNA polymerases, monoamine oxidases, Na/K-ATPase, neuraminidase, and reverse transcriptase.
- Examples of drug action on transporter systems include the inhibitors of reuptake of several neurotransmitters, including dopamine, GABA, norepinephrine, and serotonin.

## Receptors That Function as Transmembrane Enzymes

- These receptors mediate the first steps in signaling by insulin and growth factors, including epidermal growth factor (EGF) and platelet-derived growth factor (PDGF). They are membrane-spanning macromolecules with recognition sites for the binding of insulin and growth factors located externally and a cytoplasmic domain that usually functions as a tyrosine kinase. Binding of the ligand causes conformational changes (e.g., dimerization) so that the tyrosine kinase domains become activated, ultimately leading to phosphorylation of tissue-specific substrate proteins.
- Guanyl cyclase-associated receptors: stimulation of receptors to atrial natriuretic peptide activates the guanyl cyclase and ↑ cyclic GMP (cGMP)

## Receptors for Cytokines

- These include the receptors for erythropoietin, somatotropin, and interferons.
- Their receptors are membrane spanning and on activation can activate a distinctive set of cytoplasmic tyrosine kinases (Janus kinases [JAKs]).
- JAKs phosphorylate signal transducers and activators of transcription (STAT) molecules.
- STATs dimerize and then dissociate, cross the nuclear membrane, and modulate gene transcription.

## Bridge to Biochemistry

See Chapter 9 of the Biochemistry Lecture Notes for additional discussion of signal transduction.

## Clinical Correlate

Drugs acting via NO include nitrates (e.g., nitroglycerin) and M-receptor agonists (e.g., bethanechol). Endogenous compounds acting via NO include bradykinin and histamine.

## DRUG DEVELOPMENT AND TESTING

### The Food and Drug Administration (FDA)

The FDA regulates both the efficacy and safety of drugs but not of foods, nutritional supplements, and herbal remedies.

Table I-2-1. Drug Development and Testing

Preclinical	Phase 1	Phase 2	Phase 3	Phase 4
Two different animal species	~50 healthy volunteers	~200 patients	~2,000 patients	Post-marketing surveillance (after FDA approval)
Safety and biologic activity	Safety and dosage	Evaluate effectiveness	Confirm effectiveness, common side-effects	Common as well as rare side effects

### Teratogenicity

- The FDA has classified drugs into five categories (A, B, C, D, and X).
- Class A has no risks, and Class X designates absolute contraindication.
- It is based on animal studies and, when available, human studies.
- In Class D, benefits outweigh the risk.

Table I-2-2. FDA Classification of Drugs and Pregnancy

Category	Risk	
	Animals	Humans
A	-	-
B	+/-	-/o
C	+/o	o
D	+	+
X	+	+

- = Studies have proven absence of teratogenicity;  
 o = no studies available;  
 + = studies have proven teratogenicity

### Chapter Summary

- \* Plots of dose or log dose against response to a drug (agonist) can be used to assess the drug's affinity to a receptor, its potency (the amount of drug required to achieve half its maximal effect), and its efficacy (the maximal effect). Full agonists achieve full efficacy; partial agonists do not. Therefore, when a partial agonist is added to a system in which a full agonist is acting at its maximal efficacy, the partial agonist acts as a competitive inhibitor, as if it were an antagonist. These effects can be studied graphically.
- \* Antagonists are compounds that inhibit the activity of an agonist but have no effect of their own. Generally, antagonists act competitively by sharing a binding site on the receptor, but some act noncompetitively. Whether an antagonist acts competitively or noncompetitively can also be determined graphically.
- \* Antagonism may be pharmacologic (shared receptor), physiologic (acting on different systems having opposing physiologic responses), or chemical.
- \* Some effector molecules potentiate (i.e., enhance) the effect of an agonist.
- \* Quantal curves are plots of the percentage of a population responding to a specific drug versus the concentration (or log concentration) of that drug. They are used to gauge the median effective pharmacological dose (ED50) or the median toxic dose (TD50). These values can be used to evaluate the relative safety of a drug (the therapeutic index).
- \* Drugs may act on intracellular receptors, membrane receptors directly coupled to ion channels, receptors linked via coupling proteins to intracellular effectors, receptors influencing cGMP and nitric oxide signaling, receptors that function as enzymes or transporters, receptors that function as transmembrane enzymes, or receptors for cytokines.
- \* The FDA regulates the efficacy and safety of drugs but not of foods, herbs, or nutritional supplements. Before being approved by the FDA, a drug must first undergo preclinical animal studies and then phase 1, 2, 3, and 4 clinical studies. FDA also classifies drugs and their relative risks of teratogenicity during pregnancy.



# Practice Questions

---


## Review Questions

1. A patient was given a 200 mg dose of a drug IV, and 100 mg was eliminated during the first two hours. If the drug follows first-order elimination kinetics, how much of the drug will remain 6 hours after its administration?
  - A. None
  - B. 25 mg
  - C. 50 mg
  - D. 75 mg
  - E. 100 mg
2. Drugs that are administered IV are
  - A. Rapidly Absorbed
  - B. Subject to first-pass metabolism
  - C. 100% bioavailable
  - D. Rapidly excreted by the kidneys
  - E. Rapidly metabolized by the liver
3. Drugs that are highly bound to albumin:
  - A. Effectively cross the BBB
  - B. Are easily filtered at the glomerulus
  - C. Have a large  $V_d$
  - D. Often contain quaternary nitrogens
  - E. Can undergo competition with other drugs for albumin binding sites
4. Most drugs gain entry to cells by:
  - A. Passive diffusion with zero-order kinetics
  - B. Passive diffusion with first-order kinetics
  - C. Active transport with zero-order kinetics
  - D. Active transport with first-order kinetics
  - E. Passive diffusion through membrane pores



5. A subject in whom the renal clearance of inulin is 120 mL/min is given a drug, the clearance of which is found to be 18 mL/min. If the drug is 40% plasma protein bound, how much filtered drug must be reabsorbed in the renal tubules?
- A. None
  - B. 18 mL/min
  - C. 36 mL/min
  - D. 54 mL/min
  - E. 72 mL/min
6. If a drug is known to be distributed into total body water, what dose (mg) is needed to obtain an initial plasma level of 5 mg/L in a patient weighing 70 kg?
- A. 210
  - B. 150
  - C. 110
  - D. 50
  - E. 35
7. Which of the following is a phase II drug metabolism reaction associated with a genetic polymorphism?
- A. Acetylation
  - B. Glucuronidation
  - C. Oxidation
  - D. Reduction
  - E. Glutathione conjugation
8. A woman is taking oral contraceptives (OCs). Which of the following drugs is unlikely to reduce the effectiveness of the OCs?
- A. Carbamazepine
  - B. Phenytoin
  - C. Ketoconazole
  - D. Phenobarbital
  - E. Rifampin

9. The data presented in the figure below show that:



- A. Drugs A and B have equal efficacy  
B. Drug B and C have equal efficacy  
C. Drug B is a partial agonist  
D. Drugs A and C have the same affinity and efficacy  
E. Drugs A and B have equal potency
10. A 500-mg dose of a drug has therapeutic efficacy for 6 h. If the half-life of the drug is 8 h, for how long would a 1-g dose be effective?
- A. 8 h  
B. 12 h  
C. 14 h  
D. 16 h  
E. 24 h
11. A drug achieves a plasma level of 16 mg/L shortly after the administration of the first oral dose. If the half-life and the dosing interval are both 6 hours, what is the approximate plasma level shortly before the administration of the 5<sup>th</sup> dose?
- A. 15 mg/L  
B. 24 mg/L  
C. 28 mg/L  
D. 30 mg/L  
E. 31 mg/L

12. Which statement is accurate for the drug shown in the example below?
- $$100 \text{ mg}^{2\text{hr}} \rightarrow 50 \text{ mg}^{2\text{hr}} \rightarrow 25 \text{ mg}^{2\text{hr}} \rightarrow 12.5 \text{ mg}$$
- A. The rate of elimination is constant
  - B. The elimination half-life varies with the dose
  - C. The volume of distribution varies with the dose
  - D. The clearance varies with the dose
  - E. The rate of elimination varies directly with the dose
13. Normally, acetaminophen has a  $V_d = 70\text{L}$  and  $Cl = 350 \text{ mL/min}$ . If acetaminophen was administered to a patient with 50% renal function, what parameter would differ from normal?
- A. Loading dose would be higher
  - B. Maintenance dose would be higher
  - C.  $t_{1/2}$  would be shorter
  - D.  $V_d$  would be 35L
  - E.  $Cl$  would be 700 mL/min
14. Pharmacokinetic characteristics of propranolol include  $V_d = 300 \text{ L}/70 \text{ kg}$ ,  $Cl = 700 \text{ mL/min}$ , oral bioavailability  $f = 0.25$ . What is the dose needed to achieve a plasma level equivalent to a steady-state level of 20 g/L?
- A. 4 mg
  - B. 8 mg
  - C. 12 mg
  - D. 24 mg
  - E. 48 mg
15. With IV infusion, a drug reaches 50% of its final steady state in 6 hours. The elimination half-life of the drug must be approximately:
- A. 2 h
  - B. 6 h
  - C. 12 h
  - D. 24 h
  - E. 30 h
16. At 6 h after IV administration of bolus dose, the plasma level of a drug is 5 mg/L. If the  $V_d = 10 \text{ L}$  and the elimination half-life = 3 h, what was the dose administered?
- A. 100 mg
  - B. 150 mg
  - C. 180 mg
  - D. 200 mg
  - E. 540 mg

17. An IV infusion of a drug is started 400 mg/h. If  $Cl = 50$  L/h, what is the anticipated plasma level at steady state?
- 2 mg/L
  - 4 mg/L
  - 8 mg/L
  - 16 mg/L
  - 32 mg/L

## Answers

- Answer B.** One half of the dose is eliminated in the first two hours so its elimination half-life equals two hours. With the passage of each half-life the amount in the body (or in the blood) will decrease to 50% of a former level. Thus, at 6 hours after administration, three half-lives have passed: 1) 200 mg to 100 mg, 2) 100 mg to 50 mg, and 3) 50 mg to 25 mg.
- Answer C.** By definition, IV administration does not involve absorption because there is no movement from the site of administration into the blood. The IV route avoids first-pass metabolism which is common with orally administered drugs. First-pass greatly reduces the bioavailability of many drugs. Drugs given IV have 100% bioavailability ( $f = 1$ ) since the entire dose is in the systemic circulation. No conclusions can be drawn about renal or hepatic elimination of a drug knowing only that it was administered IV.
- Answer E.** Since most drugs are lipid-soluble they will need a carrier in the blood, most commonly albumin. Drugs bound to albumin do not get filtered at the glomerulus or cross the blood-brain barrier. Binding to plasma proteins keeps drugs in the plasma resulting in a lower  $V_d$ . Highly protein bound drugs are good candidates for interactions with other drugs that are also highly bound (e.g., warfarin plus sulfonamides).
- Answer B.** The permeation of most drugs through cellular membranes is by the process of passive diffusion, a nonsaturable process that follows first-order kinetics. Concentration gradient and lipid solubility are important determinants for the rate of diffusion. Only a few drugs are substrates for active transport processes such as active tubular secretion (e.g., penicillins) or penetrate membranes via aqueous pores (ethanol).
- Answer D.** The formula to use is  $Cl = ff \times GFR$ . The drug is 40% protein bound so the  $ff = 60\%$ .  $120 \text{ mL/min} \times 60\% = 72 \text{ mL/min}$  theoretical clearance of the drug. Since only 18 mL/min was actually cleared, there must have been tubular reabsorption of the drug.  $72 - 18 = 54 \text{ mL/min}$  of reabsorbed drug.
- Answer A.** This is a "loading dose" question. The equation for loading dose or the volume of distribution equation can be used ( $LD = V_d \times C_p$ ). Since the patient weighs 70 kg and 60% of body weight is water, he has 42 L ( $70 \text{ L} \times 60\%$ ) of total body water.  $LD = 42 \text{ L} \times 5 \text{ mg/L} = 210 \text{ mg}$ .

7. **Answer A.** Phase II drug metabolism involves the transfer of chemical groupings (e.g. acetyl, glucuronide, glutathione) to drugs or their metabolites via conjugation reactions involving transferase enzymes. Acetylation reactions are associated with a genetic polymorphism (slow acetylator). These individuals are slow to metabolize drugs via acetylation and are particularly susceptible to drug-induced SLE when taking hydralazine, procainamide, or isoniazid. Both oxidation and reduction are phase I metabolism reactions.
8. **Answer C.** Azole antifungals (e.g. ketoconazole) are inhibitors of cytochrome P450 enzymes, especially CYP3A4, the most abundant isozyme form in the human liver. The 3A4 isozyme metabolizes a wide range of drugs. Ketoconazole would actually raise the plasma levels of oral contraceptives increasing the risk of side effects but it would not reduce their effectiveness. All other drugs listed are P450 inducers. As such, they would tend to lower plasma levels and decrease effectiveness of oral contraceptives.
9. **Answer B.** The typical log dose response figure with the parallel nature of the curves suggests that the three drugs are interacting with the same receptor system. Drugs A and B are full agonists because they achieve the maximal response. They have the same efficacy. Drug A is more potent than drugs B or C. Drug B is more potent than drug C. Drug C is a partial agonist with less efficacy than the full agonists.
10. **Answer C.** The fact that the drug has therapeutic efficacy for 6 h has no direct relationship to its half-life—it simply means that the drug is above its minimal effective concentration for 6 h. Doubling the dose (to 1 g) means that the drug level will be above the minimum for a longer period. Because the elimination half-life is 8 h, 500 mg of the drug will remain in the body 8 h after a dose of 1 g. Thus, the total duration of effectiveness must be  $8 + 6 = 14$  h.
11. **Answer A.** The key word in this question is *before* the 5th dose. Immediately “after” the 5th dose, the plasma level should be approximately 30 mg/L, but just before it would be close to half of that level.
12. **Answer E.** In first-order kinetics, the elimination rate of a drug is directly proportional to its plasma concentration, which in turn is proportional to the dose. Drugs that follow first-order elimination have a constant elimination half-life similar to the example given in the question. Likewise, clearance and volume of distribution are pharmacokinetic characteristics of a drug that do not routinely change with dose, although they may vary in terms of disease or dysfunction.
13. **Answer B.** The patient in the question has renal dysfunction which would require a dosage adjustment for medications like acetaminophen given by maintenance dose but not loading dose. The maintenance dose equation ( $MD = Cl \times C^{ss} \times \tau$ ) factors in renal clearance while the loading dose equation does not. The  $t_{1/2}$  of acetaminophen would be increased in this patient due to the decrease in clearance, but the  $V_d$  would be unaffected.
14. **Answer D.**

$$\text{Loading dose} = V_d \times \frac{C_p}{f} \qquad LD = 300L \times 20 \mu\text{g/L} \div 0.25$$

$$= 6000 \mu\text{g}/0.25 \qquad = 24,000 \mu\text{g} \text{ or } 24 \text{ mg}$$
15. **Answer B.** The rules for time to steady-state are that it takes 4-5  $t_{1/2}$  to reach clinical steady-state. It also takes one  $t_{1/2}$  to get half way to steady-state. Since the drug got 50% of the way to steady-state in 6 hours, its  $t_{1/2}$  must be 6 hours.

16. **Answer A.** At 6 h after IV injection (which corresponds to two half-lives of the drug), the plasma level is 5 mg/L. Extrapolating back to zero time, “doubling” plasma level for each half-life results in an initial plasma level at zero time ( $C_0$ ) =  $5 \times 2 \times 2$  mg/L = 20 mg/L.

17. **Answer C.**  $MD = Cl \times C^{ss} \times \tau$

Since the drug was given by constant IV infusion there is no need to consider the dosing interval ( $\tau$ ). Therefore,  $400 \text{ mg/h} = 50 \text{ L/h} \times C^{ss}$

$$400 \text{ mg/h} \div 50 \text{ L/h} = 8 \text{ mg/L}$$



SECTION II

**Autonomic  
Pharmacology**





# The Autonomic Nervous System (ANS)

---

## ANATOMY OF THE ANS

The ANS is the major involuntary portion of the nervous system and is responsible for automatic, unconscious bodily functions, such as control of heart rate and blood pressure and both gastrointestinal and genitourinary functions. The ANS is divided into two major subcategories: the parasympathetic autonomic nervous system (PANS) and the sympathetic autonomic nervous system (SANS).

### Location of ANS Ganglia

Both the PANS and SANS have relay stations, or ganglia, between the CNS and the end organ, but the somatic system does not. An important anatomic difference between the SANS and PANS is that the ganglia of the former lie in two paraventral chains adjacent to the vertebral column, whereas most of the ganglia of the PANS system are located in the organs innervated. Figure II-1-1 highlights the major features of the ANS and the somatic systems and also shows the location of the major receptor types. These are:

- $N_N$ —Nicotinic receptors are located on cell bodies in ganglia of both PANS and SANS and in the adrenal medulla.
- $N_M$ —Nicotinic receptors are located on the skeletal muscle motor end plate innervated by somatic motor nerves.
- $M_{1-5}$ —Muscarinic receptors are located on all organs and tissues innervated by post-ganglionic nerves of the PANS and on thermoregulatory sweat glands innervated by the SANS.




Figure II-1-1. Anatomy of the Autonomic Nervous System

## Neurotransmitters

- Acetylcholine (ACh) is the neurotransmitter at both nicotinic and muscarinic receptors in tissues that are innervated. Note that all direct transmission from the CNS (preganglionic and motor) uses ACh, but postganglionic transmission in the SANS system may use one of the organ-specific transmitters described below.
- Norepinephrine (NE) is the neurotransmitter at most adrenoceptors in organs, as well as in cardiac and smooth muscle.
- Dopamine (DA) activates  $D_1$  receptors, causing vasodilation in renal and mesenteric vascular beds.
- Epinephrine (E, from adrenal medulla) activates most adrenoceptors and is transported in the blood.

## BLOOD PRESSURE CONTROL MECHANISMS

### Autonomic Feedback Loop


- Blood pressure is the product of total peripheral resistance (TPR) and cardiac output (CO).
- Both branches of the ANS are involved in the autonomic (or neural) control of blood pressure via feedback mechanisms.
- Changes in mean blood pressure are detected by baroreceptors, which relay information to the cardiovascular centers in the brainstem controlling PANS and SANS outflow. For

### Bridge to Physiology

For a more detailed discussion, see Section II, Chapter 2, in Physiology.

example, an increase in mean blood pressure elicits baroreceptor discharge, resulting in increased PANS activity, leading to bradycardia and decreased SANS activity, which leads, in turn, to decreased heart rate, force of contraction, and vasoconstriction. The resulting decreases in cardiac output and total peripheral resistance contribute to restoration of mean blood pressure toward its normal level.

- Conversely, decreases in blood pressure elicit ANS neural feedback involving decreased PANS outflow and increased SANS activity—actions leading to increases in cardiac output and total peripheral resistance.



**Note**

Baroreceptor reflexes can be blocked at the ganglionic synapse with  $N_N$  receptor antagonists. Alternatively, a reflex bradycardia can be blocked with muscarinic antagonists; a reflex tachycardia can be blocked with  $\beta_1$  antagonists.

Figure II-1-2. Automatic Feedback Loop

**Hormonal Feedback Loop**

- Blood pressure is also regulated via the hormonal feedback loop shown in Figure-II-1-3.
- The system is affected only by *decreases* in mean blood pressure (hypotension), which result in decreased renal blood flow.
- Decreased renal pressure causes the release of renin, which promotes formation of the angiotensins.
- Angiotensin II increases aldosterone release from the adrenal cortex, which, via its mineralocorticoid actions to retain sodium and water, increases blood volume.
- Increased venous return results in an increase in cardiac output.
- Angiotensin II also causes vasoconstriction, resulting in an increase in TPR.

## Section II: Autonomic Pharmacology

### Note

#### Antihypertensive Drugs

Both the ANS (neural) and endocrine feedback loops are invoked when patients are treated with antihypertensive drugs. Such compensatory mechanisms may result in tachycardia and both salt and water retention.








Figure II-1-3. Hormonal Feedback Loop


## Introduction to Blood Pressure/Heart Rate Tracings



### A Blood pressure:



- Increases are seen as deflections of the tracing upward 
- Decreases are seen as deflections of the tracing downward 
- Following mean blood pressure changes is enough 

### B Heart rate:



- Increases are seen as tighter tracing 
- Decreases are seen as a wider tracing 

### C Example of a drug X changing baseline parameters by decreasing mean blood pressure and increasing heart rate:




Figure II-1-4. Blood Pressure/Heart Rate Tracings

**PUPILLARY SIZE AND ACCOMMODATION MECHANISMS**




Figure II-1-5. Effects of ANS Drugs on the Eye

### Chapter Summary

The autonomic nervous system (ANS) is the major involuntary portion of the nervous system and is responsible for automatic, unconscious bodily functions. It has two major parts: the parasympathetic (PANS) and the sympathetic (SANS) systems.

Ganglia are relay systems set between the CNS and end organs. Ganglia in the SANS system are arranged in a series of parallel nodes adjacent to the vertebral column. In contrast, PANS ganglia are usually located in the innervated organ.


The major receptor types are ganglionic nicotinic ( $N_A$ ), endplate nicotinic ( $N_M$ ), muscarinic ( $M_{1-3}$ ), and adrenergic receptor of four major subtypes ( $\alpha_1, \alpha_2, \beta_1, \beta_2$ ). ACh is the neurotransmitter at all N receptors, at the M receptors innervated by postganglionic fibers of the PANS, and the thermoregulatory sweat glands innervated by the SANS. Norepinephrine (NE) is the neurotransmitter at adrenoceptors innervated by the SANS. NE and epinephrine (E) are released from the adrenal medulla. Dopamine (DA) receptor activation leads to vasodilation in some vascular beds.

Blood pressure (BP) is a product of the total peripheral resistance (TPR) times the cardiac output (CO). The CO is equal to the heart rate (HR) times the stroke volume (SV). The autonomic (neural) system helps regulate the BP through feedback control involving the baroreceptors, the cardiovascular centers in the brainstem, and the PANS and SANS, which act in an opposing but coordinated manner to regulate the pressure.

BP is also regulated by hormonal feedback (humoral). Hypotension decreases renal blood flow and activates the release of renin, which leads to the formation of angiotensin II, which in turn stimulates the release of aldosterone from the adrenal cortex. Aldosterone promotes water and salt retention, increasing blood volume and as a consequence increases SV and CO.

# Cholinergic Pharmacology

## CHOLINERGIC NEUROEFFECTOR JUNCTIONS



### Figure Legend

- ① Hemicholinium
- ② Botulinum toxin
- ③ Acetylcholinesterase (AChE) inhibitors
- ④ Receptor agonists and antagonists

Figure II-2-1. Cholinergic Neuroeffector Junction

- Choline is accumulated in cholinergic presynaptic nerve endings via an active transport mechanism linked to a Na<sup>+</sup> pump.
- Choline uptake is inhibited by hemicholinium (① in Figure II-2-1). ACh is synthesized from choline and acetyl-CoA via choline acetyltransferase (ChAT) and accumulated in synaptic vesicles.
- Presynaptic membrane depolarization opens voltage-dependent Ca<sup>2+</sup> channels, and the influx of this ion causes fusion of the synaptic vesicle membranes with the presynaptic membrane, leading to exocytosis of ACh. Botulinum toxin (② in Figure II-2-1) interacts with synaptobrevin and other proteins to prevent ACh release and is used in blepharospasm, strabismus/hyperhydrosis, dystonia, and cosmetics.



- Some cholinergic nerve endings have presynaptic autoreceptors for ACh that on activation may elicit a negative feedback of transmitter release.
- Inactivation via acetylcholinesterase (AChE) is the major mechanism of termination of postjunctional actions of ACh.
- AChE is a target for inhibitory drugs (indirect-acting cholinomimetics). Note that such drugs can influence cholinergic function only at innervated sites where ACh is released.
- Reversible AChE inhibitors (Ⓢ in Figure II-2-1) include edrophonium, physostigmine, and neostigmine. Irreversible AChE inhibitors include echothiophate, malathion, and parathion.
- Postjunctional receptors (N and M) (Ⓣ in Figure II-2-1) activated by ACh are major targets for both activating drugs (direct-acting cholinomimetics) and blocking agents.

**In A Nutshell**

- M receptor activation → ↓ CV function
- ↑ secretions and ↑ smooth muscle contraction
- All M receptor activators and blockers are nonspecific.

**Table II-2-1. Muscarinic Receptor Activation**

Target	Receptor	Response	
Eye	Sphincter	M <sub>3</sub>	Contraction—miosis
	Ciliary muscle	M <sub>3</sub>	Contraction—accommodation for near vision
Heart	SA node	M <sub>2</sub>	↓ Heart rate (HR)—negative chronotropy ↓ Conduction velocity—negative dromotropy No effects on ventricles, Purkinje system
	AV node	M <sub>2</sub>	
Lungs	Bronchioles	M <sub>3</sub>	Contraction—bronchospasm
	Glands	M <sub>3</sub>	Secretion
GI tract	Stomach	M <sub>3</sub>	↑ Motility—cramps Secretion Contraction—diarrhea, involuntary defecation
	Glands	M <sub>1</sub>	
	Intestine	M <sub>3</sub>	
Bladder	M <sub>3</sub>	Contraction (detrusor), relaxation (trigone/sphincter), voiding, urinary incontinence	
Sphincters	M <sub>3</sub>	Relaxation, except lower esophageal, which contracts	
Glands	M <sub>3</sub>	Secretion—sweat (thermoregulatory), salivation, and lacrimation	
Blood vessels (endothelium)	M <sub>3</sub>	Dilation (via NO/endothelium-derived relaxing factor)—no innervation, no effects of indirect agonists	

Table II-2-2. Nicotinic Receptor Activation

Target	Receptor	Response
Adrenal medulla	$N_N$	Secretion of epinephrine and NE
Autonomic ganglia	$N_N$	Stimulation—net effects depend on PANS/SANS innervation and dominance
Neuromuscular junction	$N_M$	Stimulation—twitch/hyperactivity of skeletal muscle

Note: N receptors desensitize very quickly upon excessive stimulation.

Table II-2-3. Cholinergic Receptor Mechanisms

$M_1$ and $M_3$	$G_q$ coupled	$\uparrow$ phospholipase C $\rightarrow$ $\uparrow$ $IP_3$ , DAG, $Ca^{2+}$
$M_2$	$G_i$ coupled	$\downarrow$ adenylyl cyclase $\rightarrow$ $\downarrow$ cAMP
$N_N$ and $N_M$	No 2nd messengers	activation (opening) of Na/K channels

### Bridge to Physiology and Anatomy

Blood vessels are solely innervated by the SANS, so the stimulation of autonomic ganglia results in vasoconstriction.

Conversely, the gastrointestinal tract is dominated by the PANS, so ganglionic stimulation causes increased gastrointestinal motility and secretions.

## MUSCARINIC RECEPTOR ACTIVATORS

### Muscarinic Agonists

Table II-2-4. Properties of Direct-Acting Cholinomimetics

Drug	Activity	AChE Hydrolysis	Clinical Uses
ACh	M and N	+++	Short half-life—no clinical use
Bethanechol	M	—	Rx—ileus (postop/neurogenic), urinary retention
Methacholine	M > N	+	Dx—bronchial hyperreactivity
Pilocarpine	M	—	Rx—glaucoma (topical), xerostomia

**Clinical Correlate**

**Alzheimer Disease**

Late-onset dementia with progressive memory loss and cognitive decline. Neuropathology includes neurofibrillary tangles, amyloid plaques, and loss of ACh neurons in Meynert's nucleus—rationale for clinical use of AChE inhibitors.

**Acetylcholinesterase Inhibitors**


**Table II-2-5. Properties of Indirect-Acting Cholinomimetics**

Drug	Characteristics	Clinical Uses
Edrophonium	Short-acting	Dx—myasthenia; used to differentiate myasthenia from cholinergic crisis
Physostigmine	Tertiary amine (enters CNS)	Rx—glaucoma; antidote in atropine overdose
Neostigmine, pyridostigmine	Quaternary amines (no CNS entry)	Rx—ileus, urinary retention, myasthenia, reversal of nondepolarizing NM blockers
Donepezil, tacrine	Lipid-soluble (CNS entry)	Rx—Alzheimer disease
Organophosphates	Lipid-soluble, irreversible inhibitors	Rx—glaucoma (echothiophate) Note: use as insecticides (malathion, parathion) and as nerve gas (sarin)

**Toxicity of AChE Inhibitors**

**As insecticides**

- Long-acting irreversible inhibitors (both carbamates and organophosphates)
- Wide use in agriculture as insecticides
- Examples: malathion and parathion



**Figure II-2-2. Activation of Organophosphate Insecticides**

**Classic Clue**

AChE inhibitor poisoning: "Dumbbells"

Diarrhea

Urination

Miosis

Bradycardia

Bronchoconstriction

Excitation (Muscle and CNS)

Lacrimation

Salivation

Sweating

**Acute toxicity**

- Excessive muscarinic and nicotinic stimulations
- Muscarinic effects:
  - Diarrhea
  - Urination
  - Miosis
  - Bradycardia

- Bronchoconstriction
- Lacrimation
- Salivation
- Sweating
- CNS stimulation
- Nicotinic effects:
  - Skeletal muscle excitation followed by paralysis
  - CNS stimulation

### Management

- Muscarinic effects: atropine
- Regeneration of AChE: pralidoxime (2-PAM)
- Time-dependent aging requires use of 2-PAM as soon as possible (see Figure II-2-3).

#### Irreversibly acting cholinomimetics:

These compounds phosphorylate the esteric site on AChE, at serine hydroxyl groups

- 1) phosphorylation; reversible by pralidoxime (2-PAM)
- 2) removal of a part of the organophosphate molecule (aging). Complex no longer reversible by 2-PAM.

R = leaving group  
P = organophosphate




Figure II-2-3. Effects of Organophosphate on AChE

#### Chronic toxicity

- Peripheral neuropathy causing muscle weakness and sensory loss
- Demyelination not due to AChE inhibition

## MUSCARINIC RECEPTOR ANTAGONISTS

### Atropine

- Prototype of the class
- As a tertiary amine, it enters CNS
- Other M blockers differ mainly in their pharmacokinetic properties

**Pharmacologic Effects**

- Atropine effects in order of increasing dose are:
  - Decreased secretions (salivary, bronchiolar, sweat)
  - Mydriasis and cycloplegia
  - Hyperthermia (with resulting vasodilation)
  - Tachycardia
  - Sedation
  - Urinary retention and constipation
  - Behavioral excitation and hallucinations
- Other classes of drugs with antimuscarinic pharmacology:
  - Antihistamines
  - Tricyclic antidepressants
  - Antipsychotics
  - Quinidine
  - Amantadine
  - Meperidine
- Treatment of acute intoxication:
  - Symptomatic ± physostigmine

**Table II-2-6. Clinical Uses and/or Characteristics of M Blockers**

Drug	Clinical Uses and/or Characteristics
Atropine	Antispasmodic, antisecretory, management of AChE inhibitor OD, antidiarrheal, ophthalmology (but long action)
Tropicamide	Ophthalmology (topical)
Ipratropium	Asthma and COPD (inhalational)—no CNS entry, no change in mucus viscosity
Scopolamine	Used in motion sickness, causes sedation and short-term memory block
Benztropine, trihexyphenidyl	Lipid-soluble (CNS entry) used in parkinsonism and in acute extrapyramidal symptoms induced by antipsychotics

**Bridge to Physiology**

**ANS Dominance**

For effector tissues with dual innervation, PANS is dominant. These include the SA and AV nodes of the heart, the pupil, GI and GU muscles, and sphincters. SANS is dominant only in terms of vascular tone and thermoregulatory sweat glands.

**NICOTINIC RECEPTOR ANTAGONISTS**

**Ganglion Blocking Agents**

- Drugs: hexamethonium and mecamylamine
- Reduce the predominant autonomic tone (see Table II-2-7)
- Prevent baroreceptor reflex changes in heart rate (see Figure II-2-4)
- Most are no longer available clinically because of toxicities (rarely, mecamylamine in hypertension)

Table II-2-7. Characteristics of Ganglion Blocking Agents

Effector	System	Effect of Ganglion Blockade
Arterioles	SANS	Vasodilation, hypotension
Veins	SANS	Dilation, ↓ venous return, ↓ CO
Heart	PANS	Tachycardia
Iris	PANS	Mydriasis
Ciliary muscle	PANS	Cycloplegia
GI tract	PANS	↓ Tone and motility—constipation
Bladder	PANS	Urinary retention
Salivary glands	PANS	Xerostomia
Sweat glands	SANS	Anhidrosis




Figure II-2-4. Algorithm: Reflex Control of Heart Rate

### Neuromuscular Blocking Drugs

See CNS Pharmacology, section IV, Drugs Used in Anesthesia.


**Chapter Summary**

- \* Acetylcholine (ACh) is synthesized from acetate and choline in the synaptic nerve via choline acetyltransferase and is stored in the synaptic vesicles and released by  $\text{Ca}^{2+}$  influx upon depolarization. The ACh then binds to a receptor on the other side of the synaptic junction, thereby transmitting the signal. Acetylcholinesterase (AChE) hydrolyzes the ACh and ends the signal. Cholinergic drugs are those that affect this process either by influencing ACh levels or by acting directly on the nicotinic or muscarinic receptors.
- \* Choline uptake is inhibited by hemicholinium. Botulinum toxin binds to synaptobrevin and prevents ACh release, and AChE inhibitors slow its rate of breakdown. Several reversible AChE inhibitors are useful pharmacologic agents; the irreversible AChE inhibitors are generally poisons.
- \* A cholinomimetic is a drug that imitates ACh. Nicotine acts as a cholinomimetic on nicotinic receptors, whereas bethanechol and pilocarpine are cholinomimetics that act on muscarinic receptors.
- \* Other drugs are ACh receptor blockers. Specific blocking agents acting on ganglionic nicotinic ( $\text{N}_\text{N}$ ) receptors are hexamethonium and mecamylamine. Those acting on the end-plate nicotinic receptors ( $\text{N}_\text{M}$ ) are tubocurarine, atracurium, and succinylcholine. Those acting on muscarinic (M) receptors include atropine, benztropine, and scopolamine.
- \* All M-receptor activators are nonspecific (they act on  $\text{M}_{1-3}$ ), and, in general, M-receptor activation decreases cardiovascular function and increases secretions and smooth muscle contraction. Table II-2-1 shows the type of M receptor involved and specific end-organ responses to M-receptor activators.
- \* Table II-2-2 summarizes the effects of nicotinic receptor activation on the adrenal medulla, the autonomic ganglia, and the neuromuscular junction. The effect of autonomic ganglia stimulation depends upon the transmission system used to connect the ganglia to the end organ. Blood vessels are innervated by SANS, resulting in vasoconstriction. PANS innervates the gut, the end result being increased motility and secretion.
- \* Table II-2-3 summarizes the receptor mechanisms used by the various receptor types.
- \* Table II-2-4 summarizes the activity, properties, and clinical uses for the direct-acting cholinomimetics, and Table II-2-5 does the same for the indirect-acting ones.
- \* Long-acting AChE inhibitors are commonly used as insecticides. Although these are less toxic for humans, they still provide a hazard, causing poisoning with both acute and chronic symptoms caused by both muscarinic and nicotinic hyperactivity (“dumbbells”).
- \* Therapy for acute poisoning by AChE inhibitors includes administration of M blockers (atropine) and pralidoxime (2-PAM), which helps reactivate AChE.
- \* Atropine is the prototype of muscarinic receptor antagonist drugs. In simple terms, increasing doses of atropine progressively decreases secretions and causes mydriasis, blurred vision, tachycardia, constipation, and urinary retention. Overdoses of over-the-counter medications containing M blockers are common causes of toxicity. Management is largely symptomatic, although physostigmine may be useful because it helps counteract both central and peripheral effects. The clinical uses and properties of the M-blocking drugs are summarized in Table II-2-6.
- \* The  $\text{N}_\text{N}$  antagonists act as ganglionic blockers, so they will affect both the SANS and PANS tracts.
- \* Ganglionic blockade prevents ANS reflexes. Table II-2-7 summarizes specific effects of ganglionic blocking agents and the transmission system employed for various specific organs.
- \* Figure II-2-4 summarizes the effects of ganglionic blockers on drugs that modify blood pressure, causing a reflex change in heart rate, and on drugs that act directly at the SA node to change the heart rate.

# Adrenergic Pharmacology

## CATECHOLAMINE SYNTHESIS, ACTION, AND DEGRADATION

The important aspects of the adrenergic neuroeffector junction are summarized in Figure II-3-1.



### Figure Legend

- ① Methyl-p-tyrosine
- ② MAO inhibitors
- ③ Releasers
- ④ Reuptake blockers
- ⑤ α<sub>2</sub> agonists and antagonists
- ⑥ Reserpine
- ⑦ Guanethidine
- ⑧ Agonists and blockers of α<sub>1</sub>, β<sub>1</sub> receptors

Figure II-3-1. Adrenergic Neuroeffector Junction

Tyrosine is actively transported into nerve endings and is converted to dihydroxyphenylalanine (DOPA) via tyrosine hydroxylase (1). This step is rate limiting in the synthesis of NE. DOPA is converted to dopamine (DA) via L-aromatic amino acid decarboxylase (DOPA decarboxylase). DA is taken up into storage vesicles where it is metabolized to NE via DA beta hydroxylase (6). Inactivation of NE via monoamine oxidase (MAO) (2) may regulate prejunctional levels of transmitter in the mobile pool (3) but not the NE stored in granules.

Presynaptic membrane depolarization opens voltage-dependent Ca<sup>2+</sup> channels. Influx of this ion causes fusion of the synaptic granular membranes, with the presynaptic membrane leading to NE exocytosis into the neuroeffector junction (7). NE then activates postjunctional receptors (8), leading to tissue-specific responses depending on the adrenoceptor subtype activated.



Termination of NE actions is mainly due to removal from the neuroeffector junction back into the sympathetic nerve ending via an NE reuptake transporter system (4). At some sympathetic nerve endings, the NE released may activate prejunctional alpha adrenoceptors (5) involved in feedback regulation, which results in decreased release of the neurotransmitter. Metabolism of NE is by catechol-O-methyltransferase (COMT) in the synapse or MAO<sub>A</sub> in the prejunctional nerve terminal.

**Table II-3-1. Adrenergic Receptor Activation**

Receptor	Response
<b>α<sub>1</sub></b> Eye—radial (dilator) muscle Arterioles (skin, viscera) Veins Bladder trigone and sphincter Male sex organs Liver Kidney	Contraction—mydriasis Contraction—↑ TPR—↑ diastolic pressure, ↑ afterload Contraction—↑ venous return—↑ preload Contraction—urinary retention Vas deferens—ejaculation ↑ Glycogenolysis ↓ Renin release
<b>α<sub>2</sub></b> Prejunctional nerve terminals Platelets Pancreas	↓ Transmitter release and NE synthesis Aggregation ↓ Insulin secretion
<b>β<sub>1</sub></b> Heart SA node AV node Atrial and ventricular muscle His-Purkinje Kidney	↑ HR (positive chronotropy) ↑ Conduction velocity (positive dromotropy) ↑ Force of contraction (positive inotropy), conduction velocity, CO and oxygen consumption ↑ Automaticity and conduction velocity ↑ Renin release
<b>β<sub>2</sub> (mostly not innervated)</b> Blood vessels (all) Uterus Bronchioles Skeletal muscle Liver Pancreas	Vasodilation—↓ TPR—↓ diastolic pressure— ↓ afterload Relaxation Dilation ↑ Glycogenolysis—contractility (tremor) ↑ Glycogenolysis ↑ Insulin secretion
<b>D<sub>1</sub> (peripheral)</b> Renal, mesenteric, coronary vasculature	Vasodilation—in kidney ↑ RBF, ↑ GFR, ↑ Na <sup>+</sup> secretion


**In A Nutshell**

**Adrenoceptor Sensitivity**

Beta receptors are usually more sensitive to activators than alpha receptors. With drugs that exert both effects, the beta responses are dominant at low doses; at higher doses, the alpha responses will predominate.

**Note**

**Dopamine Use in Shock**



Fenoldopam is a D<sub>1</sub> agonist used for severe hypertension.

Table II-3-2. Mechanisms Used by Adrenergic Receptors

$\alpha_1$	$G_q$ coupled	$\uparrow$ phospholipase C $\rightarrow$ $\uparrow$ $IP_3$ , DAG, $Ca^{2+}$
$\alpha_2$	$G_i$ coupled	$\downarrow$ Adenylyl cyclase $\rightarrow$ $\downarrow$ cAMP
$\beta_1 \beta_2 D_1$	$G_s$ coupled	$\uparrow$ Adenylyl cyclase $\rightarrow$ $\uparrow$ cAMP

## DIRECT-ACTING ADRENOCEPTOR AGONISTS

### $\alpha_1$ Agonists




Figure II-3-2. Effect of Alpha Activators on Heart Rate and Blood Pressure

- Systemically,  $\uparrow$  mean blood pressure via vasoconstriction
- $\uparrow$  BP may elicit a reflex bradycardia
- Cardiac output may be  $\downarrow$  but also offset by  $\uparrow$  venous return
- Drugs and uses:
  - **Phenylephrine**: nasal decongestant and ophthalmologic use (mydriasis without cycloplegia)
  - **Methoxamine**: paroxysmal atrial tachycardia through vagal reflex

### $\alpha_2$ Agonists

Stimulate prejunctional receptors in the CNS to decrease sympathetic outflow. Primary use is in mild to moderate HTN.

- Drugs and uses: **clonidine** and **methyldopa** (mild to moderate hypertension)
- See Cardiovascular section.

### $\beta$ Agonists




Figure II-3-3. Effect of Beta Receptor Activation on Heart Rate and Blood Pressure

- Systemically,  $\downarrow$  mean BP via vasodilation ( $\beta_2$ ) and  $\uparrow$  HR ( $\beta_1$ )
- Drugs and uses:
  - Isoproterenol ( $\beta_1 = \beta_2$ ): bronchospasm, heart block, and bradyarrhythmias. Side effects include flushing, angina, arrhythmias.
  - Dobutamine ( $\beta_1 > \beta_2$ ): congestive heart failure
  - Selective  $\beta_2$  agonists:
    - Salmeterol, albuterol, and terbutaline used in asthma
    - Ritodrine, used in premature labor

### Mixed-Acting Agonists: Norepinephrine vs. Epinephrine

#### Norepinephrine ( $\alpha_1, \alpha_2, \beta_1$ )






Figure II-3-4. Effect of Norepinephrine on Heart Rate and Blood Pressure

Epinephrine ( $\alpha_1, \alpha_2, \beta_1, \beta_2$ )




- $\beta_1$ :  $\uparrow$  HR,  $\uparrow$  SV,  $\uparrow$  CO,  $\uparrow$  pulse pressure
- $\beta_2$ :  $\downarrow$  TPR,  $\downarrow$  BP

Figure II-3-5a. Effect of Low-dose Epinephrine on Heart Rate and Blood Pressure



- $\beta_1$ :  $\uparrow$  HR,  $\uparrow$  SV,  $\uparrow$  CO,  $\uparrow$  pulse pressure
- $\beta_2$ :  $\downarrow$  TPR,  $\downarrow$  BP
- $\alpha_1$ :  $\uparrow$  TPR,  $\uparrow$  BP

Figure II-3-5b. Effect of Medium-Dose Epinephrine on Heart Rate and Blood Pressure



- $\alpha_1$ :  $\uparrow$  TPR,  $\uparrow$  BP
- Potential reflex bradycardia
- $\beta_1$ :  $\uparrow$  HR,  $\uparrow$  SV,  $\uparrow$  CO,  $\uparrow$  pulse pressure
- $\beta_2$ :  $\downarrow$  TPR,  $\downarrow$  BP

Figure II-3-5c. Effect of High-dose Epinephrine is Similar to Norepinephrine

- Dose-dependent effects:
  - Low-dose:  $\beta_1, \beta_2$  stimulation (see Figure II-3-5a)
  - High-dose:  $\alpha_1, \beta_1 (\beta_2)$  (see Figure II-3-5c)
- $\beta_2$ -specific effects:
  - Smooth muscle relaxation: bronchioles, uterus, blood vessels
  - Metabolic effects:
    - $\uparrow$  glycogenolysis (muscle and liver)
    - $\uparrow$  gluconeogenesis
    - $\uparrow$  mobilization and use of fat

### Classic Clue

Indirect-acting adrenoceptor agonists act only on effector tissues innervated by SANS.

Denervated effector tissues are nonresponsive because these drugs act either to release transmitter from nerve terminals or to inhibit neurotransmitter reuptake.

### In A Nutshell

#### Forms of MAO

MAO type A: mainly in liver, but *Any*where (metabolizes NE, 5HT, and tyramine).

MAO type B: mainly in *Brain* (metabolizes DA).

- Differentiation of high-dose epinephrine versus norepinephrine:
  - Epinephrine reversal: Use of  $\alpha_1$  blocker to reverse hypertension to hypotension in a patient receiving too much epinephrine
  - Hypertension was due to predominant  $\alpha_1$  tone on the vasculature
  - Hypotension results from unmasking  $\beta_2$  receptors

### Uses of Norepinephrine and Epinephrine

- Cardiac arrest
- Adjunct to local anesthetic
- Hypotension
- Anaphylaxis (epinephrine only)
- Asthma (epinephrine only)

### INDIRECT-ACTING ADRENERGIC RECEPTOR AGONISTS

- Releasers:
  - Displace norepinephrine from mobile pool
  - Drug interaction: MAO<sub>A</sub> inhibitors (hypertensive crisis)
  - **Tyramine** (red wine, cheese)
    - Oral bioavailability is limited by MAO-A metabolism in gut and liver
    - MAO-A inhibition ↑ bioavailability, resulting in hypertensive crisis
  - **Amphetamines**
    - Clinical use of methyl phenidate in narcolepsy and ADHD
    - Psychostimulant due to central release of DA, NE, 5HT
  - **Ephedrine** (cold medication)
- Reuptake inhibitors:
  - Cocaine
  - Tricyclic antidepressant (in part)

### $\alpha$ RECEPTOR ANTAGONISTS

- ↓ TPR, ↓ mean BP
- May cause reflex tachycardia and salt and water retention
- Major uses:
  - Hypertension
  - Pheochromocytoma (nonselective  $\alpha$  blocker)
  - Benign prostatic hyperplasia (BPH; selective  $\alpha_1$  blocker)
- Drugs:
  - Nonselective blocker:
    - Phentolamine, competitive inhibitor
    - Phenoxybenzamine, noncompetitive inhibitor

- Selective  $\alpha_1$  blocker:
  - Prazosin, doxazosin, terazosin, tamsulosin
- Selective  $\alpha_2$  blocker:
  - Yohimbine: used in postural hypotension and impotence
  - Mirtazapine: used as antidepressant

## $\beta$ RECEPTOR ANTAGONISTS

- $\beta_1$  blockade:
  - $\downarrow$  HR,  $\downarrow$  SV,  $\downarrow$  CO
  - $\downarrow$  renin release
  - $\downarrow$  aqueous humor production
- $\beta_2$  blockade:
  - May precipitate bronchospasm (in asthmatics) and vasospasm (in patients with vasospastic disorders)
  - Metabolic effects
    - Blocks glycogenolysis, gluconeogenesis
    - $\uparrow$  LDLs, TGs

Table II-3-3. Characteristics of Some Beta Blockers

Drugs	$\beta_1$ -Selective	ISA	Sedation	Blood Lipids
Acebutolol	+	++	+	-
Atenolol	+	-	-	$\uparrow\uparrow$
Metoprolol	+	-	+	$\uparrow\uparrow$
Pindolol	-	++	+	-
Propranolol	-	-	+++	$\uparrow\uparrow$
Timolol	-	-	++	$\uparrow\uparrow$

- Cardioselectivity ( $\beta_1$ ):
  - Less effect on vasculature, bronchioles, uterus, and metabolism
  - Safer in asthma, diabetes, peripheral vascular diseases
- Intrinsic sympathomimetic activity (ISA):
  - Act as partial agonists
  - Less bradycardia ( $\beta_1$ )
  - Slight vasodilation or bronchodilation ( $\beta_2$ )
  - Minimal change in plasma lipids ( $\beta_2$ )
- Pharmacokinetic properties:
  - No CNS entry of atenolol
- General uses of beta-blockers:
  - Angina, hypertension, post-MI (all drugs)
  - Antiarrhythmics (class II: propranolol, acebutolol, esmolol)

### Clinical Correlate

Chronic use of beta blockers (e.g., in angina, HTN) leads to receptor upregulation.

During withdrawal from use, it is important to taper dose to avoid excessive cardiovascular effects (rebound effects) of endogenous amines.

### Clinical Correlate

#### Glucagon and the Heart

Positive inotropic and chronotropic, not via activation of  $\alpha_1$  receptors, but through glucagon receptors that are G-protein linked to adenylyl cyclase  $\rightarrow$  basis for its use in beta-blocker overdose.

- Glaucoma (timolol)
- Migraine, thyrotoxicosis, performance anxiety, essential tremor (propranolol)
- Combined alpha-1 and beta blocking activity:
  - Labetalol and carvedilol
  - Uses in CHF
- K<sup>+</sup>-channel blockade and β-blocking activity
  - Sotalol
  - Use as antiarrhythmic (class III)

### Chapter Summary

Neurotransmission across adrenergic junctions is mediated by norepinephrine (NE). Adrenergic effectors may act indirectly by influencing NE synthesis, monoamine oxidase (MAO) enzymes, the mobile NE pool, the NE transporter, prejunctional  $\alpha$ -adrenoceptors, granule uptake, or release of NE, or they may act directly on the postjunctional receptor as agonists or antagonists.

Excess NE normally subjects tyrosine hydroxylase to feedback inhibition, making this enzyme the rate-limiting step in the synthetic pathway of NE and epinephrine. Tyrosine conversion to DOPA can be inhibited by methyl-p-tyrosine, a tyrosine hydroxylase inhibitor.

MAO inhibitors regulate presynaptic NE levels.

Amphetamine, ephedrine, and tyramine act, in part, by releasing NE from the mobile pool (NE stored outside granules but within the neuron).

Cocaine and the tricyclic antidepressants act by inhibiting NE reuptake, which normally removes NE from the environment and makes it unavailable as a transmitter and also conserves it for future use.

Prejunction availability of NE can also be decreased by inhibiting NE release from the granules. This can be achieved by drugs such as clonidine or methyl-dopa, which are activators of the prejunctional  $\alpha_2$ -adrenoceptor; by drugs such as guanethidine, which act directly on the granules; or by drugs such as reserpine, which reduce NE levels by inhibiting granule uptake.

Table II-3-1 summarizes the distribution and physiologic effects associated with the activation of alpha 1 and 2, beta 1 and 2, and D<sub>1</sub> receptors. Table II-3-2 summarizes the mechanism through which these receptors work.

The major direct-acting adrenoceptor agonist drugs are described. The alpha agonist phenylephrine increases mean BP, has no effect on pulse pressure, and elicits a reflex bradycardia. Isoproterenol, a beta agonist, decreases mean BP, increases pulse pressure, and causes marked tachycardia. Cardiovascular effects of norepinephrine (NE) are similar to phenylephrine, but it is also a cardiac  $\beta_1$  adrenoceptor activator. The cardiovascular effects of epinephrine (E) are betalike at low doses and alphaslike at high doses.

The nonselective alpha blockers (phentolamine, phenoxybenzamine) are described. The  $\alpha_1$ -selective blockers (e.g., prazosin) are used in hypertension and BPH.

The properties, clinical uses, and adverse effects of the nonselective beta receptor antagonist propranolol are described. A comparison of beta adrenoceptor antagonists that are cardioselective and those that have intrinsic sympathomimetic activity is made (Table II-3-3). Drugs that block both alpha and beta adrenoceptors are identified.

# Autonomic Drugs: Glaucoma Treatment and ANS Practice Problems

## GLAUCOMA TREATMENT




Figure II-4-1. Anatomy of the Eye Showing Irido-Corneal Angle Where Aqueous Humor Is Recirculated

## Glaucoma

### Open-Angle Glaucoma

A chronic condition with increased intraocular pressure (IOP) due to decreased reabsorption of aqueous humor, leading to progressive (painless) visual loss and, if left untreated, blindness. IOP is a balance between fluid formation and its drainage from the globe. Strategies in drug treatment of glaucoma include the use of beta blockers to decrease formation of fluid by ciliary epithelial cells and the use of muscarinic activators to improve drainage through the canal of Schlemm (see Table II-4-1).



### Closed-Angle Glaucoma

An acute (painful) or chronic (genetic) condition with increased IOP due to blockade of the canal of Schlemm. Emergency drug management prior to surgery usually involves cholinomimetics, carbonic anhydrase inhibitors, and/or mannitol.

### Treatment

Table II-4-1. Mechanism of Action of Drugs Used to Treat Glaucoma

Drug	Drug Class	Mechanism of Action
Pilocarpine, echothiophate	Cholinomimetic	Activation of M receptors causes contraction of ciliary muscle, which increases flow through the canal of Schlemm; echothiophate is an organophosphate AChE inhibitor → ↑ outflow
Timolol	Beta blockers	Block actions of NE at ciliary epithelium ↓ aqueous humor formation


### Note

Antimuscarinic drugs and  $\alpha_1$  agonists are contraindicated in closed-angle glaucoma.

### ANS PRACTICE PROBLEMS

Answers and explanations to the following practice questions (pages 66–71) are found on pages 83–84.

Figure II-4-2



- R is
- epinephrine
  - norepinephrine
  - phenylephrine
  - isoproterenol
  - terbutaline




Figure II-4-3

- U is
- A. epinephrine
  - B. norepinephrine
  - C. phenylephrine
  - D. isoproterenol
  - E. tyramine





Figure II-4-4


- S is
- A. epinephrine
  - B. norepinephrine
  - C. phenylephrine
  - D. isoproterenol
  - E. terbutaline

Figure II-4-5




- H is
- epinephrine
  - norepinephrine
  - phenylephrine
  - isoproterenol
  - albuterol

Figure II-4-6



Drug X is most like


- epinephrine
- isoproterenol
- norepinephrine
- phenylephrine
- terbutaline



X and Y are, respectively:

- A. isoproterenol and propranolol
- B. epinephrine and phenoxybenzamine
- C. norepinephrine and phentolamine
- D. terbutaline and phenylephrine
- E. acetylcholine and hexamethonium

Figure II-4-7




What is drug X?

- A. Hexamethonium
- B. Neostigmine
- C. Atropine
- D. Scopolamine
- E. Ipratropium

What would you expect to see if the infused drug was neostigmine?

Figure II-4-8



Given the following information:

- Contractile force is measured in an isolated arterial preparation, and heart rate is measured in an isolated heart preparation.
- One drug is added at each specified time.
- No washout between drugs

- A. Bethanechol
- B. Epinephrine
- C. Phenoxybenzamine
- D. Pindolol
- E. Methoxamine

Time 1:  
Time 2:  
Time 3:  
Time 4:  
Time 5:

Figure II-4-9

	RIGHT EYE	LEFT EYE
Without treatment	○	○
With tyramine	○	○
With epinephrine	○	○

The circles above represent the size of the pupils of a patient's eyes, without treatment and with two different treatments. The responses are compatible with the conclusion that the left eye had:

- A. beta adrenergics blocked
- B. alpha adrenergics blocked
- C. cholinesterase inhibited
- D. muscarinic receptors blocked
- E. sympathetic denervation

Figure II-4-10




Figure II-4-11

**Chapter Summary**

The drugs used to treat open-angle glaucoma and their modes of action are summarized in Table II-4-1. The effects of autonomic drugs affecting the cardiovascular system are summarized visually in Figures II-4-2 through II-4-11.



# Autonomic Drug List and Practice Questions

---

## Cholinergic Receptor Activators

Direct activators: bethanechol (M), methacholine (M and N), nicotine (N), pilocarpine (M)

AChE inhibitors: reversible—edrophonium, physostigmine, neostigmine

AChE inhibitors: irreversible—echothiophate, malathion, parathion

## Cholinergic Receptor Antagonists

Muscarinic blockers: atropine, benztropine, ipratropium, scopolamine

Ganglionic blockers: hexamethonium, mecamylamine

## Adrenergic Receptor Activators

$\alpha_1$  agonists: phenylephrine, methoxamine

$\alpha_2$  agonists: clonidine, methyl dopa

$\beta$  agonists: isoproterenol ( $\beta_1 = \beta_2$ ), dobutamine ( $\beta_1 > \beta_2$ )

$\beta_2$  agonists: albuterol, ritodrine, terbutaline, salmeterol

Mixed: dopamine ( $D_1, \beta_1, \alpha_1$ ), epinephrine ( $\alpha_1, \alpha_2, \beta_1, \beta_2$ ), norepinephrine ( $\alpha_1, \alpha_2, \beta_1$ )

Indirect-acting: amphetamine, cocaine, ephedrine, tyramine

## Adrenergic Receptor Antagonists

$\alpha_1$  antagonists: doxazosin, prazosin, terazosin

$\alpha_2$  antagonists: yohimbine, mirtazapine

Mixed  $\alpha$  antagonists: phenoxybenzamine, phentolamine

$\beta_1$  (cardioselective) antagonists: acebutolol, atenolol, metoprolol

$\beta_1, \beta_2$  (nonselective): pindolol, propranolol, timolol

$\alpha_1$  and  $\beta$  antagonists: carvedilol, labetalol



### Review Questions

1. Alpha-1 agonists cause reflex bradycardia, which can be blocked by
  - A. atenolol
  - B. atropine
  - C. mirtazepine
  - D. phenylephrine
  - E. propranolol
  
2. Which one of the following effects is caused by the ingestion of mushrooms that contain pilocarpine?
  - A. Tachycardia
  - B. Bronchodilation
  - C. Diarrhea
  - D. Hypertension
  - E. Hyperthermia
  
3. Increasing the concentration of norepinephrine in adrenergic synapses leads to
  - A. activation of dopa decarboxylase
  - B. increased release of norepinephrine
  - C. activation of presynaptic G<sub>i</sub> coupled receptors
  - D. stimulation of MAO
  - E. activation of tyrosine hydroxylase
  
4. Urination in the human subject is decreased by
  - A. muscarinic agonists
  - B. muscarinic antagonists
  - C. AChase inhibitors
  - D. Nicotinic antagonists
  - E. Spider venom

5. A 5-year-old child becomes ill while visiting relatives who have a farm in Arkansas. His symptoms include severe abdominal cramps with vomiting and diarrhea and profuse lacrimation and salivation. Pupillary constriction is marked. The most likely cause is exposure to
  - A. herbicides
  - B. antifreeze
  - C. lead-based paint
  - D. insecticides
  - E. rat poison
  
6. The activation of muscarinic receptors in bronchiolar smooth muscle is associated with
  - A. activation of adenylyl cyclase
  - B. decrease in cAMP formation mediated by G-proteins
  - C. increase in  $IP_3$  and DAG
  - D. inhibition of protein kinase C
  - E. opening of  $Na^+/K^+$  cation channels
  
7. Ganglion blocking agents are of little clinical value today but they are important drugs to know for solving cardiovascular drug problems because they can block
  - A. all muscarinic receptors
  - B. all nicotinic receptors
  - C. all autonomic reflexes
  - D. the direct actions of drugs on blood vessels
  - E. the direct actions of drugs on the heart
  
8. An 11-year-old boy was brought to the ER by some of his friends because he “started going crazy” after eating seeds from a plant while “trying to get high.” The boy was incoherent; his skin was hot and dry. His pupils were dilated and unresponsive to light. Blood pressure was 180/105, pulse 150, and rectal temp 40°C. The presumptive diagnosis was drug toxicity due to the ingestion of a compound similar to
  - A. cannabis
  - B. digoxin
  - C. mescaline
  - D. phencyclidine
  - E. scopolamine

9. Reflex tachycardia caused by the systemic administration of albuterol can be blocked by what drug?
- dobutamine
  - prazosin
  - phenylephrine
  - metoprolol
  - low-dose epinephrine
10. Cardiovascular effects of a new drug (X) that activates autonomic receptors are shown in the table below:

Parameter	Control	Drug X
Systolic BP	120 mm Hg	110 mm Hg
Diastolic BP	85 mm Hg	55 mm Hg
Heart rate	60/min	120/min

The most probable receptor affinities of drug X are

- $\alpha_1, \alpha_2$
  - $\alpha_1, \alpha_2, \beta_1$
  - $\beta_1, \beta_2$
  - $M_2$
  - $N_M$
11. Thermoregulatory sweat glands in the body utilize what type of pathway?
- Cholinergic nerves and muscarinic receptors
  - Adrenergic nerves and alpha-1 receptors
  - Adrenergic nerves and beta-2 receptors
  - Cholinergic nerves and  $N_M$  receptors
  - Neurohumorally-released epinephrine
12. Activation of postsynaptic  $M_2$  receptors on the heart is associated with
- activation of adenylyl cyclase
  - decrease in cAMP formation
  - increase in  $IP_3$  and DAG
  - inhibition of protein kinase C
  - opening of  $Na^+/K^+$  cation channels

13. The data in the table below show the effects of four drugs (#1–4) on mean blood pressure administered as individual agents before and after treatment with prazosin. The arrows denote the direction and intensity of drug actions on blood pressure.

Condition	Drug #1	Drug #2	Drug #3	Drug #4
Before prazosin			↓↓	
After prazosin			↓↓	↓

The order of drug #1 through drug #4 is best represented by

- A. epinephrine—tyramine—isoproterenol—norepinephrine  
 B. tyramine—isoproterenol—norepinephrine—epinephrine  
 C. norepinephrine—isoproterenol—epinephrine—tyramine  
 D. isoproterenol—epinephrine—tyramine—norepinephrine  
 E. norepinephrine—tyramine—isoproterenol—epinephrine
14. Prior to an eye exam a patient is given a drug that causes mydriasis but has no effect on accommodation. What is the most likely identity of this drug?
- A. mecamylamine  
 B. neostigmine  
 C. pilocarpine  
 D. phenylephrine  
 E. tropicamide
15. Following a myocardial infarct, a 40-year-old male patient is being treated prophylactically with propranolol. You would be concerned about the use of this drug if the patient also had what comorbid condition?
- A. Essential tremor  
 B. Glaucoma  
 C. Classic/stable angina  
 D. Supraventricular tachycardia  
 E. Diabetes
16. Following pretreatment with a muscarinic receptor blocking agent, the IV administration of norepinephrine is likely to result in
- A. HR and BP  
 B. HR and ↓ BP  
 C. ↓ HR and ↓ BP  
 D. ↓ HR and BP  
 E. no effect on HR, but BP

17. A 45-year-old man has recently been the recipient of a heart transplant. Which one of the following drugs is least likely to cause tachycardia in this patient?
- A. Amphetamine
  - B. Dobutamine
  - C. Epinephrine
  - D. Isoproterenol
  - E. Norepinephrine
18. A colleague with myasthenia gravis wants you to assist him to the ER because he is experiencing muscle weakness and has found it difficult to titrate his drug dosage because he has had the "flu." You note that he has a slight temperature, shallow respirations, and a gray-blue skin pallor. What would be the most appropriate drug to give to your colleague at this time?
- A. Albuterol
  - B. Edrophonium
  - C. Propranolol
  - D. Physostigmine
  - E. Scopolamine
19. Carvedilol is an effective antihypertensive agent that, like propranolol, is capable of blocking beta receptors. An important difference between the two drugs is that carvedilol
- A. is a selective blocker of cardiac  $\beta_1$  receptors
  - B. has intrinsic sympathomimetic activity
  - C. is available only as eye drops
  - D. has  $\alpha_1$  receptor blocking actions
  - E. stimulates  $\beta_2$  receptors in bronchioles
20. Neostigmine differs from pilocarpine in having effects on
- A. bladder tone
  - B. bowel motility
  - C. heart rate
  - D. salivary glands
  - E. skeletal muscle

## Questions 21–23

The table below shows the effects of three receptor activators on heart rate in anesthetized animals, administered as individual drugs and following pretreatment with one of four different receptor antagonists. The arrows denote the direction of effects on heart rate; the symbol (–) denotes no change from normal HR.

Antagonist Pretreatment	Agonist 1	Agonist 2	Agonist 3
None		↓	↓
Atropine		–	
Prazosin		–	
Propranolol	–	↓	↓
Mecamylamine		–	

Identify the agonist drugs from the following list:

- A. Acetylcholine
  - B. Low-dose epinephrine
  - C. Norepinephrine
  - D. Phenylephrine
  - E. Physostigmine
21. Agonist 1
22. Agonist 2
23. Agonist 3

### Answers

1. **Answer: B.** Bradycardia due to vagal stimulation is elicited by activation of muscarinic receptors in the heart. Atropine, which is an antagonist at M receptors, blocks bradycardia elicited by stimulation of the vagus, including reflex bradycardia due to increases in mean BP caused by vasoconstrictors.
2. **Answer: C.** Pilocarpine is present in several mushroom species including *Amanita muscaria*, the ingestion of which is associated with the stimulation of M receptors (parasympathomimetic effects). Activation of muscarinic receptors in the GI tract causes diarrhea. The activation by pilocarpine of M receptors present on vascular endothelial cells would lead to hypotension (not hypertension) via the release of NO. All of the other effects listed are typical of muscarinic antagonists.
3. **Answer: C.** In sympathetic nerve endings presynaptic  $\alpha_2$  receptors are coupled to inhibitory G-proteins. These receptors serve an autoregulatory function to inhibit further neurotransmitter release and also to decrease the synthesis of norepinephrine.
4. **Answer: B.** Urinary retention is a well known adverse effect of drugs that have antagonist effects on muscarinic receptors. In addition to the prototypic drug atropine, M blockers include drugs used in Parkinson disease, such as benztropine. Acetylcholine directly and AChE inhibitors (edrophonium, physostigmine) indirectly activate M receptors in the GU system, causing bladder contraction with voiding and incontinence. Activation, not inhibition, of nicotinic receptors in ANS ganglia would lead to the stimulation of PANS functions.
5. **Answer: D.** The symptoms of cholinergic excess seen in this child are indicative of exposure to insecticides such as the organophosphate parathion, which cause irreversible inhibition of acetylcholinesterase. Other symptoms may include CNS excitation and stimulation of the skeletal NMJ, ultimately leading to paralysis of respiratory muscles—"DUMBBELSS." In addition to symptomatic support, management of AChE inhibitor poisoning involves the use of atropine and 2-PAM.
6. **Answer: C.** Muscarinic receptors present in bronchiolar smooth muscle are of the  $M_3$  subtype coupled via  $G_q$  proteins to phospholipase C. Activation of this enzyme causes hydrolysis of phosphatidylinositol bisphosphate, with release of  $IP_3$  and DAG (the latter activates protein kinase C). Decreased formation of cAMP mediated via a  $G_i$  protein occurs with activation of  $M_2$  receptors such as those in the heart. Cation channel opening occurs in response to activation of nicotinic receptors.
7. **Answer: C.** Ganglion blockers (hexamethonium, mecamylamine) block  $N_N$  receptors at autonomic ganglia and the adrenal medulla. As such, they can block all autonomic reflexes including those elicited by changes in blood pressure. They have no effect on nicotinic receptors at the neuromuscular junction ( $N_M$ ) or on the direct actions of drugs on the blood vessels or heart.
8. **Answer: E.** The signs and symptoms experienced by this boy are highly suggestive of the ingestion of a compound with strong muscarinic receptor-blocking actions. The leaves and seeds of jimsonweed (*Datura stramonium*) contain anticholinergic compounds, including atropine, hyoscyamine, and scopolamine—approximately 50 seeds may cause severe toxicity. In addition to symptomatic support, management of poisoning (or drug overdose) due to M blockers may involve use of the AChE inhibitor physostigmine.

9. **Answer: D.** Although used primarily via inhalation for asthma, systemic effects of albuterol include vasodilation due to its  $\beta_2$  receptor activation. This can result in a decrease in TPR and mean BP, which elicits a reflex tachycardia. Reflex tachycardia could be blocked at the heart with a beta blocker such as metoprolol or by ganglion blockers (mecamylamine) which prevent all autonomic reflexes. Dobutamine stimulates beta-1 receptors causing tachycardia. Phenylephrine stimulates alpha-1 receptors which would raise TPR and BP and evoke a reflex bradycardia that doesn't block tachycardia caused by albuterol. Prazosin blocks alpha-1 receptors decreasing TPR and BP and causing a reflex tachycardia. Low-dose epinephrine stimulates beta-1 and beta-2 receptors and will cause tachycardia.
10. **Answer: C.** A decrease in mean blood pressure, an increase in pulse pressure, plus a marked increase in heart rate are characteristic of a drug such as isoproterenol. PVR and mean BP are decreased because of activation of  $\beta_2$  receptors in the vasculature. Systolic BP decreases less than diastolic BP because of activation of  $\beta_1$  receptors in the heart, leading to an increase in stroke volume, as well as the increase in heart rate.
11. **Answer: A.** Thermoregulatory sweat glands are innervated only by the sympathetic nervous system. The pathway is unusual in that the postganglionic neuron releases acetylcholine. Thus, the receptors on sweat glands are muscarinic ( $M_3$ ). The term *neurohumoral* means "nerve-blood." The only site in the ANS where neurohumoral transmission occurs is the adrenal medulla, where sympathetic nerve activity elicits the release of catecholamines (mostly epinephrine) into the blood. Epinephrine cannot bind to muscarinic receptors.
12. **Answer: B.** Postsynaptic muscarinic receptors on the heart ( $M_2$ ) are  $G_i$  protein coupled to inhibition of adenylyl cyclase and decreased formation of cAMP.
13. **Answer: E.** Of the drugs listed, only isoproterenol causes a decrease in mean blood pressure, because it activates beta receptors and has no effect on alpha receptors. This permits identification of drug #3 as isoproterenol. Prazosin is an alpha blocker, so one can anticipate that this drug would antagonize any increases in blood pressure that result from activation of  $\alpha_1$  receptors in the vasculature. Epinephrine (high dose), norepinephrine, and tyramine all exert pressor effects via activation of  $\alpha_1$  receptors. However, only epinephrine is active on  $\beta_2$  receptors, and this action would be revealed by vasodilation and a reversal of its pressor effects following treatment with an alpha blocker—"epinephrine reversal." Thus, drug #4 can be identified as epinephrine.
14. **Answer: D.** Mydriasis can be caused by either a muscarinic antagonist or an alpha-1 agonist. Cycloplegia (paralysis of accommodation) is caused by a muscarinic antagonist, but accommodation is unaffected by an alpha-1 agonist such as phenylephrine. Remember accommodation is a parasympathetic function only. Ganglionic blockade with mecamylamine would cause mydriasis and cycloplegia similar to a muscarinic blocker.
15. **Answer: E.** Propranolol is a nonselective beta blocker that causes hypoglycemia by blocking glycogenolysis and gluconeogenesis in the the liver and skeletal muscle. This is of particular concern in a patient with diabetes. The other conditions listed are all potential uses for beta blockers, including essential tremor where it is important to use a nonselective beta blocker.
16. **Answer: A.** Norepinephrine activates  $\alpha_1$  and  $\beta_1$  receptors, causing increases in PVR and CO. The increase in mean BP can elicit reflex bradycardia (vagal outflow leads to stimulation of cardiac M receptors), which may overcome the direct stimulatory effects of NE on the heart. However, reflex bradycardia is not possible following pretreatment with an M blocker. Thus, HR increases because of the direct activation of cardiac  $\beta_1$  receptors by NE.



17. **Answer: A.** This question is to remind you that indirect-acting sympathomimetics require innervation of the effector organ to exert effects. In this case, amphetamine would not be effective because the transplanted heart lacks sympathetic innervation; thus, there is no “mobile pool” of NE capable of being released by a drug. However, transplanted hearts retain receptors, including those ( $\beta_1$ ) responsive to direct-acting sympathomimetics. Heart transplants are not responsive to AChE inhibitors because they, too, are indirect acting and require vagal innervation to exert effects on the heart.
18. **Answer: B.** Edrophonium is a very short-acting (reversible) AChE inhibitor that has been used in the diagnosis of myasthenia gravis. The drug is useful for distinguishing between muscle weakness attributable to excessive cholinergic receptor stimulation (usually due to overdose of a AChE inhibitor) and the symptoms of myasthenia (reflecting inadequate treatment). If symptoms improve with a single dose of edrophonium, then an increase in the dose of neostigmine or pyridostigmine is indicated. If symptoms worsen, then the dose of neostigmine should be reduced.
19. **Answer: D.** The effectiveness of carvedilol in the management of hypertension and in congestive heart failure appears to be due to a combination of antagonistic actions at both alpha and beta adrenoceptors. Carvedilol is not a  $\beta_1$  selective blocking agent (unlike atenolol and metoprolol), and (unlike pindolol and acebutolol) it lacks intrinsic sympathomimetic activity.
20. **Answer: E.** As an inhibitor of AChE, neostigmine exerts effects to enhance the actions of ACh at all innervated effector sites where ACh is a neurotransmitter. These include all ANS ganglia, PANS postganglionic neuroeffector junctions, and SANS innervation of thermoregulatory sweat glands. Pilocarpine activates M receptors and has no effects at conventional dose levels on nicotinic receptors such as those in ANS ganglia and the skeletal NMJ.
21. **Answer: B; 22. Answer: D; 23. Answer: C**

Agonist 1 increases HR, presumably through direct activation of cardiac  $\beta_1$  receptors because the effect is blocked by propranolol but is not influenced by the alpha blocker (prazosin), the ganglion blocker (mecamylamine), or blockade of M receptors (atropine). Only two of the listed drugs directly activate cardiac receptors: epinephrine and norepinephrine. For NE, any direct cardiac stimulation is counteracted by reflex bradycardia resulting from the increase in mean BP via its activation of  $\alpha_1$  receptors in blood vessels (it has no effects on  $\beta_2$  vascular receptors). Therefore, agonist 1 is identified as epinephrine which activates both  $\beta_1$  and  $\beta_2$  receptors directly at low doses.

To identify agonists 2 and 3, recognize that although the alpha blocker prazosin simply neutralizes the effect of agonist 2 on HR, it reverses the effect of agonist 3. This could occur only if agonist 3 was capable of  $\beta_1$  receptor activation in the heart. Direct cardiac stimulation could occur with norepinephrine (agonist 3) but not with phenylephrine (agonist 2), which is a selective alpha-1 agonist.

**Explanations to Figures II-4-2 through II-4-11: Drug Identification from Effects on Heart Rate and Blood Pressure.**

**Figure II-4-2:** The effects of Drug R are changed by treatment with either an alpha or beta-blocker, so Drug R must have activity at both receptors (**choices C, D, and E** are ruled out). A pressor dose of epinephrine would be “reversed” by an alpha-blocker, not just decreased! Drug R is norepinephrine.

**Figure II-4-3:** The effects of Drug U are changed by treatment with the alpha-blocker, but not by the beta-blocker. Drug U must be an alpha-activator with no beta actions—the only choice is phenylephrine.

**Figure II-4-4:** The effects of Drug S are changed by treatment with the beta-blocker, but not by the alpha blocker (**choices A, B, and C** are ruled out). Terbutaline is  $\beta_2$  selective and would not increase heart rate directly. Drug S is isoproterenol. Note that option A would have been a possibility but one would have to assume a low-dose of epinephrine.

**Figure II-4-5:** The effects of Drug H are changed by treatment with either an alpha- or beta-blocker, so Drug H must have activity at both receptors (**choices C, D, and E** are ruled out). “Reversal” of a pressor effect can only occur if the drug has  $\beta_2$  activity (**choice B** is ruled out). Drug H is epinephrine.

**Figure II-4-6:** Mecamylamine blocked reflexed tachycardia induced by Drug X, which dropped blood pressure by vasodilation. Propranolol prevented all responses. Drug X is a  $\beta_2$  agonist (terbutaline).

**Figure II-4-7:** Drug X decreases TPR and BP, eliciting a reflex sympathetic discharge (note delay in response), resulting in increased CO. There is no direct effect on CO (**choices A, B, C, and E** are ruled out). Drugs X and Y are terbutaline and phenylephrine. Note that the alpha agonist does not antagonize the decrease in respiratory resistance (a  $\beta_2$  response).

**Figure II-4-8:** ACh (used as a drug) decreases blood pressure and heart rate, but the latter effect is overcome and reversed by a sympathetic reflex. Because Drug X abolishes only the reflex tachycardia, it must be the ganglion blocker hexamethonium (**choice A**). Remember, AChE inhibitors do not vasodilate because there is no parasympathetic innervation of the vasculature!

**Figure II-4-9:** No autonomic reflexes are possible in isolated preparations! Arterial contraction due to the alpha agonist (**choice E**) is reversed by the alpha-blocker (**choice C**). Arteriolar relaxation and tachycardia due to epinephrine (**choice B**) is reversed by the beta-blocker (**choice D**). Bethanechol (**choice A**) causes both arteriolar relaxation and bradycardia.

**Figure II-4-10:** Classic example showing that denervated tissues do not respond to indirect-acting agonists. In this case, tyramine fails to cause mydriasis in the left eye, but this eye is more responsive than the right eye to epinephrine (denervation supersensitivity).

**Figure II-4-11:** Block of tachycardia due to Drug P by hexamethonium is indicative of a sympathetic reflex that follows a decrease in BP due to a vasodilator (**choice B**). “Reversal” of bradycardia due to Drug Q by hexamethonium indicates a vagal reflex elicited by vasoconstriction (e.g., alpha activation) masking cardiac stimulation (e.g., beta activation) typical of norepinephrine (**choice C**). Tachycardia due to Drug R is unaffected by any antagonist, indicative of a beta activator (**choice D**). “Reversal” of tachycardia due to Drug S by hexamethonium indicates a sympathetic reflex masking a vagotomimetic action typical of a muscarinic activator (**choice A**); this is confirmed by the effect of atropine.



SECTION III

**Cardiac and Renal  
Pharmacology**



# Fundamental Concepts

## CARDIAC ACTION POTENTIAL

### Fast-Response Fibers: Cardiac Muscle, His-Purkinje System




Figure III-1-1. Cardiac Action Potentials in Fast-Response Fibers

#### Phase 0

- $\text{Na}^+$  channels open—sodium enters the cell down its concentration gradient (fast  $I_{\text{Na}}$ ), causing membrane depolarization.
- Rate of depolarization depends on number of  $\text{Na}^+$  channels open, which in turn depends on resting membrane potential of the cell.
- Class I antiarrhythmic drugs can slow or block phase 0 in fast-response fibers.

#### Phase 1

- $\text{Na}^+$  channels are inactivated.
- In some His-Purkinje cells, transient outward  $\text{K}^+$  currents and inward  $\text{Cl}^-$  currents contribute to the “notch” and overshoot.
- Antiarrhythmic drugs have no significant effects on these transient currents.

**Phase 2**

- Plateau phase in which a slow influx of  $\text{Ca}^{2+}$  ( $I_{\text{Ca-L}}$ ) is “balanced” by a late-appearing outward  $\text{K}^+$  current (the delayed rectifier current  $I_{\text{K}}$ ).
- Antiarrhythmic drugs have no significant effects on these currents during this phase of the action potential (AP).

**Phase 3**

- Repolarization phase in which the delayed rectifier  $\text{K}^+$  current rapidly increases as the  $\text{Ca}^{2+}$  current dies out because of time-dependent channel inactivation.
- Class III antiarrhythmic drugs slow this repolarization phase.
- Note that during phases 0 through 3 a slow  $\text{Na}^+$  current (“window current”) occurs, which can help prolong the duration of the action potential.

**Phase 4**

- Return of membrane to resting potential—maintained by activity of the  $\text{Na}^+/\text{K}^+$ -ATPase.

**Responsiveness**

- Capacity of a cell to depolarize, associated with the number of  $\text{Na}^+$  channels in a ready state (see Figure III-1-4, page 90).
- This in turn depends on resting membrane potential: the more negative the resting potential (RP), the faster the response.

**Conductance**

Rate of spread of an impulse, or conduction velocity—three major determinants:

- Rate of phase 0 depolarization—as  $V_{\text{max}}$  decreases, conduction velocity decreases and vice versa.
- Threshold potential—the less negative, the slower the conduction velocity.
- Resting potential—the more negative the RP, the faster the conduction.

**Slow-Response Fibers (SA and AV Nodes, Specialized Cells)**




Figure III-1-2. Cardiac Action Potentials in Slow-Response Fibers

- No appreciable  $\text{Na}^+$  current during phase 0 in these cells because the Na channels are either absent or in an inactive form because of the existing voltage.
- Depolarization depends on activation of  $\text{Ca}^{2+}$  channels ( $I_{\text{Ca-L}}$  and  $I_{\text{Ca-T}}$ ).
- Class IV antiarrhythmic drugs can slow or block phase 0 in slow-response fibers.
- During repolarization, the  $\text{Ca}^{2+}$  currents are opposed and overcome by the delayed rectifier  $\text{K}^+$  current. The relative magnitudes of these opposing currents determine the “shape” of the action potential.
- The major distinctive feature of slow fibers is their spontaneous depolarization, shown by the rising slope of phase 4 of the AP, referred to as the pacemaker potential or “pacemaker current.” Although not completely understood, pacemaker potential is a composite of inward  $\text{Na}^+$  ( $I_f$ ) and  $\text{Ca}^{2+}$  ( $I_{\text{Ca-T}}$ ) currents and outward  $\text{K}^+$  currents ( $I_K$ ).
- Class II and IV antiarrhythmic drugs can slow phase 4 in pacemaker fibers.

### Automaticity

- The ability to depolarize spontaneously confers automaticity on a tissue.
- The fastest phase 4 slope will determine the pacemaker of the heart, which is normally the SA node.

### Refractoriness

- The inability to respond to a stimulus—property of all cardiac cells.

#### Effective Refractory Period (ERP)

- No stimulus, of any magnitude, can elicit a response.
- Lasts into late stage 3 of the AP because  $\text{Na}^+$  channels are effectively inactivated and not in the “ready” state.
- Blockers of  $\text{K}^+$  channels prolong the ERP.

#### Relative Refractory Period (RRP)

- A strong stimulus can elicit a response, but the timing will be out of sync with the rest of the heart and arrhythmias may occur.
- Ratio of ERP to the action potential duration (APD) is a measure of refractoriness, as illustrated in Figure III-1-3. Decreases in ERP favor the formation and propagation of premature impulses.




Figure III-1-3. Relationship of ERP to APD



## Na<sup>+</sup> CHANNEL

### Activation




Figure III-1-4. Mechanism of Action of Voltage-Gated Na<sup>+</sup> Channels

- This voltage-gated channel, which is responsible for the fast Na current ( $I_{Na}$ ), exists in three conformations:
  - Resting or ready state
  - Open or active state
  - Inactivated or refractory state
- The channel has two gates: M (activating) and h (inactivating), both of which are sensitive to voltage changes.
- Inactivation of the h gate is slower; therefore, it stays open longer and the Na channel is active.

### Recovery

- Rate of recovery of the Na channel is dependent on resting potential (RP).
- Fastest rate of recovery occurs at normal RP, and recovery slows as membrane voltage increases.
- Rate of recovery is slower in ischemic tissue because cells may be partly depolarized at rest. This reduces the number of channels able to participate in the next depolarization, which leads to a decrease in conduction rate in ischemic tissue.
- Na channel blockers also slow the rate of recovery in such tissues.

## ANS REGULATION OF HEART RATE

- Nodal tissue, especially that of the SA node, is heavily innervated by both PANS and SANS fibers activating  $M_2$  and  $\beta_1$  receptors, respectively.
- Phase 4 slope is increased by an increase in cAMP resulting from  $\beta_1$  receptor activation and slowed by a decrease in cAMP resulting from  $M_2$  receptor activation.
- Increase in cAMP will:
  - Increase upstroke velocity in pacemakers by increase of  $I_{Ca-L}$
  - Shorten AP duration by increase of  $I_K$
  - Increase HR by increase of  $I_f$ , thus increasing slope of phase 4
- Decrease in cAMP:
  - Does the opposite plus produces a  $K^+$  current ( $I_{K/ACh}$ ), which slows the rate of diastolic depolarization and thus decreases HR
  - Beta blockers prevent cAMP formation, with primary effects on SA and AV nodal tissues.

### Chapter Summary

The sequences of ionic events in the action potential of cardiac cells are described.

Depolarization (phase 0) is due to  $Na^+$  influx in fast fibers and due to  $Ca^{2+}$  influx in SA and AV nodal cells. Class I antiarrhythmic drugs block  $Na^+$  influx and class IV antiarrhythmics block  $Ca^{2+}$  influx.

Repolarization (phase 3) in all cardiac cells is due to  $K^+$  efflux (delayed rectifier current) and this is blocked by class IA and class III antiarrhythmic drugs. Pacemaker currents (phase 4) are blocked by class II and class IV drugs.

Responsivity, capacity of a cell for depolarization, depends on resting membrane potential; conductance is the rate of potential spread; refractoriness is the inability to respond to excitation.

Figure III-1-4 depicts the M and h gates of cardiac  $Na^+$  channels. Three conformations exist—resting (ready), active (open), and inactive (refractory). Class I drugs are least active when  $Na^+$  channels are in the resting state (state-dependent actions).

Actions of class II antiarrhythmics (beta blockers) involve antagonism of SANS-mediated increases in cAMP, especially at SA and AV nodal cells to slow phase 0 and 4 of the action potential.



# Antiarrhythmic Drugs

## CLASS I: Na<sup>+</sup> CHANNEL BLOCKERS

### Class 1A

- Antiarrhythmic: block fast Na<sup>+</sup> channels ( $\downarrow I_{Na}$ )
- Preferentially in the open or activated state—"state-dependent" blockade
- $\uparrow$  action potential duration (APD) and effective refractory period (ERP)
- Also blocks K<sup>+</sup> channel (prolongs repolarization)
- Drugs:
  - **Quinidine**
    - In addition to the above, causes muscarinic receptor blockade, which can  $\uparrow$  HR and AV conduction.
    - May also cause vasodilation via alpha block with possible reflex tachycardia.
    - Orally effective, wide clinical use in many arrhythmias; in atrial fibrillation, need initial digitalization to slow AV conduction.
    - Adverse effects: cinchonism (GI, tinnitus, ocular dysfunction, CNS excitation), hypotension, prolongation of QRS and  $\uparrow$  QT interval associated with syncope (torsades).
    - Drug interactions: hyperkalemia enhances effects and vice versa; displaces digoxin from tissue binding sites, enhancing toxicity.
  - **Procainamide**
    - Less muscarinic receptor block
    - Metabolized via *N*-acetyltransferase (genotypic variation) to *N*-acetyl procainamide (NAPA), an active metabolite
    - Adverse effects: systemic lupus erythematosus (SLE)-like syndrome (30% incidence) more likely with slow acetylators; hematotoxicity (thrombocytopenia, agranulocytosis); CV effects (torsades)

### Class 1B

- Antiarrhythmic: block fast Na<sup>+</sup> channels ( $\downarrow I_{Na}$ )
- Block inactivated channels—preference for tissues partly depolarized (slow conduction in hypoxic and ischemic tissues). This results in an increased threshold for excitation and less excitability of hypoxic heart muscle.
- $\downarrow$  APD—due to block of the slow Na<sup>+</sup> "window" currents, but this increases diastole and extends the time for recovery.

### Note

For the exam, you should understand which effect is antiarrhythmic (slows heart) and which is proarrhythmic (speeds up heart).

### Note

Quinidine is a weak base, and antacids increase its absorption, thus greatly increasing its toxicity.

- Drugs and uses:
  - **Lidocaine**
    - Post-MI
    - Open-heart surgery
    - Digoxin toxicity
    - Side effects: CNS toxicity (seizures); least cardiotoxic of conventional antiarrhythmics
    - IV use because of first-pass metabolism
  - **Mexiletine and tocainide**
    - Same uses as lidocaine
    - Oral formulations

### Class 1C

- Block fast  $\text{Na}^+$  channels ( $\downarrow I_{\text{Na}}$ ), especially His-Purkinje tissue
- No effect on APD
- No ANS effects
- Drug:
  - **Flecainide**
    - Limited use because of proarrhythmic effects, leading to  $\uparrow$  in sudden death post-MI and when used prophylactically in VT

### CLASS II: BETA BLOCKERS

- Prevent  $\beta$ -receptor activation, which would normally  $\uparrow$  cAMP
- $\downarrow$  SA and AV nodal activity
- $\downarrow$  Slope of phase 4 (diastolic currents) of AP in pacemakers
- Drugs:
  - Propranolol (nonselective) and the cardioselective drugs: acebutolol and esmolol
  - Uses:
    - Prophylaxis post-MI and in supraventricular tachyarrhythmias (SVTs)
    - Esmolol (IV) is used in acute SVTs

### CLASS III: $\text{K}^+$ CHANNEL BLOCKERS

- $\downarrow I_{\text{K}}$  (delayed rectifier current) slowing phase 3 (repolarization) of AP
- $\uparrow$  APD and ERP, especially in Purkinje and ventricular diseases
- Drugs:
  - **Amiodarone**
    - Mimics classes I, II, III, and IV
    - Increase APD and ERP in all cardiac tissues
    - Uses: any arrhythmias
    - $t_{1/2} > 80$  days

- Binds extensively to tissues (large  $V_d$  and multiple effects)
- Side effects:
  - Pulmonary fibrosis
  - Blue pigmentation of the skin (“smurf skin”)
  - Phototoxicity
  - Corneal deposits
  - Hepatic necrosis
  - Thyroid dysfunction
- **Sotalol:**
  - ↓  $I_K$ , slowing phase III
  - $\beta_1$  blockade, leading to ↓ HR, ↓ AV conduction
  - Use: life-threatening ventricular arrhythmia

## CLASS IV: $Ca^{2+}$ CHANNEL BLOCKERS

- Block slow cardiac  $Ca^{2+}$  channels
- ↓ phase 0, ↓ phase 4
- ↓ SA, ↓ AV nodal activity
- Drugs:
  - **Verapamil and diltiazem**
    - Prototype  $Ca^{2+}$ -channels blockers (see Antihypertensive Drugs and Antianginal Drugs chapters in this section)
    - Uses: supraventricular tachycardias
    - Side effects: constipation (verapamil), dizziness, flushing, hypotension, AV block
    - Drug interaction:
      - Additive AV block with  $\beta$ -blockers, digoxin
      - Verapamil displaces digoxin from tissue-binding sites

## UNCLASSIFIED

- **Adenosine**
  - Activates adenosine receptors: causes  $G_i$ -coupled decrease in cAMP
  - ↓ SA and AV nodal activity
  - Uses: DOC for paroxysmal supraventricular tachycardias and AV nodal arrhythmias
  - Administered IV:  $t_{1/2}$  <10 seconds
  - Side effects: flushing, sedation, dyspnea
  - Adenosine is antagonized by theophylline
- **Magnesium**
  - Use: torsades
  - Drugs causing torsades include:
    - Potassium-channel blockers (class IA and class III)
    - Antipsychotics (thioridazine)
    - Tricyclic antidepressants

## Clinical Correlate

### Long QT Syndrome

A familial condition associated with increased risk of ventricular arrhythmias may result from mutation in the gene encoding cardiac potassium channels. Class IA and class III antiarrhythmic drugs may increase the risk of torsades in such patients.

### Treatment of Torsades

- Correct hypokalemia.
- Correct hypomagnesemia.
- Discontinue drugs that prolong the QT interval.
- Attempt to shorten APD with drugs (e.g., isoproterenol) or electrical pacing.

## Clinical Correlate

### Potassium

Both hyperkalemia and hypokalemia are arrhythmogenic.

### Chapter Summary

- \* The class I antiarrhythmic drugs block  $\text{Na}^+$  channels. Class IA drugs are state-dependent blockers of fast  $\text{Na}^+$  channels, and they increase the action potential duration (APD). Quinidine, in addition, is an M blocker and can increase the heart rate and AV conduction. Procainamide has less M block than quinidine and no alpha block. The uses and contraindications of quinidine and procainamide are provided.
- \* Class IB drugs are less state-dependent blockers of fast  $\text{Na}^+$  channels, and they decrease the APD. The uses for lidocaine, mexiletine, and tocainide are discussed, as are the metabolism and adverse effects of lidocaine.
- \* The class IC drug flecainide blocks fast  $\text{Na}^+$  channels, especially of His-Purkinje cells, and has no effect on the APD and no ANS effects.
- \* Class II antiarrhythmic drugs are beta-blockers that decrease SA and AV nodal activity, decrease the phase 4 slope, and prevent  $\beta_1$  adrenoceptor activation, thereby circumventing the normal increase in cAMP. Propranolol is nonselective; acebutolol and esmolol are selective. Their antiarrhythmic use is discussed.
- \* Class III antiarrhythmic drugs are  $\text{K}^+$ -channel blockers that increase the APD and effective refractor period (ERP), especially in Purkinje and ventricular tissues. Amiodarone and sotalol are the examples discussed.
- \* Class IV antiarrhythmic drugs are  $\text{Ca}^{2+}$ -channel blockers that decrease the SA and AV nodal activity and the slope of phase 4 of the action potential in pacemakers. The uses and adverse effects of verapamil are indicated.
- \* Adenosine and magnesium are two unclassified antiarrhythmic drugs. Adenosine decreases SA and AV node activity and increases the AV node refractory period. Magnesium has possible use in torsades. Drugs (other than classes Ia and III antiarrhythmics) associated with torsades include thioridazine and tricyclic antidepressants.

# Antihypertensive Drugs

---

## DRUG STRATEGY

- ↓ TPR
- ↓ CO
- ↓ body fluid volume
- ↓ BP may result in homeostatic regulation:
  - Reflex tachycardia (↑ sympathetic activity)
  - Edema (↑ renin activity)

## DRUGS ALTERING SYMPATHETIC ACTIVITY

- $\alpha_2$  agonists: **clonidine** and **methyldopa** (prodrug)
  - $\alpha_2$  stimulation:
    - ↓ in sympathetic outflow
    - ↓ TPR but also ↓ HR
  - Uses:
    - Mild-to-moderate hypertension (both)
    - Opiate withdrawal (clonidine)
    - Hypertensive management in pregnancy (methyldopa)
  - Side effects:
    - Positive Coombs test (methyldopa)
    - CNS depression (both)
    - Edema (both)
  - Drug interactions:
    - Tricyclic antidepressants ↓ antihypertensive effects of  $\alpha_2$  agonists
- Drugs interfering with storage vesicles
  - **Reserpine**
    - Destroys vesicles:
      - ↓ CO and ↓ TPR (↓ norepinephrine in periphery)
      - ↓ norepinephrine, ↓ dopamine, ↓ serotonin in CNS
    - Side effects:
      - Depression (often severe)
      - Edema
      - ↑ gastrointestinal secretions



- **Guanethidine**
  - Accumulated into nerve endings by reuptake
  - Binds vesicles
  - Inhibits norepinephrine release
  - Side effects:
    - Diarrhea
    - Edema
  - Drug interactions: tricyclic antidepressants block reuptake and the action of guanethidine
- $\alpha_1$  blockers
  - ↓ arteriolar and venous resistance
  - Reflex tachycardia
  - Drugs: **prazosin, doxazosin, terazosin**
  - Uses:
    - Hypertension
    - BPH: ↓ urinary frequency and nocturia by ↓ the tone of urinary sphincters
  - Side effects:
    - “First-dose” syncope
    - Orthostatic hypotension
    - Urinary incontinence
  - Advantage: good effect on lipid profile (↑ HDL, ↓ LDL)
- $\beta$  blockers
  - Mechanism (See ANS section)
  - Side effects:
    - Cardiovascular depression
    - Fatigue
    - Sexual dysfunction
    - ↑ LDLs and TGs
  - Cautions in use:
    - Asthma
    - Vasospastic disorders
    - Diabetics (alteration of glycemia and masking of tachycardia due to hypoglycemic events)

## DIRECT-ACTING VASODILATORS

### Drugs Acting Through Nitric Oxide

- **Hydralazine**
  - ↓ TPR via arteriolar dilation
  - Use: moderate-to-severe hypertension
  - Side effects:
    - SLE-like syndrome and slow acetylators
    - Edema
    - Reflex tachycardia



## DIURETICS (See Chapter 6)

Thiazide and loop diuretics are commonly used in the management of hypertension.

## ANGIOTENSIN-CONVERTING INHIBITORS (ACEIs) AND ANGIOTENSIN-RECEPTOR BLOCKERS (ARBs)




Figure III-3-2. The Angiotensin System

- Drugs:
  - ACEIs: **captopril** (and other “-prils”)
    - Block formation of angiotensin II
    - Resulting in prevention of AT<sub>1</sub>-receptor stimulation
    - ↓ aldosterone, vasodilation
    - ACEIs prevent bradykinin degradation
  - ARBs: **losartan** (and other “-sartans”)
    - Block AT<sub>1</sub> receptors
    - Same results as ACEIs on BP mechanisms
    - ARBs do not interfere with bradykinin degradation
  - Renin inhibitor: Aliskiren
    - Blocks formation of angiotensin I
    - Same results as ACEIs on BP mechanisms
    - Aliskiren does not interfere with bradykinin degradation
- Uses:
  - Mild-to-moderate hypertension (all)
  - Protective of diabetic nephropathy (ACEI/ARBs)
  - CHF (ACEI/ARBs)

- Side effects:
  - Dry cough (ACEIs)
  - Hyperkalemia
  - Acute renal failure in renal artery stenosis
  - Angioedema
- Contraindication: pregnancy

## INDICATIONS FOR USE OF ANTIHYPERTENSIVE DRUGS IN COMORBID CONDITIONS

Table III-3-1. Use of Antihypertensive Drugs in Comorbid Conditions

Indication	Suitable Drug(s)
Angina	Beta blockers, CCBs
Diabetes	ACEIs, ARBs
Heart failure	ACEIs, ARBs, Beta blockers
Post-MI	Beta blockers
BPH	Alpha blockers
Dyslipidemias	Alpha blockers, CCBs, ACEIs/ARBs

## TREATMENT OF PULMONARY HYPERTENSION

- **Bosentan**
  - Endothelin (ET)-1 is a powerful vasoconstrictor through ET-A and -B receptors
  - Bosentan is an ETA receptor antagonist
  - Administered orally
  - Side effects are associated with vasodilation (headache, flushing, hypotension, etc.)
  - Contraindication: pregnancy
- Prostacyclin (PGI<sub>2</sub>): **epoprostenol**
  - Administered via infusion pumps
- **Sildenafil**
  - Inhibits type V PDE
  - ↑ cGMP
  - Pulmonary artery relaxation
  - ↓ pulmonary hypertension

**Chapter Summary**

- \* Hypertension (HTN) is a major risk factor for stroke, heart failure, renal disease, peripheral vascular disease, and coronary artery disease. Factors inducing HTN include decreased vagal tone, increased sympathetic tone, increased renin-angiotensin activity, and excess water retention.
- \* Treatments for HTN aim to reduce sympathetic tone and blood volume and/or relax vascular smooth muscle. However, homeostatic mechanisms may lead to compensatory increases in heart rate and/or salt and water retention.
- \* The metabolic characteristics, clinical uses, and potential adverse effects of sympathoplegic drugs, which decrease peripheral resistance by decreasing sympathetic tone, are discussed. Those sympathoplegic drugs that act indirectly via the CNS include methyldopa and clonidine. Sympathoplegic drugs also may act directly as adrenergic neuron blockers, alpha blockers, or beta blockers. Examples of each class are provided.
- \* Direct-acting vasodilators lower the peripheral vascular resistance mainly by causing arteriolar dilation. Drugs discussed are nitroprusside, hydralazine, minoxidil, and diazoxide.
- \* Calcium channel blockers (CCBs) enhance vasodilation by blocking L-type  $\text{Ca}^{2+}$  channels in cardiac and vascular tissues. Drugs considered are verapamil, diltiazem, and dihydropyridines.
- \* Drugs that act via the renin-angiotensin system are the angiotensin-converting enzyme (ACE) inhibitors (e.g., captopril) and the angiotensin-II (AT-1) blockers (ARBs; e.g., losartan). Figure III-3-2 illustrates the angiotensin system and the pharmacologic effects of these drugs. Their clinical uses and adverse effects are discussed.
- \* Both thiazide and loop diuretics are used to treat HTN. The diuretics are discussed in more detail elsewhere.
- \* Table III-3-1 summarizes the use of antihypertensives in comorbid conditions.
- \* Bosentan, epoprostenol, and sildenafil are used in pulmonary hypertension.

# Drugs for Heart Failure




Figure III-4-1. The Failing Heart

Pharmacotherapy aimed at:

- ↓ preload: diuretics, ACEIs, ARBs, and venodilators
- ↓ afterload: ACEIs, ARBs, and arteriodilators
- ↑ contractility: digoxin, beta agonists
- ↓ remodeling of cardiac muscle: ACEIs, ARBs, spironolactone

Whereas digoxin does not improve survival, ACEIs, ARBs, carvedilol, and spironolactone have been proven beneficial in CHF. ACEIs and ARBs are currently drugs of choice for the chronic management of CHF. Inotropes are more beneficial in management of acute CHF.

## INOTROPES




Figure III-4-2. Mechanism of Action of Inotropes

- Digoxin
  - Direct effect: inhibition of cardiac  $\text{Na}^+-\text{K}^+$  ATPase
    - Results in ↑ intracellular  $\text{Na}^+$
    - ↓  $\text{Na}^+/\text{Ca}^{2+}$  exchange
    - ↑ intracellular  $\text{Ca}^{2+}$
    - ↑  $\text{Ca}^{2+}$  release from sarcoplasmic reticulum
    - ↑ actin-myosin interaction
    - ↑ contractile force
  - Indirect effect: inhibition of neuronal  $\text{Na}^+-\text{K}^+$  ATPase
    - Results in ↑ vagal and sympathetic stimulation


- Pharmacokinetics:
  - Long  $t_{1/2}$ : need loading dose (LD)
  - Renal clearance: caution in renal impairment
  - Tissue protein binding (large  $V_d$ ): displacement by other drugs (verapamil, quinidine)
- Uses:
  - CHF
  - Supraventricular tachycardias, except Wolff-Parkinson-White syndrome (*see margin note*)
- Side effects:
  - Early signs include anorexia, nausea, ECG changes
  - Later signs include disorientation, visual effects (halos)
  - In toxic doses, any cardiac arrhythmias
- Management of toxicity
  - Use of Fab antibodies toward digoxin
  - Supportive therapy (electrolytes and antiarrhythmics class IB)
- Drug interactions:
  - Diuretics:  $\downarrow K^+$ ,  $\downarrow Mg^{2+}$ ,  $\uparrow Ca^{2+}$
  - Quinidine and verapamil
- Phosphodiesterase inhibitors: inamrinone and milrinone
  - $\uparrow$  cAMP in heart muscle; results in  $\uparrow$  inotropy
  - $\uparrow$  cAMP in smooth muscle; results in  $\downarrow$  TPR
- Sympathomimetics: dobutamine and dopamine

## OTHER DRUGS

- Diuretics
  - Loops for associated backward failure
  - Spironolactone to  $\downarrow$  remodeling (with ACEIs)
- Metoprolol and carvedilol
- Nesiritide
  - Recombinant form of human B-type natriuretic peptide (rh BNP)
  - Binds to natriuretic peptide receptors, thus  $\uparrow$  cGMP, resulting in vasodilation
  - Used in acutely decompensated CHF

### Note

#### Wolff-Parkinson-White Syndrome



- Do:
  - block accessory pathway with  $I_A$  or III
- Don't:
  - slow AV conduction (avoid digoxin,  $\beta$ -blocker,  $Ca^{2+}$ -channel blocker, adenosine)



### Chapter Summary

Heart failure is an inability of the heart to pump with sufficient vigor to maintain an adequate cardiac output. The mechanisms involved are discussed and are illustrated in Figure III-4-1.

Drugs used to treat heart failure include those that decrease preload (e.g., diuretics, ACEIs, ARBs, and venodilators), those that decrease afterload (e.g., ACEIs, ARBs, and arteriodilators), and those that increase cardiac contractility (e.g., digoxin and beta agonists).

Digoxin enhances cardiac contraction by inducing a series of responses initiated by inhibiting the  $\text{Na}^+/\text{K}^+$  ATPase. Figure III-4-2 shows how inhibition of cardiac membrane  $\text{Na}^+/\text{K}^+$  ATPase leads to increased contractility.

Although digoxin ameliorates symptoms, it does not increase survival. That is why the ACEIs are the primary choice for treating heart failure.

Digoxin has potential toxic effects that are in part dependent upon the electrolyte balance.

Bipyridines, sympathomimetics, diuretics, beta blockers, and nesiritide also have uses in treating heart failure. Rationales for their use are indicated. A newer drug, nesiritide, is a recombinant form of natriuretic peptide.

# Antianginal Drugs

## RATIONALE FOR USE

Angina pectoris is the principal syndrome of ischemic heart disease, anginal pain occurring when oxygen delivery to the heart is inadequate for myocardial requirement.

- Classic angina (angina of effort or exercise) is due to coronary atherosclerotic occlusion
- Vasospastic or variant angina (Prinzmetal) is due to a reversible decrease in coronary blood flow

## Drug Strategies in Classic and Vasospastic Angina

Drug strategies in classic and vasospastic angina involve:

- ↑ oxygen delivery by ↓ vasospasm (nitrates and CCBs).
- ↓ oxygen requirement by ↓ TPR, CO, or both (nitrates, CCBs, and beta blockers).

## NITRATES




Figure III-5-1. Nitrates and the Nitric Oxide Pathway

### Clinical Correlate

#### Sildenafil (Viagra)

Phosphodiesterase 5 (PDE5) is found in blood vessels supplying the corpora cavernosa. Sildenafil inhibits PDE 5 → ↑ cGMP → vasodilation → ↑ blood flow → ↑ erectile response. If used concomitantly with nitrates or other potent vasodilators, the excessive fall in blood pressure may lead to death from cardiovascular causes, including myocardial infarct.

- Nitrates are prodrugs of nitric oxide
- Venodilation → ↓ preload → ↓ cardiac work → ↓ oxygen requirement
- Nitrates ↓ infarct size and post-MI mortality
- Drugs:
  - **Nitroglycerin:** sublingual, transdermal, and IV formulations
  - Isosorbide: oral, extended release for chronic use
  - Side effects:
    - Flushing, headache, orthostatic hypotension
    - Reflex tachycardia and fluid retention
    - Methemoglobinemia
  - Cautions and contraindications:
    - Tachyphylaxis with repeated use
    - Cardiovascular toxicity with sildenafil (*see Clinical Correlate, left*)

### CALCIUM CHANNEL BLOCKERS (CCBs)

- All CCBs can be used.
- Nifedipine is important for vasospastic angina.
- *See Antihypertensive chapter in this section.*

### BETA BLOCKERS AND CARVEDILOL

- Used in angina of effort
- β-blockers are contraindicated in vasospastic angina
- Carvedilol is clinically equivalent to isosorbide in angina of effort




Figure III-5-2. Mechanisms of Smooth Muscle Contraction and Relaxation and Drugs Affecting Them

### Chapter Summary

Angina is the principal syndrome caused by ischemic heart disease. The three variants are classic, Prinzmetal, and unstable.

The drug strategies are to increase oxygen supply by decreasing vasospasm (nitrates and calcium channel antagonists [CCBs]) and to decrease cardiac oxygen requirements by decreasing peripheral vascular resistance and/or cardiac output (nitrates, CCBs, and beta blockers).

Nitrates increase NO concentrations. Increased NO activates guanylyl cyclase; this increases cGMP levels, which dephosphorylates myosin light chains, decreasing their association with actin and thereby promoting smooth muscle relaxation. These mechanisms are summarized in Figure III-5-1.

NO-enhancing drugs used to treat angina include nitroglycerin and isosorbide.

The adverse effects of the nitrates are also considered.

CCBs decrease contractility and increase vasodilation by preventing the influx of  $\text{Ca}^{2+}$  required for muscle contraction. The sequence of reactions involved is summarized in Figure III-5-2. The CCBs considered are the dihydropyridines (e.g., nifedipine), verapamil, and diltiazem.

Beta blockers act directly on the heart by decreasing the heart rate, the force of contraction, and cardiac output, thereby decreasing the work performed.

# Diuretics




Figure III-6-1. Actions of Diuretics at the Various Renal Tubular Segments

## Hypokalemia and Alkalosis

Diuretics that block  $\text{Na}^+$  reabsorption at segments above the collecting ducts will increase sodium load to the collecting tubules and ducts ("downstream"). This results in increased loss of  $\text{K}^+$   $\rightarrow$  hypokalemia, and in the case of both loop and thiazide diuretics the associated loss of  $\text{H}^+$  results in alkalosis.

## OSMOTIC DIURETICS

- Mannitol (IV) inhibits water reabsorption throughout the tubule.
- It increases urine volume.
- Uses:
  - ↓ IOP in glaucoma
  - ↓ intracerebral pressure
  - Anuric states
  - Rhabdomyolysis
- Side effects: acute hypovolemia

## CARBONIC ANHYDRASE INHIBITORS




Figure III-6-2. Actions of Carbonic Anhydrase Inhibitors

- Drugs: **acetazolamide** and **dorzolamide**
- Mechanism: carbonic anhydrase inhibition, results in:
  - ↓  $H^+$  formation inside PCT cell
  - ↓  $Na^+/H^+$  antiport
  - ↑  $Na^+$  and  $HCO_3^-$  in lumen
  - ↑ diuresis
- Uses:
  - Glaucoma
  - Acute mountain sickness
  - Metabolic alkalosis
- Side effects:
  - Bicarbonaturia and acidosis
  - Hypokalemia
  - Hyperchloremia

- Paresthesias
- Renal stones
- Sulfonamide hypersensitivity

## LOOP DIURETICS




Figure III-6-3. Actions of Loop Diuretics on the Thick Ascending Loop (TAL)

- Drugs: **ethacrynic acid** and **furosemide**
- Mechanism:  $\text{Na}^+/\text{K}^+/2\text{Cl}^-$  transporter inhibition, results in:
  - $\downarrow$  intracellular  $\text{K}^+$  in TAL
  - $\downarrow$  back diffusion of  $\text{K}^+$
  - $\downarrow$  positive potential
  - $\downarrow$  reabsorption of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$
  - $\uparrow$  diuresis
- Uses:
  - Acute pulmonary edema
  - Heart failure
  - Hypertension
  - Refractory edemas
  - Acute renal failure
  - Anion overdose
  - Hypercalcemic states
- Side effects:
  - Sulfonamide hypersensitivity (furosemide)
  - Hypokalemia and alkalosis

### Note

#### Allergies to Sulfonamide-Containing Drugs

Cross allergenicity with:

- Carbonic anhydrase inhibitors
- All loop diuretics, except ethacrynic acid
- Thiazides
- Sulfa antibiotics
- Celecoxib



- Hypocalcemia
- Hypomagnesemia
- Hyperuricemia
- Ototoxicity (ethacrynic acid > furosemide)
- Drug interactions
  - Aminoglycosides (enhanced ototoxicity)
  - Lithium (chronic loop administration, ↓ clearance)
  - Digoxin (↑ toxicity due to electrolyte disturbances)

## THIAZIDES




Figure III-6-4. Actions of Thiazides on the Distal Convolved Tubule (DCT)

- Drugs: **hydrochlorothiazide** and **indapamide**
- Mechanism:  $\text{Na}^+/\text{Cl}^-$  transporter inhibition, results in:
  - ↑ luminal  $\text{Na}^+$  and  $\text{Cl}^-$  in DCT
  - ↑ diuresis
- Uses:
  - Hypertension, CHF
  - Nephrolithiasis (calcium stones)
  - Nephrogenic diabetes insipidus
- Side effects:
  - Sulfonamide hypersensitivity
  - Hypokalemia and alkalosis
  - Hypercalcemia
  - Hyperuricemia
  - Hyperglycemia
  - Hyperlipidemia (except indapamide)

- Drug interactions and cautions:
  - Digoxin ( $\uparrow$  toxicity due to electrolyte disturbances)
  - Avoid in patients with diabetes mellitus

## K<sup>+</sup>-SPARING AGENTS




Figure III-6-5. Actions of Potassium-Sparing Agents on Collecting Tubules

- Drugs:
  - **Spironolactone:** aldosterone-receptor antagonist
    - Uses:
      - Hyperaldosteronic state
      - Adjunct to K<sup>+</sup>-wasting diuretics
      - Antiandrogenic uses (female hirsutism)
      - Congestive heart failure
    - Side effects:
      - Hyperkalemia and acidosis
      - Antiandrogen
  - Amiloride and triamterene: Na<sup>+</sup>-channel blockers
    - Use: adjunct to K<sup>+</sup>-wasting diuretics, lithium-induced nephrogenic diabetes insipidus (amiloride)
    - Side effects: hyperkalemia and acidosis

### Note

Eplerenone is a selective aldosterone receptor blocker devoid of antiandrogenic effect.

Table III-6-1. Summary of the Modes of Action and Effects of the Various Classes of Diuretics

Drug	Mechanisms of Action	Urinary Electrolytes	Blood pH
Acetazolamide	Inhibition of carbonic anhydrase in PCT	↑ Na <sup>+</sup> ↑ K <sup>+</sup> ↑↑ HCO <sub>3</sub> <sup>-</sup>	Acidosis
Ethacrynic acid, furosemide, torsemide	Inhibition of Na <sup>+</sup> /K <sup>+</sup> /2Cl <sup>-</sup> cotransporter in TAL	↑↑ Na <sup>+</sup> ↑ K <sup>+</sup> ↑ Ca <sup>2+</sup> ↑ Mg <sup>2+</sup> ↑ Cl <sup>-</sup>	Alkalosis
Hydrochlorothiazide, indapamide, metolazone	Inhibition of Na <sup>+</sup> /Cl <sup>-</sup> cotransporter in DCT	↑ Na <sup>+</sup> ↑ K <sup>+</sup> ↑ Cl <sup>-</sup> ↓ Ca <sup>2+</sup>	Alkalosis
Amiloride, triamterene, spironolactone, eplerenone	Block Na <sup>+</sup> channels, block aldosterone receptors in collecting tubule	↑ Na <sup>+</sup> (small) ↓ K <sup>+</sup>	Acidosis

### Chapter Summary

Diuretics are used to treat HTN, heart failure, edema, renal dysfunction, hypercalcemia, renal stones, glaucoma, and mountain sickness. In addition to their diuretic action, the loop and thiazide diuretics also cause vasodilation.

Figure III-6-1 illustrates the water and ion exchange occurring in the various segments of a renal tubule and the site of action of the different classes of diuretics.

The positive and negative effects of IV mannitol, an osmotic diuretic, are discussed.

Carbonic anhydrase inhibitors (e.g., acetazolamide) act in the proximal tubule to decrease absorption of Na<sup>+</sup> and bicarbonate. The mechanisms involved are summarized in Figure III-6-2. The clinical uses and adverse effects are listed.

Loop diuretics (e.g., ethacrynic acid and furosemide) inhibit the Na<sup>+</sup>/K<sup>+</sup>/2Cl<sup>-</sup> cotransporter on the luminal membrane of the thick ascending loop. The mechanisms causing their diuretic actions (Figure III-6-3) and their clinical uses and adverse effects are discussed.

The thiazides (e.g., hydrochlorothiazide and indapamide) inhibit the Na<sup>+</sup>/Cl<sup>-</sup> cotransporter on the luminal membrane of the distal convoluted tubule. The mechanisms leading to their diuretic actions (Figure III-6-4) and their clinical uses and adverse effects are discussed.

Spironolactone, amiloride, and triamterene are K<sup>+</sup>-sparing, weak diuretics that act at the collecting tubule and duct level. The mechanisms leading to their diuretic actions (Figure III-6-5) and their clinical uses and adverse effects are discussed.

Table III-6-1 summarizes the mechanisms of action, the urinary electrolyte patterns, and the resultant blood pH associated with administration of the various classes of diuretics.

# Antihyperlipidemics

- ↑ risk of atherosclerosis is associated with hypercholesterolemia
- ↑ risk of cardiovascular and cerebrovascular diseases
- Treatment goal is to ↓ LDL cholesterol and atheroma plaque formation




Figure III-7-1. Site of Action of Statins, Niacin, and Gemfibrozil on the Synthesis of Lipids

## HMG-CoA REDUCTASE INHIBITORS

- Drugs: **lovastatin** and other “-statins”
- Mechanisms:
  - HMG-CoA reductase inhibition, results in:
    - ↓ liver cholesterol
    - ↑ LDL-receptor expression
    - ↓ plasma LDL
  - ↓ VLDL synthesis results in: ↓ triglyceridemia

- Side effects:
  - Myalgia, myopathy (check creatine kinase)
  - Rhabdomyolysis
  - Hepatotoxicity (check liver function tests)
- Drug interaction:
  - Gemfibrozil (↑ rhabdomyolysis)
  - Cytochrome P450 inhibitors enhance toxicity of statins

## BILE ACID SEQUESTRANTS

- Drugs: **cholestyramine** and **colestipol**
- Mechanism: complexation of bile salts in the gut, results in:
  - ↓ enterohepatic recirculation of bile salts
  - ↑ synthesis of new bile salts by the liver
  - ↓ liver cholesterol
  - ↑ LDL-receptor expression
  - ↓ blood LDL
- Side effects:
  - ↑ VLDL and triglycerides
  - Gastrointestinal disturbances
  - Malabsorption of lipid-soluble vitamins
- Drug interactions with orally administered drugs (warfarin, thiazides, digoxin, etc.)
- Contraindication: hypertriglyceridemia

## NICOTINIC ACID (NIACIN, VITAMIN B<sub>3</sub>)

- Mechanism: inhibition of VLDL synthesis, results in:
  - ↓ plasma VLDL
  - ↓ plasma LDL
  - ↑ plasma HDL
- Side effects:
  - Flushing, pruritus, rashes (use aspirin)
  - Hepatotoxicity

## GEMFIBROZIL

- Mechanism: activation of lipoprotein lipases, results in:
  - ↓ VLDL and IDL
  - Modest ↓ LDL
  - ↑ HDL in most patients
  - In some patients with combined hyperlipidemias can ↑ LDL
  - Used in hypertriglyceridemia

- Side effects:
  - Gallstones
  - Myositis

## EZETIMIBE

- Mechanism: prevents intestinal absorption of cholesterol, results in ↓ LDL
- Side effect: gastrointestinal distress

Table III-7-1. Summary of Antihyperlipidemic Uses

Lab Findings	Drugs
↑ Cholesterol	Cholestyramine, colestipol, ezetimibe
↑ Triglycerides	Gemfibrozil
↑ Cholesterol and triglycerides	Statins and niacin, ezetimibe

## ORLISTAT

- Therapeutic use: weight loss
- Mechanism: inhibits pancreatic lipase → ↓ triglyceride breakdown in the intestine
- Side effects: oily stools (steatorrhea), diarrhea; ↓ absorption of lipid-soluble vitamins

### Chapter Summary

- \* An aberrant serum lipid profile is associated with increased risk of atherosclerosis and cardiac heart disease.
- \* Lovastatin and the other statins inhibit the rate-limiting step in cholesterol synthesis, HMG-CoA reductase. This lowers liver cholesterol, plasma LDL, and the hepatic synthesis of VLDL and apo B. Statins also cause a small increase in HDL, and atorvastatin lowers triglycerides (TGs). The adverse effects are listed.
- \* Cholestyramine and colestipol are bile acid sequestrants that enhance cholesterol loss into the feces, thereby stimulating new bile salt synthesis, which lowers liver cholesterol levels and consequently plasma LDL levels. Their adverse effects are also listed.
- \* Nicotinic acid inhibits the hepatic synthesis of VLDL and apoprotein. It also increases HDL levels and decreases plasma VLDL, LDL, and TG levels. The adverse effects are listed.
- \* Gemfibrozil activates lipoprotein lipase, thus decreasing VLDL, TG, and LDL levels. The adverse effects are listed.
- \* Ezetimibe prevents cholesterol absorption.



# Cardiac and Renal Drug List and Practice Questions

**Table III-8-1. The Major Cardiovascular and Renal Drugs**

Antiarrhythmics	Antihypertensives	Antianginals
IA quinidine, procainamide	$\alpha_2$ agonists: clonidine, methyldopa	Nitrates: nitroglycerin, isosorbide
1B lidocaine	Neuron blockers: reserpine, guanethidine	CCBs: verapamil, nifedipine
1C flecainide	$\alpha$ blockers: prazosin, doxazosin, etc.	$\beta$ blockers: propranolol, etc.
II propranolol, acebutolol (ISA) esmolol	$\beta$ blockers: propranolol, metoprolol, acebutolol, labetalol, etc.	
III amiodarone, sotalol	ACEIs: captopril, etc., and ARBs: losartan, etc. Renin inhibitor: aliskiren	
IV verapamil, diltiazem	Vasodilators: hydralazine, nitroprusside, diazoxide, minoxidil	
Adenosine	CCBs: verapamil, nifedipine, etc.	
	Pulmonary hypertension: bosentan, epoprostenol, sildenafil	
Diuretics	Drugs for Heart Failure	Antihyperlipidemics
CA inhibitors: acetazolamide	Digoxin	Resins: cholestyramine, colestipol
Loops: ethacrynic acid, furosemide	Bipyridines: inamrinone, milrinone; $\beta$ agonists: dobutamine, dopamine	Statins
Thiazides—hydrochlorothiazide, indapamide	$\alpha$ $\beta$ blocker: carvedilol	Other: nicotinic acid, ezetimibe, gemfibrozil
$K^+$ sparing: amiloride, triamterene, spironolactone, eplerenone	Diuretics, vasodilators, nesiritide	Orlistat



### Review Questions

1. A patient has a genetic polymorphism such that they cannot rapidly metabolize drugs by acetylation. You would be most concerned about this polymorphism if the patient was taking which drug?
  - A. Sotalol
  - B. Clonidine
  - C. Nitroglycerin
  - D. Hydralazine
  - E. Prazosin
  
2. Which side effect is associated with spironolactone?
  - A. Alkalosis
  - B. Hirsutism
  - C. Hyperkalemia
  - D. Hypercalcemia
  - E. Hyperglycemia
  
3. Lidocaine is an effective antiarrhythmic because it
  - A. suppresses excitability in hypoxic areas of the heart
  - B. prolongs the QT interval
  - C. prolongs the PR interval
  - D. depresses the slope of phase 0 in slow response tissues
  - E. acts on inhibitory G-protein coupled receptors
  
4. Sildenafil has been prescribed for years to treat erectile dysfunction. Recently, this drug is also being used for what condition?
  - A. vasospastic angine
  - B. supraventricular tachycardia
  - C. cyanide poisoning
  - D. Raynaud disease
  - E. pulmonary hypertension

5. A patient with hypertension also suffers from essential tremor. Optimal treatment of the patient should include management with
  - A. prazosin
  - B. clonidine
  - C. metoprolol
  - D. lidocaine
  - E. propranolol
  
6. Selective  $\beta$ -1 blockers are preferred over nonselective beta blockers in some patients because they
  - A. cause less cardiodepression
  - B. are less likely to cause bronchoconstriction
  - C. are more effective for migraine prophylaxis
  - D. are more effective as antiarrhythmics
  - E. have greater prophylactic value post-MI
  
7. Which drug will utilize the same signaling pathway as endogenous bradykinin on smooth muscle?
  - A. minoxidil
  - B. nitroprusside
  - C. theophylline
  - D. phenylephrine
  - E. cocaine
  
8. A 75-year-old patient suffering from congestive heart failure accidentally ingests a toxic dose of digoxin. Clinical consequences due to the toxic effects of cardiac glycosides are likely to include
  - A. seizures
  - B. hypercalcemia
  - C. bicarbonaturia
  - D. intermittent claudication
  - E. visual disturbances

9. In the management of a cardiac arrhythmia, lidocaine is to be administered by way of an IV loading dose. What variable must be known to calculate an appropriate loading dose?
- A. renal clearance
  - B. bioavailability
  - C. volume of distribution
  - D. lag time
  - E. time to steady-state
10. Both dobutamine and inamrinone increase cardiac contractility by
- A. activation of adenylyl cyclase
  - B. inactivation of Na channels
  - C. inhibition of  $\text{Na}^+/\text{K}^+$ -ATPase
  - D. increasing cAMP
  - E. activation of Na/Cl cotransporter
11. Which one of the following is likely to occur following treatment of a hypercholesterolemic patient with cholestyramine?
- A. Increased recycling of bile salts
  - B. Increased circulating cholesterol
  - C. Decreased VLDL synthesis
  - D. Downregulation of LDL receptors
  - E. Elevation of plasma triglycerides
12. A new diuretic is being studied in human volunteers. Compared with placebo, the new drug increases urine volume, increases urinary  $\text{Ca}^{2+}$ , increases plasma pH, and decreases serum  $\text{K}^+$ . If this new drug has a similar mechanism of action to an established diuretic, it probably
- A. blocks the NaCl cotransporter in the DCT
  - B. blocks aldosterone receptors in the CT
  - C. inhibits carbonic anhydrase in the PCT
  - D. inhibits the  $\text{Na}^+/\text{K}^+/2\text{Cl}^-$  cotransporter in the TAL
  - E. acts as an osmotic diuretic

13. Which one of the following drugs is most likely to block  $K^+$  channels in the heart responsible for cardiac repolarization, and also blocks calcium channels in the AV node?
- A. Amiodarone
  - B. Quinidine
  - C. Lidocaine
  - D. Sotalol
  - E. Verapamil
14. The treatment of hyperlipidemic patients with nicotinic acid (niacin) results in
- A. increases in VLDL
  - B. decreases in both plasma cholesterol and TGs
  - C. inhibition of HMG-CoA reductase
  - D. decreases in HDL
  - E. no change in total cholesterol in the plasma
15. Which one of the following drugs is most likely to cause symptoms of severe depressive disorder when used in the treatment of hypertensive patients?
- A. Captopril
  - B. Hydrochlorothiazide
  - C. Prazosin
  - D. Nifedipine
  - E. Reserpine
16. Enhancement of the effects of bradykinin is most likely to occur with drugs like
- A. clonidine
  - B. diazoxide
  - C. lisinopril
  - D. losartan
  - E. propranolol

17. Outpatient prophylaxis of a patient with an SVT is best accomplished with the administration of
- A. adenosine
  - B. diltiazem
  - C. esmolol
  - D. lidocaine
  - E. mexilitene
18. Which one of the following is the most appropriate drug to use for the patient described in parentheses?
- A. Captopril (60-year-old woman with diabetic nephropathy)
  - B. Nitroprusside (50-year-old man with BP of 140/95)
  - C. Losartan (29-year-old pregnant woman)
  - D. Propranolol (40-year-old patient with peripheral vascular disease)
  - E. Milrinone (57-year-old patient with chronic CHF)
19. In a patient suffering from angina of effort, nitroglycerin may be given sublingually because this mode of administration
- A. bypasses the coronary circulation
  - B. causes less reflex tachycardia than oral administration
  - C. improves patient compliance
  - D. has a decreased tendency to cause methemoglobinemia
  - E. avoids first-pass hepatic metabolism
20. A patient with a supraventricular tachycardia has an atrial rate of 280/min with a ventricular rate of 140/min via a 2:1 AV nodal transmission. After treatment with a drug, the atrial rate slowed to 180/min, but the ventricular rate increased to 180/min! Which of the following drugs was most likely to have been given to this patient?
- A. Adenosine
  - B. Digoxin
  - C. Esmolol
  - D. Quinidine
  - E. Verapamil

## Answers

- Answer: D.** Hydralazine is metabolized by *N*-acetyltransferase (a phase II drug metabolism reaction) associated with a genetic polymorphism. Patients who are classified as slow acetylators may develop SLE-like symptoms when treated with hydralazine. Other drugs metabolized via *N*-acetyltransferase, including isoniazid and procainamide, have also been associated with lupus-like symptoms in slow acetylators.
- Answer: C.** **Spironolactone** blocks aldosterone receptors thereby inhibiting the production of  $\text{Na}^+$  channels in the collecting duct and is used as a  $\text{K}^+$ -sparing agent because the reabsorption of  $\text{Na}^+$  in the CT is coupled (indirectly) to the secretion of  $\text{K}^+$  ions. Hyperkalemia is characteristic of this drug and may lead to clinical consequences at high doses, or if patients fail to discontinue  $\text{K}^+$  supplements or ingest foodstuffs high in  $\text{K}^+$ . Because  $\text{Na}^+$  reabsorption is associated with secretion of protons, spironolactone causes retention of  $\text{H}^+$  ions, leading to acidosis. It has no significant effect on the renal elimination of  $\text{Ca}^{2+}$  or on the plasma level; of glucose.
- Answer: A.** Lidocaine, a class IB drug, effectively targets ischemic areas of the heart. Its major effect is on sodium channels in fast response fibers such as ventricular muscle. It has no significant effect on the PR or QT intervals.
- Answer: E.** Sildenafil (a PDE5 inhibitor) is used for erectile dysfunction but has been recently approved for use in pulmonary hypertension. Other useful drugs in pulmonary hypertension are epoprostenol and bosentan.
- Answer: E.** Propranolol is a nonselective beta blocker useful in a variety of cardiac conditions including hypertension. The drug is also useful in essential tremor where blocking the beta-2 receptor is beneficial. Metoprolol, beta-1 selective, is useful in hypertension but not essential tremor. Clonidine and prazosin are second-line drugs for hypertension and not effective in essential tremor. Lidocaine, an antiarrhythmic, is not effective in either condition.
- Answer: B.**  $\beta_1$ -selective blockers like atenolol and metoprolol are less likely to block receptors in the bronchiolar smooth muscle and therefore less likely to cause bronchoconstriction, especially in asthmatic patients. Nonselective beta blockers are considered to be equally as effective as selective beta-1 blockers in arrhythmias, migraine prevention, and in post-MI prophylaxis. Both types of drugs are cardiodepressant.
- Answer: B.** Bradykinin binds to endothelial receptors and causes the formation of nitric oxide, which signals through the cGMP pathway to relax smooth muscle. Nitroprusside utilizes nitric oxide and cGMP in a similar fashion to relax smooth muscle.
- Answer: E.** Digoxin toxicity is associated with CNS consequences including disorientation and visual dysfunctions such as halos around lights and blurry, yellow vision. More serious manifestations include life-threatening arrhythmias.
- Answer: C.** Back to basic principles! Recall that to calculate a loading dose you must know volume of distribution and target plasma concentration. Since lidocaine is being given IV, its bioavailability is 100% ( $f=1$ ) so no adjustment is required to the equation. Renal clearance is needed to calculate a maintenance dose, and time to steady-state applies only when using a maintenance dose. There is no lag time for an IV drug.

10. **Answer: D.** Dobutamine acts as a beta-1 agonist to activate adenylyl cyclase and increase cAMP. Inamrinone inhibits phosphodiesterase III which increases the amount of cAMP in the heart. In each case, there is an increase in intracellular  $\text{Ca}^{2+}$  being sequestered in the SR which leads to enhance contractility.
11. **Answer: E.** Cholestyramine and colestipol are resins that sequester bile acids in the gut, preventing their reabsorption. This leads to release of their feedback inhibition of 7-alpha hydroxylase and the diversion of cholesterol toward new synthesis of bile acids. Increase in high-affinity LDL receptors on hepatocyte membranes decreases plasma LDL. These drugs have a small but significant effect to increase plasma HDL rather than decrease it, but their ability to increase TGs precludes their clinical use in the management of hypertriglyceridemias.
12. **Answer: D.** The effects described are typical of loop diuretics, which inhibit the  $\text{Na}^+\text{K}^+2\text{Cl}^-$  cotransporter in the thick ascending limb. This action prevents the reabsorption of  $\text{Ca}^{2+}$  from the paracellular pathway and provides for the use of these drugs in hypercalcemia. The increased load of  $\text{Na}^+$  in the collecting tubules leads to increased excretion of both  $\text{K}^+$  and  $\text{H}^+$ , so hypokalemia and alkalosis may occur.
13. **Answer: A.** Amiodarone is a highly effective antiarrhythmic drug, in part because of its multiple actions, which include  $\text{Na}^+$  channel block, beta adrenoceptor block,  $\text{K}^+$  channel block, and  $\text{Ca}^{2+}$  channel block. Drugs that block  $\text{K}^+$  channels prolong APD and ERP and predispose toward torsades de pointes ventricular arrhythmias. Quinidine, class Ia, can block both sodium and potassium channels but not calcium channels. Lidocaine, class Ib, blocks only sodium channels. Sotalol is both a beta blocker and a potassium channel blocker. It is a class III drug that also has class II properties. Verapamil is a class IV calcium channel blocker with no effect on potassium.
14. **Answer: B.** Nicotinic acid inhibits the synthesis of the VLDL apoprotein and decreases VLDL production. Its use results in decreases of both cholesterol and triglycerides, so total cholesterol in the plasma decreases. The drug is not an inhibitor of HMG-CoA reductase, and it increases plasma HDL to a greater extent than any other available antihyperlipidemic drug.
15. **Answer: E.** In addition to decreasing the storage of NE in sympathetic nerve endings, reserpine causes a dose-dependent depletion of brain amines, including NE and serotonin. Symptoms of depression are thought to be related to a functional deficiency in noradrenergic and/or serotonergic neurotransmission in the CNS—the “amine hypothesis of depression.” Although other drugs used in the management of HTN may cause CNS effects, reserpine is the most likely drug to cause severe depression.
16. **Answer: C.** ACE inhibitors prevent the conversion of angiotensin I to angiotensin II and lower blood pressure by decreasing both the formation of aldosterone formation and the vasoconstrictive action of AII at AT-1 receptors. ACEIs also inhibit the metabolism of bradykinin, and this leads to additional hypotensive effects, because bradykinin is an endogenous vasodilator. Unfortunately, increases in bradykinin are associated with side effects, including cough and angioedema. Losartan, which blocks AT-1 receptors, does not increase bradykinin levels.

17. **Answer: B.** Supraventricular tachycardias (SVTs) are treated effectively by class II and class IV antiarrhythmics. In addition, adenosine is indicated for SVTs and nodal tachycardias but only acutely since it must be administered IV and has an extremely short duration. The primary actions of both beta blockers (esmolol) and CCBs (diltiazem) are at the AV node, but esmolol is too short-acting to be useful as prophylaxis. Lidocaine and mexiletine are both class Ib drugs that are used in ventricular arrhythmias.
18. **Answer: E.** ACEIs slow the progression of diabetic nephropathy and are indicated for management of HTN in such patients. Nitroprusside is used IV in severe HTN or hypertensive crisis, not for management of mild-to-moderate HTN. Losartan, which blocks AT-1 receptors, is associated with teratogenic effects during fetal development, as are the ACEIs. Nonselective beta blockers are not ideal for patients who suffer from peripheral vascular disease, diabetes, or asthma. Milrinone, like most inotropes, is not useful long-term in CHF patients. The drug has been shown to increase mortality with chronic use, and thus is indicated for acute CHF. Digoxin is currently the only inotrope used chronically.
19. **Answer: E.** The sublingual administration of a drug avoids its absorption into the portal circulation and hence eliminates the possibility of first-pass metabolism, which can often have a major impact on oral bioavailability. Given sublingually, nitroglycerin is more effectively absorbed into the systemic circulation and has improved effectiveness in angina by this mode of administration. Effective absorption is unlikely to decrease reflex tachycardia or propensity toward methemoglobinemia. There is no bypass of the coronary circulation—nitrates actually decrease coronary vasospasm, which makes them effective in variant angina.
20. **Answer: D.** An increase in AV conduction is characteristic of quinidine, which exerts quite marked blocking actions on muscarinic receptors in the heart. Thus, an atrial rate, formerly transmitted to the ventricles in a 2:1 ratio, may be transmitted in a 1:1 ratio after quinidine. This effect of quinidine can be offset by the prior administration of an antiarrhythmic drug that decreases AV nodal conduction, such as digoxin or verapamil. All of the drugs listed (except quinidine) slow AV nodal conduction, but adenosine and esmolol (a beta blocker) are very short-acting agents used IV only.






**SECTION IV**

**CNS Pharmacology**




# Sedative-Hypnotic-Anxiolytic Drugs

- Drugs: benzodiazepines (BZs), barbiturates, and alcohols



**Figure IV-1-1. CNS Effects Associated with Increasing Doses of Sedative-Hypnotic (S-H) Drugs**

- Cause dose-dependent CNS depression that extends from sedation to anesthesia to respiratory depression and death
- BZs reach a plateau in CNS depression; barbiturates and alcohol do not
- Mechanisms:



5 subunit types:  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\rho$ ,  $\delta$

**Figure IV-1-2. Site of Action of Drugs on the GABA<sub>A</sub> Complex**

**Clinical Correlate**

**Flumazenil**

This nonspecific BZ receptor antagonist is used to reverse the CNS depression caused by BZs used in anesthesia or in BZ overdose. Flumazenil cannot reverse the CNS depression caused by barbiturates and alcohols.

- GABA<sub>A</sub> activation ↑ Cl<sup>-</sup> influx
- GABA<sub>B</sub> activation ↑ K<sup>+</sup> efflux
- Both mechanisms result in membrane hyperpolarization
  - Benzodiazepines:
    - Potentiate GABA
    - ↑ the frequency of Cl<sup>-</sup> channel opening
    - Have no GABA mimetic activity
    - Act through BZ receptors
    - These receptors are part of the GABA<sub>A</sub> complex
    - BZ<sub>1</sub> mediates sedation
    - BZ<sub>2</sub> mediates antianxiety and impairment of cognitive functions
  - Barbiturates:
    - Prolong GABA activity
    - ↑ duration of Cl<sup>-</sup> channel opening
    - Have GABA mimetic activity at high doses
    - Do not act through BZ receptors
    - Have their own binding sites on the GABA<sub>A</sub> complex
    - Also inhibit complex I of electron transport chain
- Uses of BZs:

**Table IV-1-1. Uses of Various Benzodiazepines**

Drug	Indications
Alprazolam	Anxiety, panic, phobias
Diazepam	Anxiety, preop sedation, muscle relaxation, withdrawal states
Lorazepam	Anxiety, preop sedation, status epilepticus (IV)
Midazolam	Preop sedation, anesthesia IV
Temazepam	Sleep disorders
Oxazepam	Sleep disorders, anxiety

- Pharmacokinetics of BZs:
  - Liver metabolized to active compounds, except for oxazepam, temazepam, and lorazepam
- Uses of barbiturates:
  - Phenobarbital is used for seizures
  - Thiopental is used for induction of anesthesia
- Pharmacokinetics of barbiturates:
  - Liver metabolized, sometimes to active compounds

- General inducers of cytochrome P450s
  - Contraindication in porphyrias
- Tolerance to and dependence on sedative-hypnotics:
- Chronic use leads to tolerance
  - Cross-tolerance occurs between BZs, barbiturates, and ethanol
  - Psychologic and physical dependence occur
  - But abuse liability of BZs is < ethanol or barbiturates
  - Withdrawal signs of BZs:
    - Rebound insomnia
    - Anxiety
    - Seizures when BZs were used as antiepileptic or in high doses
  - Withdrawal signs of barbiturates and ethanol:
    - Anxiety
    - Agitation
    - Life-threatening seizures (delirium tremens with alcohol)
  - Management of withdrawal: supportive and long-acting BZs

#### Drug interactions

- GABA<sub>A</sub> drugs are:
  - Additive with other CNS depressants (possible life-threatening respiratory depression), such as anesthetics, antihistamines, opiates,  $\beta$ -blockers, etc.
  - Barbiturates induce metabolism of most lipid-soluble drugs, such as oral contraceptives, carbamazepine, phenytoin, warfarin, etc.

#### Non-BZ drugs:


- Zolpidem and zaleplon
  - BZ<sub>1</sub> receptor agonist
  - Less effect on cognitive function (BZ<sub>2</sub>-mediated)
  - Overdose reversed by flumazenil
  - Used in sleep disorders
  - Less tolerance and abuse liability
- Buspirone
  - No effect on GABA
  - 5-HT<sub>1A</sub> partial agonist
  - Used for generalized anxiety disorders
  - Nonsedative
  - Takes 1 to 2 weeks for effects

### Chapter Summary

- \* Sedative-hypnotic-anxiolytic drugs include the benzodiazepines, barbiturates, and alcohols.
- \* S-H drugs ideally should reduce anxiety without affecting mental or motor function. However, most do affect mental or motor function. Figure IV-1-1 illustrates the relative effects on these functions of classes of S-H drugs at increasing concentrations.
- \* Most S-H drugs facilitate GABA action by binding to the GABA<sub>A</sub> receptor, which has one binding site for barbiturates and alcohol and another for benzodiazepines (Figure IV-1-2). The binding of these drugs at these sites leads to increased Cl<sup>-</sup> influx, potentiating the inhibitory transmitter effects of GABA. The differences in action of the various S-H drugs relate to the differences in the binding site used. Further heterogeneity is introduced by the existence of two subtypes of benzodiazepine receptors, BZ<sub>1</sub> and BZ<sub>2</sub>.
- \* The benzodiazepines are used to treat anxiety states and sleep disorders. Dose-dependent CNS depression does occur but can be reversed by flumazenil. Chronic use can lead to tolerance and dependency with rebound effects upon withdrawal. Table IV-1-1 summarizes the various benzodiazepines and their indications.
- \* Phenobarbital is used to treat seizures, and thiopental is used as an IV anesthetic. Barbiturates induce deep CNS depression at high doses, and there is no antidote.
- \* The barbiturates induce drug-metabolizing enzymes, including the P450 system, leading to potential drug interactions. They also stimulate heme synthesis and are contraindicated in porphyrias.
- \* Tolerance, dependence, and severe withdrawal symptoms are associated with chronic barbiturate use.
- \* Zolpidem and zaleplon are nonbenzodiazepines that bind to the BZ<sub>1</sub> receptors and therefore are more specific hypnotics. Buspirone is an anxiolytic that does not work through the GABA system. It is nonsedating and does not cause dependence but takes a week or two to show antianxiety effects.

# Alcohols

- All alcohols cause CNS depression, in part through GABA mimetic activity.
- All alcohols cause metabolic acidosis.




- 1 CNS depression
- 2 severe metabolic acidosis
- 3 **nephrotoxicity**



- 1 respiratory failure
- 2 severe anion gap metabolic acidosis
- 3 **ocular damage**

Treatment for overdose: fomepizole—long acting inhibitor of alcohol dehydrogenase. High alcohol levels will also require hemodialysis.



- 1 CNS depression
- 2 metabolic acidosis (ketones)
- 3 **acetaldehyde toxicity**

↑ increasing plasma levels

- ↑ sociability
- ↑ gait disturbances
- ↑ reaction time
- ataxia
- impaired motor & mental skills
- impaired memory
- coma
- ↓ death

↑ inhibited by disulfiram

- 1 N&V
- 2 headache
- 3 hypotension
- 4 combines with folate to inactivate it
- 5 combines with thiamine to decrease availability

### Chronic Alcoholism

- 1 hypoglycemia
- 2 fatty liver and lipemia
- 3 muscle wasting (long-term alcoholic, poor food intake)
- 4 gout (lactate competes with urate for excretion)

### Clinical Correlate

#### Alcohol and Pregnancy

The fetal alcohol syndrome is characterized by growth restriction, midfacial hypoplasia, microcephaly, and marked CNS dysfunction, including the frequent occurrence of mental retardation.

### In A Nutshell

Drugs that cause disulfiram-like effects:

- Metronidazole
- Cefoperazone
- Cefotetan
- Chlorpropamide
- Griseofulvin

Figure IV-2-1. Metabolism and Pharmacologic Actions of the Alcohols





# Anticonvulsants



- Seizures result from episodic electrical discharges in cerebral neurons associated with prolonged depolarization, during which sustained, high-frequency, repetitive firing (SHFRF) occurs, followed by prolonged hyperpolarization. The goal of drug management is restoration of normal patterns of electrical activity.
- Mechanisms of action:
  - ↓ axonal conduction by preventing Na<sup>+</sup> influx through fast Na channels—carbamazepine, phenytoin
  - ↑ inhibitory tone by facilitation of GABA-mediated hyperpolarization—barbiturates, benzodiazepines
  - ↓ excitatory effects of glutamic acid—lamotrigine, topiramate (blocks AMPA receptors); felbamate (blocks NMDA receptors)
  - ↓ presynaptic Ca<sup>2+</sup> influx through type-T channels in thalamic neurons—ethosuximide and valproic acid

**Table IV-3-1. Seizure States and Drugs of Choice**

Seizure Type	Drugs of Choice
Partial—simple or complex	Valproic acid, phenytoin, carbamazepine
General—tonic-clonic	Valproic acid, phenytoin, carbamazepine
General—absence	Ethosuximide, valproic acid
Status epilepticus	Lorazepam, diazepam, phenytoin, or fosphenytoin*

\*IV fosphenytoin is more water soluble.

- Primary anticonvulsants
  - **Phenytoin**
    - Blocks axonal Na<sup>+</sup> channels in their inactivate state
    - Prevents seizure propagation
    - Uses: seizure states

- Pharmacokinetics:
  - Variable absorption
  - Nonlinear kinetics
  - Induction of cytochrome P450s
  - Zero-order kinetic of elimination
- Side effects:
  - CNS depression
  - Gingival hyperplasia
  - Hirsutism
  - Osteomalacia (↓ vitamin D)
  - Megaloblastic anemia (↓ folate)
  - Aplastic anemia (check hematology lab results)
- Teratogenicity: cleft lip and palate
- **Carbamazepine**
  - Mechanism identical to phenytoin
  - Uses:
    - Seizure states
    - DOC for trigeminal neuralgia
    - Bipolar disorder
  - Pharmacokinetics: induces cytochrome P450, including its own metabolism
  - Side effects:
    - CNS depression
    - Osteomalacia
    - Megaloblastic anemia
    - Aplastic anemia
    - Exfoliative dermatitis
    - ↑ ADH secretion (dilutional hyponatremia)
  - Teratogenicity:
    - Cleft lip and palate
    - Spina bifida
- **Valproic acid**
  - Mechanism:
    - Similar to phenytoin
    - But also inhibition of GABA transaminase
    - Blockade of T-type  $\text{Ca}^{2+}$  channels
  - Uses:
    - Seizure states
    - Mania of bipolar disorders
    - Migraines
  - Pharmacokinetics: inhibits cytochrome P450s

- Side effects:
  - Hepatotoxicity (from toxic metabolite)
  - Thrombocytopenia
  - Pancreatitis
  - Alopecia
- Teratogenicity: spina bifida
- **Ethosuximide**
  - Mechanism: blockade of T-type  $\text{Ca}^{2+}$  channels in thalamic neurons
  - Use: absence seizures
- General features of anticonvulsant drug use:
  - Anticonvulsants are additive with other CNS depressants
  - Avoid abrupt withdrawal, which may precipitate seizures
  - ↓ efficacy of oral contraceptives via induction of cytochrome P450
  - Phenobarbital is considered safest during pregnancy
- Other anticonvulsant drugs
  - **Felbamate and lamotrigine**
    - Block  $\text{Na}^{+}$  channels and glutamate receptors
    - Used in seizure states (often adjunct therapy)
    - Side effects:
      - Hepatotoxicity and aplastic anemia (felbamate)
      - Stevens-Johnson syndrome (lamotrigine)
  - **Gabapentin**
    - ↑ GABA effects
    - Used in seizure states, neuropathic pain (such as postherpetic neuralgia)

### Chapter Summary

- \* Seizures are caused by episodic electrical discharges in cerebral neurons. These trigger repetitive firing and prolonged hyperpolarization. The goal of drug management is to restore normal electrical patterns. Different classes of drugs do this by acting on different receptor/transmitter systems, which are listed.
- \* Table IV-3-1 summarizes the drugs of choice available to treat each of the several types of seizures.
- \* The mechanisms of action, metabolism, and the adverse effect of the primary anticonvulsant drugs (phenytoin, carbamazepine, ethosuximide, valproic acid, and the barbiturates and benzodiazepines) are discussed.
- \* Anticonvulsive drugs in general have additive depressive effects when used with other depressant drugs, cause a precipitation of seizures upon abrupt withdrawal, and decrease the efficiency of oral contraceptives.
- \* Newer anticonvulsants listed are felbamate, gabapentin, and lamotrigine.



# Drugs Used in Anesthesia

## GENERAL ANESTHETICS

### Inhaled Anesthetics




Figure IV-4-1. Compartmentalization of Anesthetics in the Body

Table IV-4-1. Properties of Specific Inhaled Anesthetics

Anesthetic	MAC Value	Blood-Gas Ratio	CV Effects	Specific Characteristics
Nitrous oxide	104%	0.5	Minimal	Rapid onset and recovery, no metabolism Diffusional hypoxia Spontaneous abortions
Halothane	0.8%	2.3	Sensitizes heart to catecholamines	Malignant hyperthermia, hepatitis, cardiac arrhythmias

Other halogenated "fluranes," e.g., desflurane, sevoflurane, etc.

- Anesthesia protocols include several agents in combinations.
- Inhaled anesthetics have varying potency in proportion to their lipid solubility.
- A MAC (minimal alveolar anesthetic concentration) is defined as the concentration of inhaled anesthetic, as a % of inspired air, at which 50% of patients do not respond to a surgical stimulus.
  - MAC is a measure of potency: ED50.
  - The more lipid soluble the anesthetic, the lower the MAC and the greater the potency.
  - MAC values are additive.
  - MAC values are lower in the elderly and in the presence of opiates or sedative-hypnotics.

- Rates of onset and recovery depend on the blood–gas ratio:
  - The more soluble the anesthetic in the blood, the slower the anesthesia.
  - Anesthetics with high blood–gas ratios are associated with slow onset.
  - Anesthetics with high blood–gas ratios are associated with slow recovery.
  - Anesthetics with low blood–gas ratios have fast onset and recovery.

### Intravenous Anesthetics

- Thiopental
  - Barbiturate used for induction
  - Highly lipid soluble; rapid onset; short-acting due to redistribution
- Midazolam
  - Benzodiazepine used for:
    - Preoperative sedation
    - Anterograde amnesia
    - Induction
    - Outpatient surgery
  - Depresses respiratory function
- Propofol
  - Used for induction and maintenance of anesthesia
  - Antiemetic
  - CNS and cardiac depressant
- Fentanyl
  - Opiate used for induction and maintenance of anesthesia
  - Depresses respiratory function
  - See Opioid Analgesics, chapter 5 in this section
- Ketamine
  - Dissociative anesthetic
  - NMDA-receptor antagonist
  - Induction of anesthesia
  - Emergent delirium, hallucinations
  - Cardiovascular stimulation
  - ↑ intracranial pressure

## LOCAL ANESTHETICS




Figure IV-4-2. Mode of Action of Local Anesthetics

- Local anesthetics provide regional anesthesia.
- Drugs:
  - **Esters:** procaine, cocaine, benzocaine are metabolized by plasma and tissue esterases
  - **Amides:** lidocaine, bupivacaine, mepivacaine are metabolized by liver amidases
- Mechanisms:
  - Nonionized form crosses axonal membrane
  - From within, ionized form blocks the inactivated  $Na^+$  channel
  - Slows recovery and prevents propagation of action potentials
- Nerve fiber sensitivity:
  - Nerve fibers most sensitive to blockade are of smaller diameter and have high firing rates
  - The order of sensitivity is:

type B and C > type  $A_\delta$  > type  $A_\beta$  and  $A_\gamma$  > type  $A_\alpha$

- Recovery is in reverse order
- Absorption:
  - Coadministration of  $\alpha_1$  agonists:
    - $\downarrow$  local anesthetic absorption into the systemic circulation
    - Prolong effects and  $\downarrow$  toxicity
- Side effects:
  - Neurotoxicity
  - Cardiovascular toxicity
  - Allergies (esters via PABA formation)

### Note

#### $Na^+$ Channel Toxins

- Tetrodotoxin (from puffer fish) and saxitoxin (algae toxin, "red tide")
  - Block activated  $Na^+$  channels
  - $\downarrow Na^+$  influx
- Ciguatoxin (exotic fish) and batrachotoxin (frogs)
  - Bind to activated  $Na^+$  channels
  - Cause inactivation
  - Prolong  $Na^+$  influx

### Note

#### Esters and Amides

Local anesthetics that are esters have just one "i" in their names (e.g., procaine, cocaine); amide local anesthetics have more than one "i" (e.g., lidocaine, bupivacaine).

### Note

Cocaine intrinsically causes vasoconstriction by blocking norepinephrine uptake.



## SKELETAL MUSCLE RELAXANTS




Figure IV-4-3. Nicotinic ACh Receptor of the Neuromuscular Junction

- Nicotinic receptors have five subunits.
- Two ACh bind each to two  $\alpha$  subunits in order to open the  $\text{Na}^+$  channel.
- This depolarizes the muscle.
- Used mainly in anesthesia protocols or in the ICU to afford muscle relaxation and/or immobility.
- Muscle relaxants interact with nicotinic ACh receptors at the neuromuscular junction.
- Drugs:
  - **Nondepolarizing (competitive)**
    - Nicotinic antagonists
    - D-Tubocurarine prototype
    - Reversible with AChE inhibitors
    - Progressive paralysis (face, limbs, respiratory muscle)
    - No effects on cardiac and smooth muscle
    - No CNS effects
    - Specific drugs:
      - Atracurium**
        - \* Rapid recovery
        - \* Safe in hepatic or renal impairment
        - \* Spontaneous inactivation to laudanosine
        - \* Laudanosine can cause seizures
      - Mivacurium**
        - \* Very short duration
        - \* Metabolized by plasma cholinesterases
  - **Depolarizing (noncompetitive)**
    - Nicotinic agonist
    - Specific drug: **succinylcholine**
    - Two phases:
      - Phase I:* depolarization, fasciculation, prolong depolarization, flaccid paralysis
      - Phase II:* desensitization
    - AChE inhibitors  $\uparrow$  phase I; have no effect on phase II
    - Rapidly hydrolyzed by pseudocholinesterase: short duration

- Cautions:
  - Atypical pseudocholinesterase
  - Hyperkalemia
  - Malignant hyperthermia

### **Centrally Acting Skeletal Muscle Relaxants**

- Benzodiazepines through GABA<sub>A</sub> receptors
- Baclofen through GABA<sub>B</sub> receptors
- Use: spasticity

### **Chapter Summary**

#### **Drugs Used in Anesthesia**

- \* The more lipid soluble the inhalation anesthetic, the greater its potency (lower MAC value). The more soluble an inhalation anesthetic in the blood (higher blood:gas ratio), the slower will be the onset to anesthesia and the slower will be the recovery.
- \* Thiopental, midazolam, propofol, fentanyl, and ketamine are intravenous anesthetics that are discussed.
- \* Local anesthetics (weak bases) infiltrate and anesthetize nerve bundles near sites of injection by binding to inactive Na<sup>+</sup> channels in their ionized forms. However, to get to the channel they must diffuse through the lipid bilayer in an unionized form. Thus, their effects are influenced by pH.
- \* The smaller and most rapidly firing nerve fibers are the most sensitive to blockade.
- \* The coadministration of alpha adrenoceptor agonists decreases local anesthetic absorption into the systemic circulation, prolonging their effects and potentially decreasing their toxicity.
- \* The adverse effects of local anesthetics are given.

#### **Sodium Channel Toxins**

- \* Tetrodotoxin, saxitoxin, ciguatoxin, and batrachotoxin are sodium-channel toxins found in various fish, frogs, or dinoflagellates.

#### **Skeletal Muscle Relaxants**

- \* The skeletal muscle relaxants provide muscle relaxation and/or immobility via N-receptor interactions. Most, including D-tubocurarine, atracurium, and mivacurium, are competitive and nondepolarizing and can be reversed by AChE inhibitors. Succinylcholine is a depolarizing, noncompetitive agonist.
- \* Spasmolytics reduce excess muscle tone or spasm in injury or CNS dysfunction. They may act in the CNS, the spinal cord, or directly on the muscle. Benzodiazepines and baclofen reduce the tonic output of spinal motor neurons. Dantrolene blocks Ca<sup>2+</sup> release from the muscle sarcoplasmic reticulum.

### **Bridge to Pathology/Genetics**

#### **Malignant Hyperthermia**

A life-threatening syndrome characterized by muscle rigidity, hyperthermia, hypertension, acidosis, and hyperkalemia. Associated with the use of skeletal muscle relaxants, especially succinylcholine, used in anesthesia regimens. Genotypic susceptibility may be related to mutations in the genes encoding ryanodine receptors and/or a protein component of the L-type calcium channel in skeletal muscle.

#### **Treatment**

Dantrolene acts directly on skeletal muscle to decrease contractility by blocking Ca<sup>2+</sup> release from the sarcoplasmic reticulum. It is used in states that include extreme muscle rigidity, such as malignant hyperthermia associated with inhaled anesthetics and skeletal muscle relaxants or neuroleptic malignant syndrome associated with antipsychotics.



# Opioid Analgesics

---

- Endogenous opiate peptides represented by endorphins, enkephalins, and dynorphins
- Three receptor families:  $\mu$ ,  $\kappa$ , and  $\delta$
- Presynaptic and postsynaptic inhibition through  $G_i$  coupling
- Mu pharmacology most important
- Morphine is the prototype  $\mu$  agonist
- Pharmacology of **morphine**:
  - Analgesia:  $\uparrow$  pain tolerance and  $\downarrow$  perception and reaction to pain
  - Sedation
  - Respiratory depression:  $\downarrow$  response to  $\uparrow$   $pCO_2$  (do not give  $O_2$ ; give naloxone)
  - Cardiovascular: minimal effects on heart, but vasodilation (avoid in head trauma)
  - Smooth muscle
    - Longitudinal relaxes
    - Circular constricts
      - GI:  $\downarrow$  peristalsis, constipation, cramping
      - GU: urinary retention, urgency to void
      - Biliary:  $\uparrow$  pressure
      - Pupils: miosis
  - Cough suppression: antitussive action, independent of analgesia and respiratory depression
  - Nausea and vomiting: stimulation of the chemoreceptor trigger zone (CTZ) in the area postrema
  - $\uparrow$  histamine release
- Pharmacokinetics of morphine:
  - Glucuronidation
  - Morphine-6-glucuronide is highly active
  - Caution in renal dysfunction
- Other opioids and analgesics (*see* Table IV-5-1).

## Clinical Correlate

### Contraindications and Cautions for Opioids

- Head injuries (possible increased intracranial pressure)
- Pulmonary dysfunction (except pulmonary edema)
- Hepatic/renal dysfunction (possible accumulation)
- Adrenal or thyroid deficiencies (exaggerated responses)
- Pregnancy (possible neonatal depression or dependence), except meperidine

Table IV-5-1. Other Opioids and Analgesics

Receptor Action	Drug	Characteristics
Full agonists	Meperidine	<ul style="list-style-type: none"> <li>• Also antimuscarinic</li> <li>• No miosis</li> <li>• Tachycardia</li> <li>• No spasm GI/GU/gallbladder</li> <li>• Metabolized by cytochrome P450 to normeperidine, a serotonin reuptake inhibitor; normeperidine may cause serotonin syndrome and seizures</li> </ul>
	Methadone	<ul style="list-style-type: none"> <li>• Used in maintenance of opiate addict</li> </ul>
	Codeine	<ul style="list-style-type: none"> <li>• Cough suppressant</li> <li>• Analgesia</li> <li>• Used in combination with NSAIDs</li> </ul>
Partial agonist	Buprenorphine	<ul style="list-style-type: none"> <li>• Precipitation of withdrawal</li> </ul>
Mixed agonist-antagonists	Nalbuphine, pentazocine	<ul style="list-style-type: none"> <li>• <math>\kappa</math> agonist</li> <li>• spinal analgesia</li> <li>• dysphoria</li> <li>• <math>\mu</math> antagonist</li> <li>• precipitation of withdrawal</li> </ul>
Antagonists	Naloxone	<ul style="list-style-type: none"> <li>• IV, reversal for respiratory depression</li> </ul>
	Naltrexone	<ul style="list-style-type: none"> <li>• PO, ↓ craving for alcohol and used in opiate addiction</li> </ul>
	Methylnaltrexone	<ul style="list-style-type: none"> <li>• Treatment of opioid-induced constipation (does not cross BBB and won't precipitate withdrawal)</li> </ul>

- Side effects of opioid analgesics:
  - Acute toxicity: classic triad
    - Pinpoint pupils
    - Respiratory depression
    - Coma
  - Management of acute toxicity:
    - Supportive
    - IV naloxone
- Abuse liability of opioid analgesics:
  - Tolerance: pharmacodynamic; occurs to all effects, except miosis and constipation
  - Dependence: physical and psychologic
  - Withdrawal:
    - Yawning
    - Lacrimation, rhinorrhea, salivation
    - Anxiety, sweating, goose bumps
    - Muscle cramps, spasms, CNS-originating pain

- Management of withdrawal:
  - Supportive
  - Methadone
  - Clonidine
- Opiate-related drugs with specific indications
  - Loperamide: diarrhea
  - Dextromethorphan: cough

### Chapter Summary

- \* Opioid agonists and/or antagonists act in part by binding to the receptors for the endogenous opiopeptides. These are G-protein-linked, multisubunit structures to which the various opioids bind as full or partial agonists or as antagonists. The resultant complex array of potential mechanisms, sites of action, types of effects, kinetics, and contraindications are discussed. Table IV-5-1 summarizes the receptor actions and other relevant characteristics of eight opioid drugs.



# Drugs Used in Parkinson Disease and Psychosis

---

## DOPAMINERGIC NEURAL PATHWAYS

In the CNS, dopamine (DA) is a precursor to NE in diffuse noradrenergic pathways and is an inhibitory neurotransmitter in the following major dopaminergic pathways:

- Nigrostriatal tract
  - Cell bodies in the substantia nigra project to the striatum, where they release DA, which inhibits GABA-ergic neurons. In Parkinson disease, the loss of DA neurons in this tract leads to excessive ACh activity → extrapyramidal dysfunction.
  - DA receptor antagonists → pseudo-Parkinsonism (reversible).
  - DA agonists may cause dyskinesias.
- Mesolimbic-mesocortical tracts—cell bodies in midbrain project to cerebrocortical and limbic structures.
  - Functions include regulation of affect, reinforcement, cognitive functions, and sensory perception. Psychotic disorders and addiction are partly explained by ↑ DA in these pathways.
  - Drugs that ↑ DA functions → ↑ reinforcement and, at high doses, may cause psychoses.
  - DA antagonists → ↓ cognitive function.
- Tuberoinfundibular
  - Cell bodies in hypothalamus project to anterior pituitary and release DA → ↓ prolactin.
  - DA agonists are used in hyperprolactinemic states.
  - DA antagonists may cause endocrine dysfunction, including gynecomastia and amenorrhea/galactorrhea.
- Chemoreceptor trigger zone
  - Activation of DA receptors → ↑ emesis.
  - DA agonists (e.g., apomorphine) are emetic, and DA antagonists are antiemetic.

## DOPAMINE RECEPTORS

- D<sub>1</sub>-like: G<sub>s</sub> coupled
- D<sub>2</sub>-like: G<sub>i</sub> coupled
  - D<sub>2A</sub>: nigrostriatal
  - D<sub>2C</sub>: mesolimbic



## DRUGS USED IN PARKINSON DISEASE

- Signs and symptoms of Parkinson disease include:
  - Bradykinesia
  - Muscle rigidity
  - Resting tremor
- Pathology: degeneration of nigrostriatal dopamine tracts with imbalance between dopamine (↓) and ACh (↑)




Figure IV-6-1. CNS Targets for Antiparkinsonian Drugs

- Pharmacologic strategy: restore normal dopamine and ↓ ACh activity at muscarinic receptors in the striatum
- Drugs increasing dopamine function:
  - **Levodopa**
    - Prodrug converted to dopamine by aromatic amino acid decarboxylase (AAAD)
    - Usually given with carbidopa
    - Side effects:
      - Dyskinesias
      - “On-off” effects
      - Psychosis
      - Hypotension
      - Vomiting




Figure IV-6-2. Inhibitors of Levodopa Metabolism

- **Tolcapone and entacapone**
  - COMT converts dopamine to 3-O-methyl-dopa, a partial agonist at dopamine receptors.
  - These drugs inhibit COMT and enhance levodopa uptake and efficacy.
  - Tolcapone is hepatotoxic.
- **Selegiline**
  - MAO<sub>B</sub>-selective inhibitor (no tyramine interactions)
  - Initial treatment and adjunct to levodopa
  - Side effects: dyskinesias, psychosis, insomnia (metabolized to amphetamine)
- Dopamine-receptor agonists:
  - **Bromocriptine**
    - Use: hyperprolactinemia and acromegaly
    - Side effects: dyskinesias and psychosis
  - **Pramipexole and ropinirole**
- Drugs decreasing ACh function:
  - Include **benztropine, trihexyphenidyl, and diphenhydramine**, which are muscarinic blockers
  - Actions: ↓ tremor and rigidity but have little effects on bradykinesia
  - Side effects: atropine-like
- **Amantadine**
  - Antiviral, which block muscarinic receptors and ↑ dopamine release
  - Side effects: atropine-like and livedo reticularis

## ANTIPSYCHOTIC DRUGS

### Schizophrenia

- Positive symptoms:
  - Thought disorders
  - Delusions
  - Hallucinations
  - Paranoia
- Negative symptoms:
  - Amotivation
  - Social withdrawal
  - Flat affect
  - Poverty of speech
- “Dopamine hypothesis”:
  - Symptoms arise because of excessive dopaminergic activity in mesolimbic system.
  - Dopamine agonists cause psychosis.
  - Dopamine antagonists have antipsychotic actions.
- Serotonin is increasingly seen as a part of the etiology of schizophrenia.
- Mechanism: blockade of dopamine and/or 5HT<sub>2</sub> receptors
- Uses
  - Schizophrenia
  - Schizoaffective states
  - Bipolar disorder
  - Tourette syndrome and Huntington disease
  - Drug or radiation emesis
- Side effects from dopamine blockade:
  - Dyskinesias (extrapyramidal symptoms [EPS])
    - Acute EPS:
      - Pseudoparkinsonism, dystonia, akathisia
      - Management: antimuscarinic drugs (benztropine or diphenhydramine)
    - Chronic EPS:
      - Tardive dyskinesia (TD)
      - Management: discontinuation/switch to atypical
  - Dysphoria
  - Endocrine dysfunction:
    - Temperature regulation problems (neuroleptic malignant syndrome [NMS], treated with dantrolene and bromocriptine)
    - ↑ prolactin (galactorrhea, amenorrhea, gynecomastia)
    - ↑ eating behaviors (weight gain)
- Side effects from muscarinic blockade (particularly tachycardia and ↓ seizure threshold)
- Side effects from alpha blockade (particularly hypotension)

Summary of Antipsychotic Drug Pharmacology

Table IV-6-1. Characteristic Properties of Antipsychotic Drugs

Drug Group Examples	EPS*	M Block	Sedation	Alpha Block	Other Characteristics
<b>TYPICALS</b>					
Chlorpromazine	++	++	+++	+++	NA
Thioridazine	+	+++	+++	+++	<ul style="list-style-type: none"> <li>• Cardiotoxicity (torsades—"quinidine-like")</li> <li>• Retinal deposits</li> </ul>
Fluphenazine	+++	+	+	+	NA
Haloperidol	+++	+	+	+	Most likely cause of neuroleptic malignant syndrome (NMS) and TD
<b>ATYPICALS</b>					
Clozapine	+/-	++	+	+++	<ul style="list-style-type: none"> <li>• Blocks D<sub>2c</sub> and 5HT<sub>2</sub> receptors</li> <li>• No TD</li> <li>• Agranulocytosis—(weekly WBC count) requirement for weekly blood test, weight gain</li> <li>• Increased salivation ("wet pillow" syndrome)</li> <li>• Seizures</li> </ul>
Olanzapine	+/-	+	+	++	Blocks 5HT <sub>2</sub> receptors, improves negative symptoms
Risperidone	+	+/-	++	++	Blocks 5HT <sub>2</sub> receptors, improves negative symptoms
Aripiprazole	+	+/-	+/-	+/-	Partial agonist of D <sub>2</sub> receptor; blocks 5HT <sub>2</sub> receptors

Other atypicals: Quetiapine, Ziprasidone

\*Extrapyramidal symptoms

**Clinical Correlate**

**Parenteral Forms**

Parenteral formulations of certain antipsychotic drugs (e.g., fluphenazine, haloperidol) are available for rapid initiation of treatment and for maintenance therapy in noncompliant patients. Depot forms of both drugs exist.

## Chapter Summary

### Dopaminergic Neural Pathways

- \* Dopamine (DA) in the nigrostriatal tract helps regulate kinesis by inhibiting GABA-ergic and cholinergic neurons. The loss of DA neurons in this tract leads to excessive ACh activity and Parkinsonism. DA receptor antagonists cause a reversible pseudo-Parkinsonism; agonists may cause dyskinesia.
- \* DA neurons in the midbrain projecting into the cerebrocortical and limbic regions regulate affect, reinforcement, psychomotor function, and sensory perception. DA agonists enhance psychomotor activity and reinforcement and at high doses may cause psychoses. DA antagonists decrease psychomotor function.
- \* In the hypothalamus, DA released into the pituitary decreases prolactin release. DA agonists (e.g., pergolide) are used to treat hyperprolactinemia; antagonists may cause endocrine dysfunction.
- \* The activation of DA receptors in the chemoreceptor trigger zone increases emesis; thus, DA agonists are emetic, and antagonists are antiemetic.

### Antiparkinsonian Drugs

- \* Parkinsonism is due to an imbalance between DA and ACh activity in the nigrostriatal tract. Drugs attempt to restore this balance either by increasing DA or decreasing ACh levels. Figure IV-6-1 illustrates the CNS sites targeted in antiparkinsonism therapy.
- \* Drugs used to increase DA function are levodopa, tolcapone, entacapone, bromocriptine, pramipexole, and selegiline. Drugs that decrease ACh function are benztropine, trihexyphenidyl, diphenhydramine, and amantadine. The properties of each are described.

### Antipsychotic Drugs

- \* Although the prevailing concept is that schizophrenia is due to hyperdopaminergic activity in the CNS, not all antischizophrenic drugs act as DA antagonists; some instead modify serotonin function.
- \* The typical antipsychotic drugs (e.g., chlorpromazine, thioridazine, fluphenazine, and haloperidol) act primarily as DA antagonists, blocking  $D_{2A}$  receptors. Side effects include the induction of pseudo-Parkinsonism, akathisia, and/or acute dystonic effects. Their use and symptom management are discussed, as are other adverse effects including toxicity, tardive dyskinesia, and neuroleptic malignant syndrome.
- \* Atypical antipsychotics (e.g., clozapine, risperidone, and olanzapine) act as antagonists at  $5HT_2$  receptors and seem to have fewer adverse effects. Aripiprazole is a  $D_2$  partial agonist.
- \* Table IV-6-1 summarizes the characteristics of the antipsychotic drugs.

# Drugs Used for Depression, Bipolar Disorders, and Attention Deficit Hyperactivity Disorder (ADHD)

---

- “Amine hypothesis” of depression:
  - Reserpine: depletes NE, 5HT, DA, and causes severe depression
  - Acute mechanism of antidepressants:  $\uparrow$  NE,  $\uparrow$  5HT
  - However, antidepressant effect takes several weeks to occur.

## DRUGS USED IN DEPRESSION

### MAO Inhibitors

- Drugs: **phenelzine** and **tranylcypromine**
- Mechanism: inhibition of MAO<sub>A</sub> and MAO<sub>B</sub>
- Use: atypical depressions
- Drug interactions
  - $\uparrow$  NE: hypertensive crisis
    - Symptoms:  $\uparrow$  BP, arrhythmias, excitation, hyperthermia
    - Drugs: releasers (i.e., tyramine), tricyclic antidepressants (TCAs),  $\alpha_1$  agonists, levodopa
  - $\uparrow$  5HT: serotonin syndrome
    - Symptoms: sweating, rigidity, myoclonus, hyperthermia, ANS instability, seizures
    - Drugs: SSRIs, TCAs, and meperidine

### Tricyclic Antidepressants (TCAs)

- Drugs: **amitriptyline**, **imipramine**, and **clomipramine**
- Mechanism: nonspecific blockade of 5HT and NE reuptake
- Uses:
  - Major depressions
  - Phobic and panic anxiety states
  - Obsessive-compulsive disorders (OCDs)
  - Neuropathic pain
  - Enuresis
- Side effects: muscarinic and  $\alpha$  blockade
- Toxicity: the “3 Cs”: coma, convulsions, and cardiotoxicity
- Drug interactions:
  - Hypertensive crisis with MAO inhibitors
  - Serotonin syndrome with SSRIs, MAO inhibitors, and meperidine
  - Prevent antihypertensive action of  $\alpha_2$  agonists and guanethidine

### Selective Serotonin Reuptake Inhibitors (SSRIs)

- Drugs: **fluoxetine, paroxetine, sertraline**
- Mechanism: selective blockade of 5HT reuptake
- Uses:
  - Major depression
  - OCD
  - Bulimia
  - Anxiety disorders
  - Premenstrual dysphoric disorder (PMDD)
- Side effects: anxiety, agitation, bruxism, sexual dysfunction, weight loss
- Toxicity: serotonin syndrome
- Drug interactions: serotonin syndrome with MAO inhibitors, meperidine, and TCAs

### **Clinical Correlate**

Varenicline is a partial agonist of nicotinic receptors and is used in smoking cessation.

### Other Antidepressants

- Trazodone: associated with cardiac arrhythmias and priapism
- Venlafaxine: nonselective reuptake blocker devoid of ANS side effects
- **Bupropion**: dopamine reuptake blocker; used in smoking cessation
- Mirtazapine:  $\alpha_2$  antagonist, associated with weight gain

## **LITHIUM AND BIPOLAR DISORDERS**

- **Lithium** remains DOC for bipolar disorders.
- Usually antidepressants/antipsychotics also required
- Mechanism:
  - Prevents recycling of inositol ( $\downarrow$  PIP<sub>2</sub>)
  - $\downarrow$  cAMP
- Side effects:
  - Narrow therapeutic index; requires therapeutic monitoring
  - Tremor, flu-like symptoms, life-threatening seizures
  - Hypothyroidism with goiter ( $\downarrow$  TSH effects and inhibition of 5'-deiodinase)
  - Nephrogenic diabetes insipidus ( $\downarrow$  ADH effect); manage with amiloride
- Teratogenicity: Ebstein's anomaly (malformed tricuspid valve)
- Other drugs used in bipolar disorders: valproic acid, carbamazepine

## **DRUGS USED IN ADHD**

- **Methylphenidate**: amphetamine-like
  - Side effects: agitation, restlessness, insomnia, cardiovascular toxicity
- **Atomoxetine**: selective NE reuptake inhibitor
  - Side effects: See TCA section, above.

### **Chapter Summary**

- \* The amine hypothesis of depression postulates that symptoms are caused by a functional deficiency of CNS NE and/or 5HT. This is based on the observation that most antidepressants affect the metabolism of these amines. Again, there are exceptions.
- \* The uses, drug interactions, and adverse effects of the monoamine oxidase inhibitors, tricyclic antidepressants, selective serotonin reuptake inhibitors, and other antidepressants are discussed.
- \* Lithium, the mainstay for bipolar disorder treatment, often needs supplementation with antidepressant and/or sedative drugs. The uses, mechanisms of action, and adverse effects of lithium therapy as well as backup drugs used for treatment of bipolar disorder are considered.
- \* Atomoxetine and methylphenidate are used in the treatment of ADHD.





# Drugs of Abuse

## DRUGS OF ABUSE

Table IV-8-1. Properties of Drugs of Abuse

CNS Stimulants	Cocaine	Amphetamines
Neurotransmitters involved	NE, DA, 5HT	
Mechanism(s) of action	Blocks DA, NE, and 5HT reuptake in CNS; local anesthetic action from Na <sup>+</sup> channel blockade	Blockade of reuptake of NE and DA, release amines from mobile pool, weak MAO inhibitors
Effects	<ol style="list-style-type: none"> <li>1. Increase NE: sympathomimetic effect with increased heart rate and contractility, blood pressure changes, mydriasis, and central excitation, hyperactivity</li> <li>2. Increase DA: psychotic episodes, paranoia, hallucinations, possible dyskinesias, and endocrine disturbances</li> <li>3. Increase 5HT: behavioral changes, aggressiveness, dyskinesias, and decreased appetite</li> </ol>	
Toxicity	<ol style="list-style-type: none"> <li>1. Excess NE: cardiac arrhythmias, generalized ischemia with possible MI and strokes; acute renal and hepatic failures</li> <li>2. Excess DA: major psychosis, cocaine delirium</li> <li>3. Excess 5HT: possible serotonin syndrome</li> <li>4. All of the above: convulsion, hyperpyrexia, and death</li> </ol>	
Withdrawal	Craving, severe depression, anhedonia, anxiety; manage with antidepressants	
CNS Depressants	Benzodiazepines	Barbiturates and Ethanol
Neurotransmitters involved	GABA	
Mechanism of action	Potentiation of GABA interaction with GABA <sub>A</sub> receptors involves BZ <sub>1</sub> and BZ <sub>2</sub> binding sites	Prolongation of GABA, GABA mimetic at high doses, on GABA <sub>A</sub> receptors
Effects	Light to moderate CNS depression	Any plane of CNS depression

(Continued)

**Table IV-8-1. Properties of Drugs of Abuse (continued)**

Toxicity	Sedation, anterograde amnesia; in severe OD (or IV use), reverse with <b>flumazenil</b>	Severe CNS depression, respiratory depression, and death
Withdrawal	Rebound insomnia, rebound anxiety	Agitation, anxiety, hyperreflexia, and life-threatening seizures + in ethanol withdrawal delusions/ hallucinations—delirium tremens (DTs)

Opioids	Morphine, Heroin, Methadone, Fentanyl, Other Opioids	
Neurotransmitters involved	NE, DA, 5HT, GABA, and many others	
Mechanism of action	Activate opioid $\mu$ , $\kappa$ , and $\delta$ receptors. Potent $\mu$ receptor activators have the most intense abuse and dependence liability, possibly effected via an increase in dopaminergic transmission in the mesolimbic tracts	
Effects	Euphoria, analgesia, sedation, cough suppression, and constipation; strong miosis (except meperidine)	
Toxicity	Severe respiratory depression (reverse with <b>naloxone</b> ), nausea, vomiting	
Withdrawal	Lacrimation, yawning, sweating, and restlessness, rapidly followed with centrally originating pain, muscle cramping, and diarrhea; not life-threatening	

Hallucinogens	Marijuana	Hallucinogens
Neurotransmitters involved	Many	5HT
Mechanism of action	Interaction of THC with $CB_1$ and $CB_2$ cannabinoid receptors in CNS and periphery	Interaction with several subtypes of 5HT <sub>1</sub> receptors
Effects	Sedation, euphoria, $\uparrow$ HR, conjunctiva, delusions, hallucinations	Hallucinogen, sympathomimetic, causes dysesthesias
Toxicity	Associated with smoking, possible flashbacks	Poorly described, flashbacks likely
Withdrawal	Irritability, anxiety	Poorly characterized

**Miscellaneous Abused Drugs**

1. PCP: NMDA-receptor antagonist; extremely toxic, horizontal and vertical nystagmus, paranoia, rhabdomyolysis; overdose is common, with convulsions and death
2. Ketamine: similar to but milder than PCP, with hallucinations, glutamate-receptor antagonist
3. Anticholinergics: scopolamine, atropine-like
4. MDMA (“Ecstasy”), MDA, MDEA: amphetamine-like with strong 5HT pharmacology and therefore hallucinogenic; generally neurotoxic
5. Inhalants: solvent abuse, multiple organ damage; see Toxicology, section XI

### Chapter Summary

- \* Table IV-8-1 summarizes the properties of drugs of abuse. These include the CNS stimulants (cocaine and amphetamines), the CNS depressants (benzodiazepines, barbiturates, and ethanol), the opioids (morphine, heroin, methadone, fentanyl, and others), the hallucinogens (marijuana and other hallucinogens), PCP, ketamine, anticholinergics (scopolamine), MDMA-MDA-MDEA (all amphetamine-like), and inhalants.



# CNS Drug List and Practice Questions

Table IV-9-1. CNS Drug List

<b>Sedative-Hypnotics</b>	<b>Anticonvulsants</b>
Barbiturates: phenobarbital Benzodiazepines: alprazolam, diazepam, lorazepam, oxazepam	Carbamazepine, ethosuximide, valproic acid, phenytoin, diazepam, lorazepam, gabapentin, lamotrigine, felbamate, topiramate, tiagabin, vigabatrin
Others: buspirone, zolpidem, zaleplon BZ receptor antagonist: flumazenil	<b>Anesthetics (Inhaled)</b>
	Halothane, nitrous oxide
<b>Anesthetics (IV)</b>	<b>Neuromuscular Blocking Agents</b>
Fentanyl, ketamine, midazolam, propofol, thiopental	Depolarizing: succinylcholine Nondepolarizing: atracurium, mivacurium, tubocurarine
<b>Local Anesthetics</b>	<b>Skeletal Muscle Relaxants</b>
Lidocaine, bupivacaine, mepivacaine, procaine, cocaine	Depolarizing: succinylcholine Nondepolarizing: D-tubocurarine, atracurium, mivacurium
<b>Opioid Analgesics</b>	<b>Antipsychotics</b>
Full agonists: morphine, meperidine, methadone, fentanyl, and heroin Partial agonists: buprenorphine, codeine Mixed agonist-antagonists: nalbuphine Antagonists: naloxone, naltrexone, methylnaltrexone	Typicals: Chlorpromazine, fluphenazine, thioridazine, haloperidol Atypicals: clozapine, risperidone, olanzapine, aripiprazole, quetiapine, ziprasidone
<b>Antiparkinsonian Drugs</b>	<b>Antidepressants</b>
DA agonists: levodopa, bromocriptine, pramipexole MAO-B inhibitor: selegiline AAAD inhibitor: carbidopa M blockers: benztropine, trihexiphenidyl COMT inhibitor: tolcapone Amantadine	MAOIs: phenelzine, tranylcypromine TCAs: amitriptyline, imipramine, clomipramine SSRIs: fluoxetine, paroxetine, sertraline Others: bupropion, mirtazapine, trazodone, venlafaxine
<b>Bipolar Disorder</b>	<b>ADHD</b>
Lithium	Methylphenydate Atomoxetine

### Review Questions

1. Lorazepam can be safely used as a preanesthetic medication in a patient undergoing liver transplantation without fear of excessive CNS depression because the drug is
  - A. excreted in unchanged form
  - B. actively secreted into the GI tract
  - C. conjugated extrahepatically
  - D. a selective anxiolytic devoid of CNS depressant actions
  - E. reversible by naloxone
2. Midazolam is an effective anesthetic because it acts by
  - A. increasing functional activity at GABA<sub>B</sub> receptors
  - B. enhancing the actions of dopamine
  - C. blocking the NMDA glutamate receptor subtype
  - D. acting as a partial agonist at 5HT receptors
  - E. facilitating GABA-mediated increases in chloride ion conductance
3. Which one of the following is an established clinical use of morphine?
  - A. Management of generalized anxiety disorders
  - B. Relief of pain associated with biliary colic
  - C. Pulmonary congestion
  - D. Treatment of cough associated with use of ACE inhibitors
  - E. Suppression of the ethanol withdrawal syndrome
4. A 40-year-old man was given a drug that binds to a subunit of the GABA<sub>A</sub> receptor. When used at a high dose, the drug can open Cl<sup>-</sup> channels independent of GABA. What drug was the man given?
  - A. Diazepam
  - B. Ethanol
  - C. Phenobarbital
  - D. Baclofen
  - E. Dronabinol
5. Which one of the following is characteristic of both phenytoin and carbamazepine?
  - A. Inhibition of hepatic cytochrome P450
  - B. First-order elimination at high therapeutic doses
  - C. Enhances the effects of oral contraceptives
  - D. Safe to use in pregnancy
  - E. Prevent sodium influx through fast sodium channels

6. A patient comes to the ER with a painful stab wound. The ER resident administers pentazocine for the pain. Soon after administration the patient experiences sweating, restlessness, and an increase in pain sensations. What is the most likely explanation for his symptoms?
- A. The patient is probably tolerant to pentazocine.
  - B. The patient is a heroin addict.
  - C. Pentazocine is an ineffective analgesic.
  - D. Pentazocine was used at the wrong dose.
  - E. Pentazocine doesn't cross the blood-brain barrier.
7. The data shown in the table below concern the effects of drugs on transmitter function in the CNS. Which one of the drugs is most likely to alleviate extrapyramidal dysfunction caused by typical antipsychotics? (The + signs denote intensity of drug actions.)

Drug	Activation of DA Receptors	Activation of GABA Receptors	Block of ACh M Receptors
A.	++++	0	0
B.	++	++	0
C.	0	0	++++
D.	0	+++++	0
E.	+	+	0

8. Tricyclic antidepressants
- A. increase the antihypertensive effect of guanethidine
  - B. have anticonvulsant activity
  - C. should not be used in patients with glaucoma
  - D. may increase oral absorption of levodopa
  - E. are sometimes used as antiarrhythmics
9. Which one of the following statements about lithium is accurate?
- A. It causes symptoms of mild hyperthyroidism in up to 25% of patients.
  - B. Plasma levels are increased by a high-Na diet.
  - C. Adverse effects include acne, polydipsia, and polyuria.
  - D. Spina bifida is major concern in fetal development.
  - E. Sedative actions calm manic patients within 24 h.
10. Ingestion of methanol in wood spirits would cause which of the following to happen?
- A. The formation of formaldehyde
  - B. Nephrotoxicity
  - C. Hypotension and vomiting
  - D. The production of glycolic acids
  - E. Inhibition of aldehyde dehydrogenase



11. What is the rationale for combining levodopa with carbidopa?
- Carbidopa stimulates dopamine receptors
  - Carbidopa increases levodopa entry into the CNS by inhibiting peripheral dopa decarboxylase
  - Carbidopa enhances levodopa absorption
  - Carbidopa enhances the peripheral conversion of levodopa to dopamine
  - Carbidopa blocks peripheral COMT
12. A 29-year-old male patient is being treated with an antidepressant drug, and his mood is improving. However, he complains of feeling “jittery” and agitated at times, and if he takes his medication in the afternoon he finds it difficult to get to sleep at night. He seems to have lost weight during the 6 months that he has been taking the drug. He has been warned not to take other drugs without consultation because severe reactions have occurred with opioid analgesics including meperidine. This patient is probably taking
- alprazolam
  - chlorpromazine
  - paroxetine
  - amitriptyline
  - trazodone
13. The ability of several drugs to inhibit the reuptake of CNS amine neurotransmitters is shown in the table below (number of arrows ↓ indicates the intensity of inhibitory actions). Which one of the drugs is most likely to have therapeutic effectiveness in the management of both obsessive-compulsive disorders (OCD) and major depressive disorders?

Drug	DA Reuptake	NE Reuptake	5HT Reuptake	GABA Reuptake
A.	↓↓	0	0	↓↓
B.	0	↓↓↓↓	↓	0
C.	0	0	↓↓↓↓	0
D.	0	0	↓	↓↓↓↓
E.	↓↓↓↓	↓↓	0	0

14. A patient suffering from attention deficit hyperactivity disorder is placed on atomoxetine. A drug that has a similar mechanism of action to atomoxetine is
- methylphenidate
  - botulinum toxin
  - clonidine
  - amitriptyline
  - entacapone

15. A patient suffering from generalized anxiety disorder (GAD) has a history of drug dependence that includes the illicit use of secobarbital (“reds”) and a variety of other drugs. Psychotherapy is indicated, but the physician also prescribes a drug that can be helpful in GAD and that has the advantage of no abuse liability. The drug prescribed was most likely to have been
- A. bupropion
  - B. buspirone
  - C. baclofen
  - D. buprenorphine
  - E. phenobarbital
16. A patient has been diagnosed as having “long QT syndrome.” The patient is experiencing significant pain following a bout with shingles. What would be an appropriate drug for his pain?
- A. Amitriptyline
  - B. Fentanyl
  - C. Acyclovir
  - D. Diazepam
  - E. Gabapentin
17. A habitual user of a schedule-controlled drug abruptly stops using it. Within 8 h, she becomes anxious, starts to sweat, and gets severe abdominal pain with diarrhea. These symptoms intensify over the next 12 h, during which time she has a runny nose, is lacrimating, and has uncontrollable yawning and intensification of muscle cramping and jerking. Assuming that these are withdrawal symptoms in the patient due to her physical dependence, the drug most likely to be involved is
- A. alprazolam
  - B. amphetamine
  - C. ethanol
  - D. meperidine
  - E. secobarbital
18. A 57-year-old patient, living at home, has severe pain due to a metastatic carcinoma that is being managed with fentanyl, delivered transdermally from a patch. He should also be taking, or at least have on hand
- A. apomorphine
  - B. docusate
  - C. loperamide
  - D. morphine
  - E. naloxone

19. A hospital nurse is taking imipramine for a phobic anxiety disorder, and her patient is being treated with chlorpromazine for a psychotic disorder. Which of the following adverse effects is likely to occur in both of these individuals?
- A. Excessive salivation
  - B. Pupillary constriction
  - C. Orthostatic hypotension
  - D. Seizure threshold
  - E. Weight loss
20. Which one of the following pairs of “drug/mechanism of action” is most accurate?
- A. Carbamazepine/facilitation of the actions of GABA
  - B. Ethosuximide/blocks Na channels in axonal membranes
  - C. Phenelzine/inhibits dopa decarboxylase
  - D. Procaine/blocks Ca channels (type T) in thalamic neurons
  - E. Lithium/inhibits recycling of inositol
21. A 30-year-old male patient is brought to the ER with the following symptoms attributed to a drug overdose: HR and BP, mydriasis, behavioral excitation, aggressiveness, paranoia, and hallucinations. Of the following drugs, which one is most likely to be responsible for these symptoms?
- A. Amphetamine
  - B. Ethanol
  - C. Fentanyl
  - D. Flunitrazepam
  - E. Marijuana
22. Which one of the following CNS receptors is directly coupled to an ion channel so that the effects of its activation do not involve second messenger systems?
- A. N (ACh)
  - B.  $\alpha$  (NE)
  - C. D<sub>2A</sub> (DA)
  - D.  $\mu$  (beta endorphin)
  - E. 5HT<sub>2</sub> (serotonin)

## Answers

- Answer: C.** Most benzodiazepines are metabolized by liver cytochrome P450. In a patient lacking liver function, benzodiazepines that are metabolized via extrahepatic conjugation (e.g., lorazepam, oxazepam) are safer in terms of the possibility of excessive CNS depression. Lorazepam is metabolized, probably in the lungs, via glucuronidation. Although benzodiazepine actions can be reversed, the drug that acts as an antagonist is flumazenil, not naloxone.
- Answer: E.** Benzodiazepines interact with components of the GABA receptor–chloride ion channel macromolecular complex. Binding of BZs leads to an increase in the frequency of chloride ion channel opening elicited by the inhibitory transmitter GABA. Benzodiazepines do not act on GABA<sub>B</sub> receptors; baclofen, a centrally acting muscle relaxant, is an agonist at these receptors. Buspirone, the selective anxiolytic, may be a partial agonist at 5HT receptors.
- Answer: C.** Morphine continues to be used in pulmonary congestion, in part because of its sedative (calming) and analgesic effects and also because of its vasodilating actions, which result in favorable hemodynamics in terms of cardiac and pulmonary function. Similarly, morphine is of value in an acute MI, especially its ability to relieve pain. However, morphine is not suitable for pain of biliary origin because it causes contraction of the sphincters of Oddi, leading to spasms. None of the other proposed indications are appropriate.
- Answer: C.** Benzodiazepines, barbiturates, and ethanol all modulate the actions of the GABA<sub>A</sub> receptor, while baclofen works at the GABA<sub>B</sub> receptor, and dronabinol works on cannabinoid receptors. Of the GABA<sub>A</sub> drugs, only barbiturates have GABA-mimicking activity and this occurs at high doses. This is one of the reasons why barbiturates are a more dangerous group of drugs than benzodiazepines since benzos lack GABA-mimicking activity.
- Answer: E.** Phenytoin has the unusual characteristic of following first-order elimination kinetics at low doses but zero-order kinetics at high doses because of saturation of the liver enzymes involved in its metabolism. Carbamazepine, like most drugs, follows first-order kinetics. Both drugs are P450 inducers and can increase the metabolism of oral contraceptives making them less effective. Both drugs are teratogenic, causing structural abnormalities during fetal development including cleft palate. Both drugs block inactivated sodium channels, preventing sodium entry, thereby prolonging the time to recovery.
- Answer: B.** Pentazocine is an agonist at κ (kappa) opioid receptors and an antagonist at μ opioid receptors. Mixed agonist-antagonists can displace μ receptor agonists such as heroin from receptors, resulting in the rapid development of symptoms of withdrawal in patients who are physically dependent on such drugs—“precipitated withdrawal.” Symptoms include yawning, lacrimation, salivation, restlessness, anxiety, sweating, goosebumps, muscle cramps, and pain.
- Answer: C.** Muscarinic receptor antagonists such as benztropine, trihexy-phenidyl, and diphenhydramine are used to manage the reversible extrapyramidal dysfunction (e.g., pseudo-Parkinsonism) that results from treatment with drugs that block DA receptors in the striatum (typical antipsychotics). Drugs that activate DA receptors, although theoretically possible, require doses that are toxic and exacerbate psychoses. Because the actions of DA in the striatum lead to inhibition of GABA-ergic neurons, drugs that activate GABA receptors are unlikely to be effective in this situation, although they may well have both anxiolytic and anticonvulsant properties.

8. **Answer: C.** In addition to blocking reuptake of NE and 5HT, pharmacodynamic actions of the tricyclic antidepressants include block of peripheral adrenergic and muscarinic receptors—the former resulting in postural hypotension and the latter, via mydriasis, exacerbating glaucoma. TCAs block the uptake of guanethidine into sympathetic nerve endings, decreasing its antihypertensive effects, and they may cause arrhythmias in overdose. They have no effect on the absorption of levodopa.
9. **Answer: C.** Lithium causes goiter in a significant number of patients; however, thyroid dysfunction does not occur in all such patients, and when it does it presents as hypothyroidism (not hyper-T). High-Na diets increase lithium elimination; low Na increases lithium plasma levels. Uncoupling of vasopressin receptors is characteristic of lithium, leading to a nephrogenic diabetes insipidus. Although potential teratogenicity is a concern during pregnancy, lithium does not cause neural tube defects but may cause abnormalities in heart valves. Lithium takes 10 to 20 days for effectiveness, and in acute mania it is often necessary to calm the patient with parenteral antipsychotic drugs such as fluphenazine or haloperidol.
10. **Answer: A.** Methanol is metabolized by alcohol dehydrogenase to formaldehyde and then further metabolized to formic acid by aldehyde dehydrogenase. Its major toxicity is severe vision damage. Ethylene glycol ingestion is associated with nephrotoxicity, while ethanol ingestion causes nausea, vomiting, and hypotension.
11. **Answer: B.** Carbidopa inhibits peripheral dopa decarboxylase which enhances uptake of levodopa into the CNS and therefore, its conversion to dopamine. Carbidopa doesn't cross the blood-brain barrier and therefore has no direct benefit at dopamine receptors.
12. **Answer: C.** The patient is probably taking an SSRI such as paroxetine. SSRIs rarely cause sedation and commonly cause agitation and the "jitters," which sometimes necessitates concomitant use of drugs that are strongly sedating, such as trazodone. SSRIs are best taken in the morning to avoid problems of insomnia, and they appear to cause weight loss, at least during the first 12 months of treatment. Severe drug interactions leading to the "serotonin syndrome" have been reported when SSRIs have been used together with MAO inhibitors, tricyclics, and the opioid meperidine.
13. **Answer: C.** Drug C appears to be a selective inhibitor of the reuptake of serotonin, and existing drugs of this class (SSRIs) are approved for use in both major depressive and obsessive-compulsive disorders. The tricyclic antidepressant clomipramine, a potent inhibitor of 5HT reuptake, was formerly the drug of choice for OCD until replaced by the SSRIs. Drugs A and E may have value in the treatment of Parkinson disease because they block the reuptake of DA. Drug D may be effective in anxiety and seizure states because it is an effective blocker of GABA reuptake.
14. **Answer: D.** Atomoxetine is used in attention deficit hyperactivity disorder (ADHD) and works by blocking the reuptake of norepinephrine into nerve terminals. This mechanism is how both cocaine and the tricyclic antidepressants such as amitriptyline work. Amphetamines such as methylphenidate are also commonly used in ADHD and work by displacing norepinephrine from the mobile pool.

15. **Answer: B.** Buspirone has selective anxiolytic activity that is slow in onset. The drug has no abuse liability and will not suppress withdrawal symptoms in patients who have become physically dependent on barbiturates, benzodiazepines, or ethanol. Bupropion is an antidepressant, also approved for management of dependence on nicotine. Baclofen is a spinal cord muscle relaxant that activates GABA<sub>B</sub> receptors. Buprenorphine is a long-acting opioid analgesic with no effectiveness in GAD, and phenobarbital is a barbiturate that may cause dependence.
16. **Answer: E.** The patient is experiencing postherpetic neuralgia. While acyclovir is effective at eradicating the herpes virus it is ineffective against the pain of shingles. Appropriate drugs are TCAs like amitriptyline and gabapentin. Patients with long QT syndrome have a genetic flaw in cardiac inward rectifying K current, leading to increased APD. Drugs that accentuate this by inhibiting the repolarizing K current (phase 3), which include thioridazine and the tricyclic antidepressants, are likely to have enhanced cardiotoxic potential in such patients. As a result, this patient should be placed on gabapentin.
17. **Answer: D.** The signs and symptoms described are typical of withdrawal from physical dependency on an opioid that has efficacy equivalent to a full agonist—in this case, meperidine. Although anxiety, agitation, and even muscle jerking may occur in withdrawal from dependence on sedative-hypnotics such as alprazolam and secobarbital, the symptoms of GI distress, rhinorrhea, lacrimation, and yawning are not characteristic (seizures are more typical). Symptoms of withdrawal from high-dose use of CNS stimulants such as amphetamine or cocaine include lassitude and severe depression of mood. The phrase “schedule-controlled” refers to FDA classifications of drugs that have abuse liability, including both licit and illicit drugs.
18. **Answer: B.** Fentanyl is a full agonist at opioid receptors and provides analgesia in cancer pain equivalent to morphine, so there is no good reason to have morphine on hand, and it would be a danger to the patient in terms of accidental overdose. Apomorphine is an emetic, hardly appropriate given the stimulatory effects of opioids on the emetic center. Likewise, loperamide is used in diarrheal states, and patients on strong opioids are almost certain to be constipated; for this reason, a stool softener like docusate should be available to the patient. The opioid antagonist naloxone is used IV in overdose situations but would not be provided to the patient for use PRN.
19. **Answer: C.** Orthostatic hypotension occurs with both tricyclic antidepressants and phenothiazines because both types of drug can block alpha-adrenergic receptors in venous beds. Their ability to block M receptors leads to xerostomia (not salivation) and mydriasis (not miosis). Tricyclics and phenothiazines also share a common tendency to decrease seizure threshold and cause weight gain (not loss).
20. **Answer: E.** Lithium inhibits the dephosphorylation of IP<sub>2</sub> (needed for the recycling of inositol), leading to depletion of membrane PIP<sub>2</sub>. Consequently, the activation of receptors by neurotransmitters such as ACh, NE, and 5HT fails to release the second messengers IP<sub>3</sub> and DAG. Carbamazepine and the local anesthetic procaine block axonal Na channels; ethosuximide may block Ca channels in thalamic neurons. Phenelzine is a nonselective inhibitor of MAO.

21. **Answer: A.** The signs and symptoms are characteristic of a CNS stimulant that facilitates the activity of amines in both the CNS and the periphery. Amphetamines promote the release of NE from sympathetic nerve endings, causing CV stimulation and pupillary dilation. In the CNS, they enhance the actions of DA, NE, and 5HT, causing behavioral excitation and a psychotic state that may be difficult to distinguish from schizophrenia. Ethanol, marijuana, fentanyl, and flunitrazepam (a benzodiazepine that has been used in “date rape”) are all CNS depressants.
22. **Answer: A.** ACh receptors in the CNS are present on less than 5% of the neuronal population. Most of them are of the muscarinic subtype,  $M_1$  (excitatory) and  $M_2$  (inhibitory), via G-protein coupled changes in cAMP. Nicotinic receptors are excitatory via direct coupling to cation channels (Na/K), and their activation does not initiate second messenger pathways. Other CNS transmitter receptors that are directly coupled to ion channels include those for GABA and glutamic acid. Almost all CNS receptors for DA, NE, 5HT, and opioid peptides are coupled to ion channels via second messenger systems.

SECTION V

# **Antimicrobial Agents**





# Antibacterial Agents

---

## PRINCIPLES OF ANTIMICROBIAL CHEMOTHERAPY

- Bactericidal
- Bacteriostatic
- Combinations:
  - Additive
  - Synergistic (penicillins plus aminoglycosides)
  - Antagonistic (penicillin plus tetracyclines)
- Mechanisms:

**Table V-1-1. Mechanism of Action of Antimicrobial Agents**

Mechanism of Action	Antimicrobial Agents
Inhibition of bacterial cell-wall synthesis	Penicillins, cephalosporins, imipenem/meropenem, aztreonam, vancomycin
Inhibition of bacterial protein synthesis	Aminoglycosides, chloramphenicol, macrolides, tetracyclines, streptogramins, linezolid
Inhibition of nucleic synthesis	Fluoroquinolones, rifampin
Inhibition of folic acid synthesis	Sulfonamides, trimethoprim, pyrimethamine

- Resistance:

Table V-1-2. Mechanisms of Resistance to Antimicrobial Agents

Antimicrobial Agents	Primary Mechanism(s) of Resistance
Penicillins and cephalosporins	Production of beta-lactamases, which cleave the beta-lactam ring structure; change in penicillin-binding proteins; change in porins
Aminoglycosides (gentamicin, streptomycin, amikacin, etc.)	Formation of enzymes that inactivate drugs via conjugation reactions that transfer acetyl, phosphoryl, or adenylyl groups
Macrolides (erythromycin, azithromycin, clarithromycin, etc.) and clindamycin	Formation of methyltransferases that alter drug binding sites on the 50S ribosomal subunit Active transport out of cells
Tetracyclines	Increased activity of transport systems that “pump” drugs out of the cell
Sulfonamides	Change in sensitivity to inhibition of target enzyme; increased formation of PABA; use of exogenous folic acid
Fluoroquinolones	Change in sensitivity to inhibition of target enzymes; increased activity of transport systems that promote drug efflux
Chloramphenicol	Formation of inactivating acetyltransferases

## INHIBITORS OF CELL-WALL SYNTHESIS

- All cell-wall synthesis inhibitors are bactericidal.




Figure V-1-1. Beta-Lactam Antibiotics

## Penicillins

- Mechanisms of action:
  - Bacterial cell wall is cross-linked polymer of polysaccharides and pentapeptides
  - Penicillins interact with cytoplasmic membrane-binding proteins (PBPs) to inhibit transpeptidation reactions involved in cross-linking, the final steps in cell-wall synthesis
- Mechanisms of resistance:
  - Penicillinases (beta-lactamases) break lactam ring structure (e.g., staphylococci)
  - Structural change in PBPs (e.g., methicillin-resistant *Staphylococcus aureus* [MRSA], penicillin-resistant pneumococci)
  - Change in porin structure (e.g., *Pseudomonas*)
- Subgroups and antimicrobial activity:
  - Narrow spectrum, beta-lactamase sensitive: penicillin G and penicillin V
    - Spectrum: streptococci, pneumococci, meningococci, *Treponema pallidum*
  - Very narrow spectrum, beta-lactamase resistant: nafcillin, methicillin, oxacillin
    - Spectrum: known or suspected staphylococci (not MRSA)
  - Broad spectrum, aminopenicillins, beta-lactamase sensitive: ampicillin and amoxicillin
    - Spectrum: gram-positive cocci (not staph), *E. coli*, *H. influenzae*, *Listeria monocytogenes* (ampicillin), *Borrelia burgdorferi* (amoxicillin), *H. pylori* (amoxicillin)
  - Extended spectrum, antipseudomonal, beta-lactamase sensitive: ticarcillin, piperacillin, azlocillin
    - Spectrum: increased activity against gram-negative rods, including *Pseudomonas aeruginosa*
- General considerations:
  - Activity enhanced if used in combination with beta-lactamase inhibitors (clavulanic acid, sulbactam)
  - Synergy with aminoglycosides against pseudomonal and enterococcal species
- Pharmacokinetics:
  - Most are eliminated via active tubular secretion with secretion blocked by probenecid; dose reduction needed only in major renal dysfunction
  - Nafcillin and oxacillin eliminated largely in bile; ampicillin undergoes enterohepatic cycling, but excreted by the kidney
  - Benzathine penicillin G—repository form (half-life of 2 weeks)
- Side effects:
  - Hypersensitivity
    - Incidence 5 to 7% with wide range of reactions (types I–IV). Urticarial skin rash common, but severe reactions, including anaphylaxis, are possible.
    - Interstitial nephritis with methicillin
    - Assume complete cross-allergenicity between individual penicillins
  - Other:
    - GI distress (NVD), especially ampicillin
    - Jarisch-Herxheimer reaction in treatment of syphilis

### Bridge to Biochemistry

#### Suicide Inhibitors

Metabolism of a substrate by an enzyme to form a compound that irreversibly inhibits that enzyme. Penicillinase inhibitors, such as clavulanic acid and sulbactam, are suicide inhibitors.

### Bridge to Immunology

#### Drug Hypersensitivity Reactions

- I. IgE mediated—rapid onset; anaphylaxis, angioedema, laryngospasm
- II. IgM and IgG antibodies fixed to cells—vasculitis, neutropenia, positive Coombs test
- III. Immune complex formation—vasculitis, serum sickness, interstitial nephritis
- IV. T-cell mediated—urticarial and maculopapular rashes, Stevens-Johnson syndrome

## Cephalosporins

- Mechanisms of action and resistance: identical to penicillins
- Subgroups and antimicrobial activity:
  - First generation: cefazolin, cephalexin
    - Spectrum: gram-positive cocci (not MRSA), *E. coli*, *Klebsiella pneumoniae*, and some *Proteus* species
    - Common use in surgical prophylaxis
    - Pharmacokinetics: none enter CNS
  - Second generation: cefotetan, cefaclor, cefuroxime
    - Spectrum: ↑ gram-negative coverage, including some anaerobes
    - Pharmacokinetics: no drugs enter the CNS, except cefuroxime
  - Third generation: ceftriaxone (IM) and cefotaxime (parenteral), cefdinir and cefixime (oral)
    - Spectrum: gram-positive and gram-negative cocci, plus many gram-negative rods
    - Pharmacokinetics: most enter CNS (not cefoperazone); important in empiric management of meningitis and sepsis
  - Fourth generation: cefepime (IV)
    - Even wider spectrum
    - Resistant to most beta-lactamases
    - Enters CNS
- Pharmacokinetics:
  - Renal clearance similar to penicillins, with active tubular secretion blocked by probenecid
  - Dose modification in renal dysfunction
  - Cefoperazone and ceftriaxone are largely eliminated in the bile
- Side effects:
  - Hypersensitivity:
    - Incidence: 2%
    - Wide range, but rashes and drug fever most common
    - Positive Coombs test, but rarely hemolysis
    - Assume complete cross-allergenicity between individual cephalosporins and partial cross-allergenicity with penicillins (about 5%)
    - Most authorities recommend avoiding cephalosporins in patients allergic to penicillins (for gram-positive organisms, consider macrolides; for gram-negative rods, consider aztreonam)
  - Other:
    - Disulfiram-like effect: cefotetan and cefoperazone

### Classic Clues

Organisms *not* covered by cephalosporins are “LAME”:

*Listeria monocytogenes*

Atypicals (e.g., *Chlamydia*, *Mycoplasma*)

MRSA

Enterococci

### Imipenem and Meropenem

- Mechanism of action:
  - Same as penicillins and cephalosporins
  - Resistant to beta-lactamases
- Spectrum:
  - Gram-positive cocci, gram-negative rods (e.g., *Enterobacter*, *Pseudomonas* spp.), and anaerobes
  - Important in-hospital agents for empiric use in severe life-threatening infections
- Pharmacokinetics:
  - Imipenem is given with cilastatin, a renal dehydropeptidase inhibitor, which inhibits imipenem's metabolism to a nephrotoxic metabolite
  - Both drugs undergo renal elimination—↓ dose in renal dysfunction
- Side effects:
  - GI distress
  - Drug fever (partial cross-allergenicity with penicillins)
  - CNS effects, including seizures with imipenem in overdose or renal dysfunction

### Aztreonam

- Mechanism of action:
  - Same as for penicillins and cephalosporins
  - Resistant to beta-lactamases
- Uses:
  - IV drug mainly active versus gram-negative rods
  - No cross-allergenicity with penicillins or cephalosporins

### Vancomycin

- Mechanism of action:
  - Binding at the D-ala-D-ala muramyl pentapeptide to sterically hinder the transglycosylation reactions involved in elongation of peptidoglycan chains
  - Does not interfere with PBPs
- Spectrum:
  - MRSA
  - Enterococci
  - *Clostridium difficile* (backup drug)
- Resistance:
  - Vancomycin-resistant staphylococcal (VRSA) and enterococcal (VRE) strains emerging
  - Enterococcal resistance involves change in the muramyl pentapeptide “target,” such that the terminal D-ala is replaced by D-lactate
- Pharmacokinetics:
  - Used IV and orally (not absorbed) in colitis
  - Enters most tissues (e.g., bone), but not CNS
  - Eliminated by renal filtration (important to decrease dose in renal dysfunction)

- Side effects:
  - “Red man syndrome” (histamine release)
  - Ototoxicity (usually permanent, additive with other drugs)
  - Nephrotoxicity (mild, but additive with other drugs)

## INHIBITORS OF BACTERIAL PROTEIN SYNTHESIS

- Site of action:




Figure V-1-2. Bacterial Protein Synthesis

- Mechanisms:

Table V-1-3. Summary of Mechanisms of Protein Synthesis Inhibition

Event	Antibiotic(s) and Binding Site(s)	Mechanism(s)
1. Formation of initiation complex	Aminoglycosides (30S) Linezolid (50S)	Interfere with initiation codon functions—block association of 50S ribosomal subunit with mRNA-30S (static); misreading of code (aminoglycosides only)—incorporation of wrong amino acid (–cidal)
2. Amino-acid incorporation	Tetracyclines (30S) Dalfopristin/ quinupristin (50S)	Block the attachment of aminoacyl tRNA to acceptor site (–static)
3. Formation of peptide bond	Chloramphenicol (50S)	Inhibit the activity of peptidyltransferase (–static)
4. Translocation	Macrolides and clindamycin (50S)	Inhibit translocation of peptidyl-tRNA from acceptor to donor site (–static)

- For mechanisms of resistance of antibiotics, see Table V-1-2.

## Aminoglycosides

- Activity and clinical uses:
  - Bactericidal, accumulated intracellularly in microorganisms via an O<sub>2</sub>-dependent uptake → anaerobes are innately resistant
  - Useful spectrum includes gram-negative rods; **gentamicin, tobramycin, and amikacin** often used in combinations
  - Synergistic actions occur for infections caused by enterococci (with penicillin G or ampicillin) and *P. aeruginosa* (with an extended-spectrum penicillin or third-generation cephalosporin)
  - **Streptomycin** used in tuberculosis; is the DOC for bubonic plague and tularemia
- Pharmacokinetics:
  - Are polar compounds, not absorbed orally or widely distributed into tissues
  - Renal elimination proportional to GFR, and major dose reduction needed in renal dysfunction
- Side effects:
  - Nephrotoxicity (6 to 7% incidence) includes proteinuria, hypokalemia, acidosis, and acute tubular necrosis—usually reversible, but enhanced by vancomycin, amphotericin B, cisplatin, and cyclosporine
  - Ototoxicity (2% incidence) from hair cell damage; includes deafness (irreversible) and vestibular dysfunction (reversible); toxicity may be enhanced by loop diuretics

## Bridge to Microbiology

### Once-Daily Dosing of Aminoglycosides

Antibacterial effects depend mainly on peak drug level (rather than time) and continue with blood levels < MIC—a postantibiotic effect (PAE).

Toxicity depends both on blood level and the time that such levels are > than a specific threshold (i.e., total dose).



- Neuromuscular blockade with ↓ release of ACh—may enhance effects of skeletal muscle relaxants

### Tetracyclines

- Activity and clinical uses:
  - Bacteriostatic drugs, actively taken up by susceptible bacteria
  - “Broad-spectrum” antibiotics, with good activity versus chlamydial and mycoplasmal species, *H. pylori* (GI ulcers), *Rickettsia*, *Borrelia burgdorferi*, *Brucella*, *Vibrio*, and *Treponema* (backup drug)
- Specific drugs:
  - **Doxycycline**: more activity overall than tetracycline HCl and has particular usefulness in prostatitis because it reaches high levels in prostatic fluid
  - **Minocycline**: in saliva and tears at high concentrations and used in the meningococcal carrier state
  - **Demeclocycline**: used in syndrome of inappropriate secretion of ADH (SIADH; blocks ADH receptor function in collecting ducts)
  - Tigecycline: used in complicated skin, soft tissue, and intestinal infections due to resistant gram + (MRSA, VREF), gram –, and anaerobes
- Pharmacokinetics:
  - Kidney for most (↓ dose in renal dysfunction)
  - Liver for doxycycline
  - Chelators: tetracyclines bind divalent cations ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Fe}^{2+}$ ), which ↓ their absorption
- Side effects:
  - Tooth enamel dysplasia and possible ↓ bone growth in children (avoid)
  - Phototoxicity (demeclocycline, doxycycline)
  - GI distress (NVD), superinfections leading to candidiasis or colitis
  - Vestibular dysfunction (minocycline)
  - Have caused liver dysfunction during pregnancy at very high doses (contraindicated)

### **Classic Clues**

#### **Phototoxicity**

- Tetracyclines
- Sulfonamides
- Quinolones

### Chloramphenicol

- Activity and clinical uses:
  - Bacteriostatic with a wide spectrum of activity
  - Currently a backup drug for infections due to *Salmonella typhi*, *B. fragilis*, *Rickettsia*, and possibly in bacterial meningitis
- Pharmacokinetics:
  - Orally effective, with good tissue distribution, including CSF
  - Metabolized by hepatic glucuronidation, and dose reductions are needed in liver dysfunction and in neonates
  - Inhibition of cytochrome P450
- Side effects:
  - Dose-dependent bone marrow suppression common; aplastic anemia rare (1 in 35,000)
  - “Gray baby” syndrome in neonates (↓ glucuronosyl transferase)

## Macrolides

- Drugs: erythromycin, azithromycin, clarithromycin
- Activity and clinical uses:
  - Macrolides are wide-spectrum antibiotics
    - Gram-positive cocci (not MRSA)
    - Atypical organisms (*Chlamydia*, *Mycoplasma*, and *Ureaplasma* species)
    - *Legionella pneumophila*
    - *Campylobacter jejuni*
    - *Mycobacterium avium-intracellulare* (MAC)
    - *H. pylori*
- Pharmacokinetics:
  - They inhibit cytochrome P450s
- Side effects:
  - Macrolides stimulate motilin receptors and cause gastrointestinal distress (erythromycin, azithromycin > clarithromycin)
  - Macrolides cause reversible deafness at high doses
- Telithromycin: a ketolide active against macrolide-resistant *S. pneumonia*

## Clindamycin

- Not a macrolide, but has the same mechanisms of action and resistance
- Narrow spectrum: gram-positive cocci (including community-acquired MRSA) and anaerobes, including *B. fragilis* (backup drug)
- Concentration in bone has clinical value in osteomyelitis due to gram-positive cocci
- Side effect: pseudomembranous colitis (most likely cause)

## Linezolid

- Mechanism of action:
  - Inhibits the formation of the initiation complex in bacterial translation systems by preventing formation of the N-formylmethionyl-tRNA-ribosome-mRNA ternary complex
- Spectrum:
  - Treatment of VRSA, VRE, and drug-resistant pneumococci
- Side effects: bone marrow suppression (platelets)

## Quinupristin–Dalfopristin

- Mechanism of action:
  - Quinupristin and dalfopristin—streptogramins that act in concert via several mechanisms
  - Binding to sites on 50S ribosomal subunit, they prevent the interaction of aminoacyl-tRNA with acceptor site and stimulate its dissociation from ternary complex
  - May also decrease the release of completed polypeptide by blocking its extrusion

## Bridge to Microbiology

### Community-Acquired Pneumonia

With no comorbidity, the most common organisms associated with community-acquired pneumonia are *M. pneumoniae*, *C. pneumoniae*, and viruses. In smokers, the pneumococcus is a more frequent pathogen. Macrolide antibiotics have activity against most strains of these organisms (other than viruses) and are therefore commonly used in the treatment of a community-acquired pneumonia.

## Clinical Correlate

### Don't Use in Pregnancy

Aminoglycosides,  
fluoroquinolones,  
sulfonamides, tetracyclines

**Note**

- Streptogramins for *E. faecium*, including VRE *faecium*, but not for *E. faecalis*
- Linezolid for both types of enterococci

- Spectrum:
  - Used parenterally in severe infections caused by vancomycin-resistant staphylococci (VRSA) and enterococci (VRE), as well as other drug-resistant, gram-positive cocci
- Side effects:
  - Toxic potential remains to be established

**INHIBITORS OF NUCLEIC ACID SYNTHESIS**

**Inhibitors of Folic Acid Synthesis**

- Drugs: sulfonamides, trimethoprim, and pyrimethamine

**Bridge to Biochemistry**

**Antimetabolites**

*Definition:* a substance inhibiting cell growth by competing with, or substituting for, a natural substrate in an enzymatic process

Sulfonamides and trimethoprim are antimetabolites, as are many antiviral agents and drugs used in cancer chemotherapy.





Figure V-1-3. Inhibitors of Folic Acid Synthesis

- Activity and clinical uses:
  - Sulfonamides alone are limited in use because of multiple resistance
  - Sulfasalazine is a prodrug used in ulcerative colitis and rheumatoid arthritis
  - Ag sulfadiazine used in burns



5-ASA = 5-aminosalicylic acid; SP = sulfapyridine

Figure V-1-4. Metabolism and Uses of Sulfasalazine

- Combination with dihydrofolate reductase inhibitors:
  - ↓ resistance
  - Synergy
- Uses of trimethoprim-sulfamethoxazole (cotrimoxazole):
  - Bacteria:
    - DOC in *Nocardia*
    - Listeria (backup)
    - Gram-negative infections (*E. coli*, *Salmonella*, *Shigella*, *H. influenzae*)
    - Gram-positive infections (*Staph.*, including community-acquired MRSA, *Strep.*)
  - Fungus: *Pneumocystis jirovecii* (back-up drugs are pentamidine and atovaquone)
  - Protozoa: *Toxoplasma gondii* (sulfadiazine + pyrimethamine)
- Pharmacokinetics:
  - Sulfonamides are hepatically acetylated (conjugation)
  - Renally excreted metabolites cause crystalluria (older drugs)
  - High protein binding
    - Drug interaction
    - Kernicterus in neonates (avoid in third trimester)
- Side effects:
  - Sulfonamides
    - Hypersensitivity (rashes, Stevens-Johnson syndrome)
    - Hemolysis in G6PD deficiency
    - Phototoxicity
  - Trimethoprim or pyrimethamine
    - Bone marrow suppression (leukopenia)

### Direct Inhibitors of Nucleic Acid Synthesis: Quinolones

- Drugs: ciprofloxacin, levofloxacin, and other “–floxacin”
- Mechanisms of action:
  - Quinolones are bactericidal and interfere with DNA synthesis
  - Inhibit topoisomerase II (DNA gyrase) and topoisomerase IV (responsible for separation of replicated DNA during cell division)
  - Resistance is increasing
- Activity and clinical uses:
  - Urinary tract infections (UTIs), particularly when resistant to cotrimoxazole
  - Sexually transmitted diseases (STDs)/pelvic inflammatory diseases (PIDs): chlamydia, gonorrhea
  - Skin, soft tissue, and bone infections by gram-negative organisms
  - Diarrhea to *Shigella*, *Salmonella*, *E. coli*, *Campylobacter*
  - Drug-resistant pneumococci (levofloxacin)

#### **Note**

The activity of quinolones includes *Bacillus anthracis*. Anthrax can also be treated with penicillins or tetracyclines.

- Pharmacokinetics:
  - Iron, calcium limit their absorption
  - Eliminated mainly by kidney by filtration and active secretion (inhibited by probenecid)
  - Reduce dose in renal dysfunction
- Side effects:
  - Tendonitis, tendon rupture
  - Phototoxicity, rashes
  - CNS effects (insomnia, dizziness, headache)
  - Contraindicated in pregnancy and in children (inhibition of chondrogenesis)

### Clinical Correlate

#### Antibiotics for *H. pylori* Gastrointestinal Ulcers

- "BMT" regimen: bismuth, metronidazole, and tetracycline
- Clarithromycin, amoxicillin, omeprazole

### Note

#### *Mycobacterium avium-intracellulare* (MAC)

- *Prophylaxis*: azithromycin (1 × week) or clarithromycin (daily)
- *Treatment*: clarithromycin + ethambutol ± rifabutin

## UNCLASSIFIED ANTIBIOTIC: METRONIDAZOLE

- Possibly producing free radicals, bactericidal
- Antiprotozoal: *Giardia*, *Trichomonas*, *Entamoeba*
- Antibacterial: strong activity against most anaerobic gram-negative *Bacteroides* species *Clostridium* species (DOC in pseudomembranous colitis), *Gardnerella*, and *H. pylori*
- Side effects:
  - Metallic taste
  - Disulfiram-like effect

## ANTITUBERCULAR DRUGS

- Combination drug therapy is the rule to delay or prevent the emergence of resistance and to provide additive (possibly synergistic) effects against *Mycobacterium tuberculosis*.
- The primary drugs in combination regimens are isoniazid (INH), rifampin, ethambutol, and pyrazinamide. Regimens may include two to four of these drugs, but in the case of highly resistant organisms, other agents may also be required. Backup drugs include aminoglycosides (streptomycin, amikacin, kanamycin), fluoroquinolones, capreomycin (marked hearing loss), and cycloserine (neurotoxic).
- *Prophylaxis*: usually INH, but rifampin if intolerant. In suspected multidrug resistance, both drugs may be used in combination.

- Mechanisms of action, resistance, and side effects:

**Table V-1-4. Summary of the Actions, Resistance, and Side Effects of the Antitubercular Drugs**

Drug	Mechanisms of Action and Resistance	Side Effects
Isoniazid (INH)	<ul style="list-style-type: none"> <li>• Inhibits mycolic acid synthesis</li> <li>• Prodrug requiring conversion by catalase</li> <li>• High level resistance—deletions in <i>katG</i> gene (encodes catalase needed for INH bioactivation)</li> </ul>	<ul style="list-style-type: none"> <li>• Hepatitis (age-dependent)</li> <li>• Peripheral neuritis (use vitamin B<sub>6</sub>)</li> <li>• Sideroblastic anemia (use vitamin B<sub>6</sub>)</li> <li>• SLE in slow acetylators (rare)</li> </ul>
Rifampin	<ul style="list-style-type: none"> <li>• Inhibits DNA-dependent RNA polymerase (nucleic acid synthesis inhibitor)</li> </ul>	<ul style="list-style-type: none"> <li>• Hepatitis</li> <li>• Induction of P450</li> <li>• Red-orange metabolites</li> </ul>
Ethambutol	<ul style="list-style-type: none"> <li>• Inhibits synthesis of arabinogalactan (cell-wall component)</li> </ul>	<ul style="list-style-type: none"> <li>• Dose-dependent retrobulbar neuritis → ↓ visual acuity and red-green discrimination</li> </ul>
Pyrazinamide		<ul style="list-style-type: none"> <li>• Hepatitis</li> <li>• Hyperuricemia</li> </ul>
Streptomycin	<ul style="list-style-type: none"> <li>• Protein synthesis inhibition (<i>see</i> Aminoglycosides)</li> </ul>	<ul style="list-style-type: none"> <li>• Deafness</li> <li>• Vestibular dysfunction</li> <li>• Nephrotoxicity</li> </ul>

### Clinical Correlate

#### INH Prophylaxis

- Exposure, TST-negative, young children
- TST conversion in past 2 years
- Tuberculin reactors with high risk: e.g., diabetes, immunosuppressive Rx, prolonged glucocorticoid Rx, HIV-positive, leukemia

## Chapter Summary

### Basic Principles

- \* Antimicrobial drugs can be either bactericidal or bacteriostatic. The effectiveness of bacteriostatic drugs depends on an intact host immune system. Antimicrobial agents may be administered singly or in combination. Some combinations induce synergy and/or delay emergence of resistance.
- \* An antimicrobial agent should have maximal toxicity toward the infecting agent and minimal toxicity for the host. Table V-1-1 summarizes the four basic antibacterial actions demonstrated by antibiotics and the agents working by each of these mechanisms.
- \* Microbial resistance can occur by the gradual selection of resistant mutants or more usually by R-factor transmission between bacteria. Table V-1-2 summarizes the common modes of resistance exhibited by microorganisms against the various classes of antimicrobial agents.

### Inhibitors of Bacterial Cell-Wall Synthesis

- \* The inhibitors of bacterial cell-wall synthesis are the beta-lactam antibiotics (the penicillins and cephalosporins; Figure V-1-1), the carbapenems, vancomycin, and aztreonam.
- \* The mechanisms of action of penicillins, the bacterial modes of resistance to penicillins, the penicillin subgroups, their biodisposition, and side effects are provided. The subgroups discussed are the penicillins that are  $\beta$ -lactamase susceptible with a narrow spectrum of activity;  $\beta$ -lactamase-resistant penicillins that have a very narrow spectrum of activity; and  $\beta$ -lactamase-susceptible penicillins that have a wider spectrum of activity. The common penicillins and their susceptible organisms are listed for each subgroup.
- \* The same parameters are considered for the cephalosporins. These have the same mode of action as the penicillins and also require an intact  $\beta$ -lactam ring structure for activity. There are four generations of cephalosporins. Each is considered in terms of range of activity, susceptibility to resistance, clinical usage, and specific antibiotics in that class.
- \* Imipenem and meropenem have the same mode of antibacterial action as the penicillins and cephalosporins but structurally are carbapenems that have the  $\beta$ -lactam ring. Their clinical uses, routes of elimination, and side effects are considered.
- \* Aztreonam is a monobactam inhibitor of early cell-wall synthesis. It is used primarily as an intravenous drug against gram-negative rods.
- \* Vancomycin inhibits an early stage of cell-wall synthesis. It has a relatively narrow range of activity, but as yet, resistance is uncommon. Its use, excretion, and side effects are considered.

### Inhibitors of Bacterial Protein Synthesis

- \* Figure V-1-2 illustrates the mechanisms of bacterial protein synthesis, and Table V-1-3 summarizes the places in the translatory sequence, as well as the mechanisms by which antibiotics operate to disrupt protein synthesis.
- \* The aminoglycosides (e.g., gentamicin and tobramycin) inhibit initiation complex formation. Their uses and properties are discussed. Streptomycin is particularly useful in the treatment of tuberculosis and is the drug of choice for treating bubonic plague and tularemia. Neomycin is toxic and can only be used topically.

(Continued)

### Chapter Summary (cont'd)

- \* The tetracyclines block the attachment of aminoacyl tRNA to the acceptor site on the bacterial ribosome. They are broad-spectrum drugs with good activity against chlamydial and mycoplasmal species, as well as against other indicated bacteria. Doxycycline is of particular use in the treatment of prostatitis, minocycline is useful for treating meningococcal carrier states, and demeclocycline is useful for treating the syndrome of inappropriate secretion of ADH (SIADH). Their biodisposition and side effects are discussed.
- \* Chloramphenicol inhibits the activity of peptidyltransferase and is currently used primarily as a backup drug. Its activity, clinical use, and side effects are considered.
- \* The macrolides (e.g., erythromycin, clarithromycin, and azithromycin) are translocation inhibitors. Their spectrums of activity, clinical uses, biodisposition, and side effects are considered. Clindamycin is not a macrolide but shares the same mechanism of action.
- \* Linezolid inhibits initiation by blocking formation of the N-formyl-methionyl-tRNA-ribosome-mRNA ternary complex. The clinical uses and side effects of this new drug are mentioned.
- \* Quinupristin and dalopristin bind to the 50S ribosomal subunit, where they interfere with the interaction of aminoacyl-tRNA and the acceptor site and also stimulate its dissociation from the ternary complex. Their clinical use and side effects are discussed.

### Antibiotics That Inhibit Folic Acid Synthesis and Nucleic Acid Metabolism

- \* The sulfonamides compete with para-aminobenzoic acid (PABA) as shown in Figure V-1-3. The methods bacteria use to develop resistance to the sulfonamides, their activity and clinical uses, biodisposition, and side effects are considered.
- \* Trimethoprim (TMP), a folate analog and inhibitor of dihydrofolate reductase (Figure V-1-3), is usually used together with sulfamethoxazole (SMX). The simultaneous inhibition of the tetrahydrofolate synthesis pathway at two steps has a synergistic effect and prevents the rapid generation of resistance. The clinical uses and side effects of TMP-SMX are discussed.
- \* The fluoroquinolones (e.g., ciprofloxacin) are nalidixic acid analogs that inhibit topoisomerase II (DNA gyrase) and topoisomerase IV. Their clinical use, the relevant drugs in this class, their biodisposition, and side effects are reported.
- \* The exact mode of metronidazole action is unknown. Its use as an antiprotozoal and antibacterial drug is discussed, as are its side effects.

### Antitubercular Drugs

- \* Infections caused by *Mycobacterium tuberculosis* are treated with combination therapy. The primary drugs used are isoniazid, rifampin, ethambutol, and pyrazinamide. Highly resistant organisms may require the use of additional agents. Backup drugs include streptomycin, fluoroquinolones, capreomycin, and cycloserine.
- \* Table V-1-4 summarizes the actions, resistance, and side effects of the antitubercular drugs.





# Antifungal Agents




Figure V-2-1. Mechanism of Action of Antifungal Drugs

## POLYENES (AMPHOTERICIN B [AMP B], NYSTATIN)

- Mechanisms:
  - Amphoteric compounds with both polar and nonpolar structural components—interact with **ergosterol** in fungal membranes to form artificial “pores,” which disrupt membrane permeability
  - Resistant fungal strains appear to have low ergosterol content in their cell membranes
- Activity and clinical uses:
  - Amphotericin B has wide fungicidal spectrum; remains the DOC (or co-DOC) for severe infections caused by *Aspergillus*, *Candida*, *Cryptococcus*, *Histoplasma*, *Mucor*, and *Sporothrix*
  - Amphotericin B—synergistic with flucytosine in candidiasis and cryptococcoses
  - Nystatin (too toxic for systemic use)—used topically for localized infections (e.g., candidiasis)
- Pharmacokinetics:
  - Amphotericin B given by slow IV infusion—poor penetration into the CNS (intrathecal possible)
  - Slow clearance (half-life >2 weeks) via both metabolism and renal elimination

- Side effects:
  - Infusion-related
    - Fever, chills, muscle rigor, hypotension (histamine release) occur during IV infusion (a test dose is advisable)
    - Can be alleviated partly by pretreatment with NSAIDs, antihistamines, meperidine, and adrenal steroids
  - Dose-dependent
    - Nephrotoxicity includes ↓ GFR, tubular acidosis, ↓ K<sup>+</sup> and Mg<sup>2+</sup>, and anemia through ↓ erythropoietin
    - Protect by Na<sup>+</sup> loading, use of liposomal amphotericin B, or by drug combinations (e.g., + flucytosine), permitting ↓ in amphotericin B dose

## **AZOLES (KETOCONAZOLE, FLUCONAZOLE, ITRACONAZOLE, VORICONAZOLE)**

- Mechanism:
  - “Azoles” are fungicidal and interfere with the synthesis of ergosterol by inhibiting 14- $\alpha$ -demethylase, a fungal P450 enzyme, which converts lanosterol to ergosterol
  - Resistance occurs via decreased intracellular accumulation of azoles
- Activity and clinical uses:
  - Ketoconazole
    - Co-DOC for *Paracoccidioides* and backup for *Blastomyces* and *Histoplasma*
    - Oral use in mucocutaneous candidiasis or dermatophytoses
  - Fluconazole
    - DOC for esophageal and invasive candidiasis and coccidioidomycoses
    - Prophylaxis and suppression in cryptococcal meningitis
  - Itraconazole and Voriconazole
    - DOC in blastomycoses, sporotrichoses, aspergillosis
    - Backup for several other mycoses and candidiasis
  - Clotrimazole and miconazole
    - Used topically for candidal and dermatophytic infections
- Pharmacokinetics:
  - Effective orally
  - Absorption of ketoconazole ↓ by antacids
  - Absorption of itraconazole ↑ by food
  - Only fluconazole penetrates into the CSF and can be used in meningeal infection. Fluconazole is eliminated in the urine, largely in unchanged form.
  - Ketoconazole and itraconazole are metabolized by liver enzymes.
  - Inhibition of hepatic P450s
- Side effects:
  - ↓ synthesis of steroids, including cortisol and testosterone → ↓ libido, gynecomastia, menstrual irregularities
  - ↑ liver function tests and rare hepatotoxicity

## OTHER ANTIFUNGALS

- Flucytosine
  - Activated by fungal cytosine deaminase to 5-fluorouracil (5-FU), which after triphosphorylation is incorporated into fungal RNA
  - 5-FU also forms 5-fluorodeoxyuridine monophosphate (5-Fd-UMP), which inhibits thymidylate synthase → ↓ thymine.
  - Resistance emerges rapidly if flucytosine is used alone.
  - Use in combination with amphotericin B in severe candidal and cryptococcal infections—enters CSF
  - Toxic to bone marrow (*see* Anticancer Drugs, Section IX).
- Griseofulvin
  - Active only against dermatophytes (orally, not topically) by depositing in newly formed keratin and disrupting microtubule structure
  - Side effects:
    - Disulfiram-like reaction
- Terbinafine
  - Active only against dermatophytes by inhibiting squalene epoxidase → ↓ ergosterol
  - Possibly superior to griseofulvin in onychomycoses
  - Side effects: GI distress, rash, headache, ↑ liver function tests → possible hepatotoxicity

### Chapter Summary

- \* In eukaryotes, fungal metabolism is somewhat similar to that in humans. Thus, most bacterial antibiotics are ineffective, and many otherwise potentially effective drugs are also toxic to their human hosts. A difference between fungi and humans susceptible to exploitation by antibiotics is the high concentration of ergosterol in their membranes.
- \* The polyenes amphotericin (amp B) are amphoteric compounds that bind to ergosterol, forming pores, which results in the leakage of intracellular contents. The activity, clinical uses, biodisposition, and side effects of these polyenes are discussed.
- \* The azoles (ketoconazole, fluconazole, clotrimazole, miconazole, and itraconazole) kill fungi by interfering with ergosterol synthesis. The mechanisms of action, clinical uses, biodisposition, and side effects are considered.
- \* Flucytosine is activated by fungal cytosine deaminase to form 5-fluorouracil (5-FU). It is sometimes used in combination with amp-B. Inasmuch as 5-FU is a classic anticancer agent, it is not surprising that flucytosine is also toxic to bone marrow.
- \* Griseofulvin and terbinafine are active against dermatophytes. Griseofulvin interferes with microtubule function; terbinafine blocks ergosterol synthesis.



# Antiviral Agents

Many antiviral drugs are antimetabolites that resemble the structure of naturally occurring purine and pyrimidine bases or their nucleoside forms. Antimetabolites are usually prodrugs requiring metabolic activation by host-cell or viral enzymes—commonly, such bioactivation involves phosphorylation reactions catalyzed by kinases.

- Site of action:




Figure V-3-1. Sites of Antiviral Drug Actions

Table V-3-1. Mechanism of Action of Antiviral Drugs

Mechanism of Action	Major Drugs
Block viral penetration/uncoating	Amantadine, enfuvirtide, maraviroc
Inhibit viral DNA polymerases	Acyclovir, foscarnet, ganciclovir
Inhibit viral RNA polymerases	Foscarnet, ribavirin
Inhibit viral reverse transcriptase	Zidovudine, didanosine, zalcitabine, lamivudine, stavudine, nevirapine, efavirenz
Inhibit viral aspartate protease	Indinavir, ritonavir, saquinavir, nelfinavir
Inhibit viral neuraminidase	Zanamivir, oseltamivir

## ANTIHERPETICS

### Acyclovir

- Mechanisms of action:
  - Monophosphorylated by viral thymidine kinase (TK), then further bioactivated by host-cell kinases to the triphosphate
  - Acyclovir-triphosphate is both a substrate for and inhibitor of viral DNA polymerase
  - When incorporated into the DNA molecule, acts as a chain terminator because it lacks the equivalent of a ribosyl 3' hydroxyl group
  - Resistance possibly due to changes in DNA polymerase or to decreased activity of TK
  - >50% of HSV strains resistant to acyclovir completely lack thymidine kinase (TK<sup>-</sup> strains)




Figure V-3-2. Common Mechanism for "ovirs" and NRTIs

- Activity and clinical uses:
  - Activity includes herpes simplex virus (HSV) and varicella-zoster virus (VZV)
  - There are topical, oral, and IV forms; has a short half-life
  - Reduces viral shedding in genital herpes; ↓ acute neuritis in shingles but has no effect on postherpetic neuralgia
  - Reduces symptoms if used early in chickenpox; prophylactic in immunocompromised patients
- Side effects:
  - Minor with oral use, more obvious with IV
  - Crystalluria (maintain full hydration) and neurotoxicity (agitation, headache, confusion—seizures in OD)
  - Is *not* hematotoxic
- Newer drugs—famciclovir and valacyclovir are approved for HSV infection and are similar to acyclovir in mechanism. They may have activity against strains resistant to acyclovir, but not TK<sup>-</sup> strains. They have a longer  $t_{1/2}$  than acyclovir.

## Ganciclovir

- Mechanisms of action:
  - Similar to that of acyclovir
  - First phosphorylation step is viral-specific; involves thymidine kinase in HSV and a phosphotransferase (UL97) in cytomegalovirus (CMV)
  - Triphosphate form inhibits viral DNA polymerase and causes chain termination
  - Resistance mechanisms similar to acyclovir
- Activity and clinical uses:
  - HSV, VZV, and CMV
  - Mostly used in prophylaxis and treatment of CMV infections, including retinitis, in AIDS and transplant patients—relapses and retinal detachment occur
- Side effects:
  - Dose-limiting hematotoxicity (leukopenia, thrombocytopenia), mucositis, fever, rash, and crystalluria (maintain hydration)
  - Seizures in overdose

## Foscarnet

- Mechanisms and clinical uses:
  - Not an antimetabolite, but still inhibits viral DNA and RNA polymerases
  - Uses identical to ganciclovir, plus > activity versus acyclovir-resistant strains of HSV
- Side effects:
  - Dose-limiting nephrotoxicity with acute tubular necrosis, electrolyte imbalance with hypocalcemia (tremors and seizures)
  - Avoid pentamidine IV (→↑ nephrotoxicity and hypocalcemia)

## TREATMENT OF HIV

### Reverse Transcriptase Inhibitors (RTIs)

- The original inhibitors of reverse transcriptases of HIV are nucleoside antimetabolites (e.g., zidovudine, the prototype) that are converted to active forms via phosphorylation reactions.
- **Nucleoside reverse transcriptase inhibitors (NRTIs):**
  - Are components of most combination drug regimens used in HIV infection
  - Are used together with a protease inhibitor (PI)
  - Highly active antiretroviral therapy (HAART) has often resulted in ↓ viral RNA, reversal of the decline in CD4 cells, and ↓ opportunistic infections
- **Nonnucleoside reverse transcriptase inhibitors (NNRTIs):**
  - RTIs that do not require metabolic activation: nevirapine, efavirenz
  - Are not myelosuppressant
  - Inhibit reverse transcriptase at a site different from the one NRTIs bind to
  - Additive or synergistic if used in combination with NRTIs and/or PIs



### Zidovudine (Azidothymidine, ZDV, AZT)

- Mechanisms of action:
  - Phosphorylated nonspecifically to a triphosphate that can inhibit reverse transcriptase (RT) by competing with natural nucleotides and can also be incorporated into viral DNA to cause chain termination.
  - Resistance occurs by mutations (multiple) in the gene that codes for RT.

### Other NRTIs

- Mechanism of action identical to that of zidovudine
- Each requires metabolic activation to nucleotide forms that inhibit reverse transcriptase
- Resistance mechanisms are similar
- Not complete cross-resistance between NRTIs
- Drugs differ in their toxicity profiles and are less bone-marrow suppressing than AZT
- Side effects:

#### **Clinical Correlate**

##### **Combinations to Avoid**

- DDI + DDC because of additive toxicity
- D4T + AZT because of competition for activation

**Table V-3-2. Side Effects of NRTIs**

Drug	Side Effects
Zidovudine, AZT	<ul style="list-style-type: none"> <li>• Hematotoxicity (major and dose-limiting)</li> <li>• Headache, asthenia, myalgia, myopathy, and peripheral neuropathy</li> </ul>
Didanosine, DDI	<ul style="list-style-type: none"> <li>• Pancreatitis (major and dose-limiting)</li> <li>• Peripheral neuropathy, hyperuricemia, liver dysfunction</li> </ul>
Zalcitabine, DDC	<ul style="list-style-type: none"> <li>• Peripheral neuropathy (major and dose-limiting)</li> <li>• GI distress, pancreatitis, neutropenia, rash</li> </ul>
Stavudine, D4T	<ul style="list-style-type: none"> <li>• Peripheral neuropathy (major and dose-limiting)</li> <li>• Myelosuppression &lt; ZDV</li> </ul>
Lamivudine, 3TC	<ul style="list-style-type: none"> <li>• Least toxic of the NRTIs, but some GI effects and neutropenia</li> <li>• Active in hepatitis B</li> </ul>

### Protease Inhibitors (PIs)

- Mechanisms of action:
  - Aspartate protease (*pol* gene encoded) is a viral enzyme that cleaves precursor polypeptides in HIV buds to form the proteins of the mature virus core.
  - The enzyme contains a dipeptide structure not seen in mammalian proteins. PIs bind to this dipeptide, inhibiting the enzyme.
  - Resistance occurs via specific point mutations in the *pol* gene, such that there is not complete cross-resistance between different PIs.
- Clinical uses:
  - The two PIs used most in the last 5 to 6 years, nearly always in combination regimens with two NRTIs, are indinavir and ritonavir.
  - Saquinavir, one of the least toxic, has very low (and variable) oral bioavailability that predisposes to resistance development.

- Side effects:
  - Indinavir
    - Crystalluria (maintain hydration)
  - Ritonavir
    - Major drug interactions: induces CYP 1A2 and inhibits the major P450 isoforms (3A4 and 2D6)
  - General: syndrome of disordered lipid and CHO metabolism with central adiposity and insulin resistance

## FUSION INHIBITORS:

### Enfuvirtide

- Mechanism of action: binds gp41 and inhibits fusion of HIV-1 to CD4+ cells

### Maraviroc

- Mechanism of action: blocks CCR5 protein on T-cell surface to prevent viral entry

## OTHER ANTIVIRALS

### Amantadine

- Mechanisms of action: blocks attachment, penetration, and uncoating of influenza A virus
- Clinical uses: prophylaxis mainly, but may ↓ duration of flu symptoms by 1–2 days
- Side effects:
  - CNS effects: nervousness, insomnia, and seizures in OD
  - Causes atropine-like peripheral effects and livedo reticularis

### Zanamivir and Oseltamivir

- Mechanisms of action:
  - Inhibit neuraminidases of influenza A and B (enzymes that prevent clumping of virions, so that more particles are available for infecting host cells)
  - Decreases the likelihood that the virus will penetrate uninfected cells
- Clinical uses: prophylaxis mainly, but may ↓ duration of flu symptoms by 2–3 days

### Ribavirin

- Mechanisms:
  - Monophosphorylated form inhibits IMP dehydrogenase
  - Triphosphate inhibits viral RNA polymerase and end-capping of viral RNA
- Clinical uses:
  - Adjunct to alpha-interferons in hepatitis C

## Clinical Correlate

### HIV Prophylaxis

*Needle stick*—ZDV+ 3TC, 1 month, but in high risk (e.g., high viral RNA copies), a combination of ZDV + 3TC + indinavir is recommended.

*Pregnancy*—ZDV full dose, trimester 2 and 3, plus 6 weeks to neonate, reduces vertical transmission by 80%—possible combination if high maternal viral RNA.

ZDV restricted to intrapartum period, or nevirapine (NNRTI) one dose at onset of delivery + one dose to neonate → ↓ transmission by 50 to 60%.

- Management of respiratory syncytial virus
- Lassa fever
- Hantavirus
- Side effects:
  - Hematotoxic
  - Upper airway irritation
  - Teratogenic

### **Human Interferons**

- Antiviral: hepatitis B, C, and D
- Antitumor: Kaposi sarcoma, CML, multiple myeloma, renal carcinoma
- Immunoregulatory: multiple sclerosis
- See Section X, Chapter 1 (Immunopharmacology)

## Chapter Summary

### General Principles

- \* Antiviral drugs are often antimetabolites that are structural analogs of purine or pyrimidine bases or their nucleoside forms. Many are prodrugs to be activated by host or viral enzymes. The steps in viral replication and the main sites of action of such antiviral drugs are illustrated in Figure V-3-1.
- \* Table V-3-1 summarizes the mechanisms of action of the major antiviral drugs.

### Antiherpetics

- \* The antiherpes drugs include acyclovir, ganciclovir, and foscarnet. Famciclovir and valacyclovir are newer drugs very similar to acyclovir. All inhibit viral DNA polymerase. Acyclovir and ganciclovir do so by first being phosphorylated by viral enzymes. As well as acting as a polymerase inhibitor, acyclovir triphosphate is incorporated into the viral DNA, where it acts as a chain terminator. The mechanisms of action, activities, clinical uses, and adverse effects are discussed.

### Reverse Transcriptase Inhibitors

- \* Nucleoside reverse transcriptase inhibitors (NRTIs) are used in most drug regimes to treat HIV infections. Commonly two NRTIs are used together with a protease inhibitor.
- \* The mechanisms, biodisposition, and adverse effects associated with zidovudine (AZT) use are described. The other nucleotide RTIs act almost identically. The NRTIs and their adverse effects are summarized in Table V-3-2.
- \* Nonnucleoside inhibitors of reverse transcriptase (NNRTIs) and a nucleotide RTI are also used in combinations for treatment in an HIV-positive patient.

### Protease Inhibitors (PIs)

- \* HIV aspartate protease has a unique dipeptide structure that has been used as a target for protease inhibitory drugs.
- \* The two protease inhibitors most used are indinavir and ritonavir. Their adverse effects are discussed.

### Fusion Inhibitors

- \* Two new drugs, enfuvirtide and maraviroc, block the entry of HIV into cells.

### Other Antivirals

- \* Amantadine blocks the attachment, penetration, and uncoating of influenza virus A; zanamivir and oseltamivir inhibit influenza viruses A and B neuraminidase, promoting viral clumping and decreasing the chance of penetration. Ribavirin becomes phosphorylated and inhibits IMP dehydrogenase and RNA polymerase. It is used to treat respiratory syncytial virus, influenza A and B, Lassa fever, Hantavirus, and as an adjunct to alpha-interferons in hepatitis C. The mechanisms, clinical uses, and side effects of these drugs are considered.



# Antiprotozoal Agents

## OVERVIEW

Table V-4-1. Major Protozoal Infections and the Drugs of Choice

Infection	Drug of Choice	Comments
Amebiasis	Metronidazole	Diloxanide for noninvasive intestinal amebiasis
Giardiasis	Metronidazole	“Backpacker’s diarrhea” from contaminated water or food
Trichomoniasis	Metronidazole	Treat both partners
Toxoplasmosis	Pyrimethamine + sulfadiazine	—
Leishmaniasis	Stibogluconate	—
Trypanosomiasis	Nifurtimox (Chagas disease) Arsenicals (African)	—

## ANTIMALARIAL DRUGS

- Clinical uses:
  - Chloroquine-sensitive regions
    - Prophylaxis: chloroquine +/- primaquine
    - Backup drugs: hydroxychloroquine, primaquine, pyrimethamine-sulfadoxine

- Specific treatment:

**Table V-4-2. Treatment of Chloroquine-Sensitive Malaria**

<i>P. falciparum</i>	Chloroquine
<i>P. malariae</i>	Chloroquine
<i>P. vivax</i>	Chloroquine + primaquine
<i>P. ovale</i>	Chloroquine + primaquine

- Chloroquine-resistant regions
  - Prophylaxis: mefloquine; backup drugs: doxycycline, atovaquone-proguanil
  - Treatment: quinine +/- either doxycycline or clindamycin or pyrimethamine
- Side effects:
  - Hemolytic anemia in G6PD deficiency (primaquine, quinine)
  - Cinchonism (quinine)

## DRUGS FOR HELMINTHIC INFECTIONS

- Most intestinal nematodes (worms)
  - Mebendazole ( $\downarrow$  glucose uptake and  $\downarrow$  microtubular structure)
  - Pyrantel pamoate (NM agonist  $\rightarrow$  spastic paralysis)
- Most cestodes (tapeworms) and trematodes (flukes)
  - Praziquantel ( $\uparrow$   $\text{Ca}^{2+}$  influx,  $\uparrow$  vacuolization)

### Chapter Summary

- \* Table V-4-1 lists the major types of protozoal infections and the drugs of choice for their treatment, with various relevant comments.
- \* Table V-4-2 lists the drugs of choice used against the various forms of malaria, and information is given about treatment and prophylaxis of malaria. Chloroquine-sensitive or -resistant areas are listed separately.
- \* The drugs used to treat helminthic infections are listed, and their mechanisms of action are noted.

# Antimicrobial Drug List and Practice Questions

Table V-5-1. Antimicrobial Drug List

<b>Penicillins</b>	<b>Cephalosporins</b>	<b>Other Cell Wall Inhibitors</b>	
Penicillin G	Cefazolin (1st)	Imipenem, meropenem	
Nafcillin, oxacillin	Cefaclor (2nd)	Vancomycin	
Amoxicillin, ampicillin	Ceftriaxone (3rd)		
Ticarcillin, piperacillin, azlocillin			
<b>Macrolides</b>	<b>Aminoglycosides</b>	<b>Tetracyclines</b>	<b>Others</b>
Erythromycin	Gentamicin	Tetracycline HCl	Metronidazole
Azithromycin	Tobramycin	Doxycycline	
Clarithromycin	Streptomycin		
<b>Fluoroquinolones</b>	<b>Antifolates</b>	<b>Antimycobacterials</b>	
Ciprofloxacin	Sulfamethoxazole	Isoniazid, rifampin	
Levofloxacin	Trimethoprim	Ethambutol, pyrazinamide	
<b>Antifungals</b>	<b>Anti-Herpes</b>	<b>Anti-HIV</b>	
Amphotericin B	Acyclovir	Zidovudine (NRTI), didanosine (NRTI)	
Ketoconazole	Ganciclovir	Lamivudine (NRTI)	
Fluconazole	Foscarnet	Indinavir (PI), ritonavir (PI) Enfuvirtide, maraviroc	



### Review Questions

1. A patient suffering from invasive aspergillosis is first administered NSAIDs, antihistamines, and adrenal glucocorticoids prior to administration of an antifungal drug. The antifungal drug works by
  - A. binding to tubulin
  - B. inhibiting squalene epoxidase
  - C. inhibiting thymine synthesis
  - D. binding to ergosterol
  - E. inhibiting 14 $\alpha$ -demethylase
2. A patient is prescribed isoniazid prophylactically since another family member currently has tuberculosis. When the patient ends up getting tuberculosis despite prophylaxis, resistance to isoniazid is suspected. In what way did this resistance likely develop?
  - A. Decreased intracellular accumulation of the drug
  - B. Inactivation of the drug via *N*-acetyltransferases
  - C. Increased synthesis of mycolic acids
  - D. Mutations in the gene coding for DNA-dependent RNA polymerase
  - E. Reduced expression of the gene that encodes a catalase
3. A 7-year-old child presents with pharyngitis and fever of 2 days' duration, and microbiology reveals small, translucent, beta-hemolytic colonies sensitive in vitro to bacitracin. Past history includes a severe allergic reaction to amoxicillin when used for an ear infection. The physician needs to treat this infection, but prefers not to use a drug that needs parenteral administration. Which one of the following agents is most likely to be appropriate in terms of both effectiveness and safety?
  - A. Azithromycin
  - B. Cefaclor
  - C. Doxycycline
  - D. Penicillin G
  - E. Vancomycin

4. A woman has a sexually transmitted disease, and the decision is made to treat her with antibiotics as an outpatient. She is warned that unpleasant reactions may occur if she consumes alcoholic beverages while taking this drug. The antibiotic can be identified as which of the following?
  - A. Ceftriaxone
  - B. Doxycycline
  - C. Metronidazole
  - D. Ofloxacin
  - E. Pen G
  
5. An 82-year-old hospitalized patient with creatinine clearance of 25 mL/min has a microbial infection requiring treatment with antibiotics. Which of the following drugs is least likely to require a dosage adjustment, either a smaller dose than usual or an increased interval between doses?
  - A. Amphotericin B
  - B. Ceftriaxone
  - C. Gentamicin
  - D. Imipenem-cilastatin
  - E. Vancomycin
  
6. What drug is most likely to be effective in most diseases caused by nematodes?
  - A. Chloroquine
  - B. Mebendazole
  - C. Metronidazole
  - D. Praziquantel
  - E. Pyrimethamine
  
7. What antibiotic effectively treats a variety of causative organisms for bacterial pneumonia, and also works at the 50S ribosomal subunit?
  - A. Azithromycin
  - B. Ceftriaxone
  - C. Doxycycline
  - D. Ofloxacin
  - E. Clindamycin

8. In bacterial meningitis, third-generation cephalosporins are commonly drugs of choice. However, in neonatal meningitis they would not provide coverage if the infection was due to which of the following organisms?
- A. Meningococci
  - B. *L. monocytogenes*
  - C. Pneumococci
  - D. *E. coli*
  - E. Group B streptococci
9. Which one of the following drugs inhibits bacterial protein synthesis, preventing the translocation step via its interaction with the 50S ribosomal subunit?
- A. Clindamycin
  - B. Gentamicin
  - C. Chloramphenicol
  - D. Imipenem
  - E. Tetracycline
10. Which of the following is a mechanism underlying the resistance of strains of *S. pneumoniae* to the widely used antibiotic ciprofloxacin?
- A. Reduced topoisomerase sensitivity to inhibitors
  - B. Increased synthesis of PABA
  - C. Formation of methyltransferases that change receptor structure
  - D. Structural changes in porins
  - E. Formation of drug-inactivating hydrolases
11. Gentamicin would be an ineffective drug for which of the following organisms?
- A. *E. coli*
  - B. *B. fragilis*
  - C. Pseudomonas
  - D. Listeria if combined with ampicillin
  - E. Proteus

12. In the treatment of a urinary tract infection in a patient known to have a deficiency of glucose-6-phosphate dehydrogenase, it would not be advisable to prescribe which of the following?
- A. Ciprofloxacin
  - B. Amoxicillin
  - C. Cephalexin
  - D. Doxycycline
  - E. Sulfamethoxazole
13. What is the most likely mechanism of resistance for methicillin-resistant staphylococcus aureus to antistaph penicillins?
- A. methylation of the binding site
  - B. active efflux of the drug from the bacteria
  - C.  $\beta$ -lactamase production
  - D. phosphorylation of the drug by bacterial enzymes
  - E. structural modifications of PBPs
14. Highly active antiretroviral therapy (HAART) in HIV infection is associated with which of the following?
- A. A decrease in viral mRNA copies/mL of blood
  - B. A decrease in the rate of emergence of drug resistance
  - C. A possible increase in CD4 cell count
  - D. A reduced incidence of opportunistic infections
  - E. All of the above
15. Oseltamivir and zanamivir are available for treatment of infections due to influenza A and B. The mechanism of their antiviral action is inhibition of which of the following?
- A. RNA polymerase
  - B. Reverse transcriptase
  - C. Thymidine kinase
  - D. Neuraminidase
  - E. Aspartate protease

16. In a patient who has an established hypersensitivity to metronidazole, what is the most appropriate drug to use for the management of pseudomembranous colitis?
- A. Ampicillin
  - B. Clindamycin
  - C. Doxycycline
  - D. Ofloxacin
  - E. Vancomycin
17. An AIDS patient who is being treated with multiple drugs, including AZT, lamivudine, indinavir, ketoconazole, and cotrimoxazole, develops breast hypertrophy, central adiposity, hyperlipidemia, insulin resistance, and nephrolithiasis. If these changes are related to his drug treatment, which of the following is the most likely cause?
- A. Azidothymidine
  - B. Indinavir
  - C. Ketoconazole
  - D. Sulfamethoxazole
  - E. Trimethoprim
18. Which one of the following drugs is most suitable in an immunocompromised patient for prophylaxis against infection due to *Cryptococcus neoformans*?
- A. Amphotericin B
  - B. Ampicillin
  - C. Fluconazole
  - D. Nystatin
  - E. Flucytosine
19. Which one of the following drugs is most likely to be associated with elevations of pancreatic enzymes, including amylase and lipase?
- A. Erythromycin
  - B. Didanosine
  - C. Isoniazid
  - D. Zidovudine
  - E. Pyrazinamide

20. The major mechanism of HSV resistance to acyclovir is
- A. a structural change in viral thymidine kinase
  - B. a mutation in the gene that encodes DNA polymerase
  - C. the loss of ability to produce viral thymidine kinase
  - D. changes in reverse transcriptase
  - E. mutations in the gene that codes for phosphotransferase
21. Despite its “age,” penicillin G remains the drug of choice in the treatment of infections caused by which of the following organisms?
- A. *B. fragilis*
  - B. *T. pallidum*
  - C. *H. influenzae*
  - D. *E. coli*
  - E. *S. aureus*
22. Which one of the following drugs is most likely to be equally effective in the treatment of amebic dysentery and “backpacker’s diarrhea”?
- A. Ciprofloxacin
  - B. Diloxanide
  - C. Metronidazole
  - D. Quinacrine
  - E. Trimethoprim-sulfamethoxazole

## Answers

1. **Answer: D.** Life-threatening invasive aspergillosis, with necrotizing pneumonia, most commonly occurs in severely immunocompromised patients. The mortality rate approaches 50%, but high intravenous doses of amphotericin B may be lifesaving. Intravenous amphotericin B causes infusion-related hypotension (via histamine release), fever, and chills, which may be attenuated by the prior administration of NSAIDs and antihistamines. Adrenal steroids may provide supplementary stress support. Amphotericin B binds to ergosterol in fungal membranes, opening pores and disrupting membrane permeability.
2. **Answer: E.** For antitubercular activity, isoniazid (INH) must first be metabolically activated via a catalase present in mycobacteria. A decrease in expression of the *cat G* gene that encodes this enzyme is the mechanism of high-level resistance to INH.
3. **Answer: A.** Azithromycin is highly effective as an oral agent in the management of pharyngitis caused by gram-positive cocci and may necessitate only a short course of therapy. In patients who have marked hypersensitivity to penicillins, it is inappropriate to use a cephalosporin, even though cefaclor is active against common oropharyngeal pathogens. Doxycycline should not be used in children. One must assume that complete cross-allergenicity exists between different members of the penicillin class of antibiotics, and, in any case, penicillin G is not usually given orally because of its lability in gastric acid. Vancomycin would need parenteral administration, and this antibiotic should be reserved for more serious bacterial infections.
4. **Answer: C.** Organisms associated with sexually transmitted diseases include chlamydia, neisseria gonorrhoea, treponema (syphilis), trichomonas, and gardnerella vaginalis. The latter two organisms are effectively treated with the drug metronidazole. Metronidazole has a chemical structure that results in a disulfiram-like effect on aldehyde dehydrogenase, causing reactions with ethanol. Patients should be cautioned not to consume alcoholic beverages while on this drug.
5. **Answer: B.** Ceftriaxone is eliminated largely via biliary excretion, and decreases in renal function do not usually require a dose reduction. All of the other antimicrobial drugs listed are eliminated by the kidney, at rates proportional to creatinine clearance, so major dose reductions would be needed in patients with renal dysfunction to avoid toxicity.
6. **Answer: B.** Mebendazole is the drug of choice for treatment of all nematode infections (hookworm, roundworm, pinworm, whipworm). Pyrantel is considered equally effective as mebendazole for nematodes. Praziquantel is used for tapeworms (cestodes) and flukes (trematodes).

7. **Answer: A.** Macrolides (azithromycin) are effective for common causes of pneumonias such as *Strep pneumoniae*, *Haemophilus influenzae*, *Mycoplasma*, *Legionella*, and *Chlamydia*. The drugs work at the 50S ribosomal subunit to inhibit translocation of the peptidyl tRNA from the acceptor to the donor site.
8. **Answer: B.** The most common pathogens implicated in bacterial meningitis in a neonate (age <1 month) are group B streptococci, followed by *E. coli*. Meningococci and pneumococci become prevalent after 1 month of age, and *H. influenzae* is becoming rarer since the availability of a vaccine. A third-generation cephalosporin (e.g., cefotaxime) would be administered because it provides coverage for most of the organisms mentioned. However, ampicillin is also needed to cover for *Listeria monocytogenes*, which occurs with an incidence of 7 to 8% in neonatal meningitis.
9. **Answer: A.** Clindamycin has a mechanism of action similar to, if not identical with, erythromycin and related macrolides. They bind to rRNA bases on the 50S subunit to prevent translocation of peptidyl-mRNA from the acceptor to the donor site. Chloramphenicol also binds to the 50S subunit but interferes with the activity of peptidyltransferase. Gentamicin and tetracyclines bind to the 30S ribosomal subunit. Imipenem is a cell-wall synthesis inhibitor, acting similarly to beta-lactams.
10. **Answer: A.** Microbial resistance to fluoroquinolones is increasing, and some strains of *Streptococcus pneumoniae* are now resistant to ciprofloxacin. The mechanism can involve changes in the structure of topoisomerase IV, one of the “targets” of fluoroquinolones, which inhibit nucleic acid synthesis. Pneumococcal resistance to penicillins is also increasing via changes in penicillin-binding proteins (PBPs). The other mechanisms listed underlie microbial resistance to other antibiotics as follows: sulfonamides (**choice B**), macrolides (**choice C**), extended-spectrum penicillins (**choice D**), and beta-lactams (**choice E**).
11. **Answer: B.** Aminoglycosides like gentamicin work on aerobic gram negative rods. They require oxygen to enter bacteria, and, as such, do not treat any anaerobes including *Bacteroides fragilis*. They can be used with penicillins such as ampicillin against *Listeria* for a synergistic effect.
12. **Answer: E.** Drugs that cause oxidative stress may precipitate acute hemolysis in patients who lack G6PD because they have a limited ability to generate NADPH, which restricts the formation of glutathione. Drugs in this category include primaquine, quinine, nitrofurantoin, sulfonamides, and TMP-SMX.
13. **Answer: E.** Antistaph penicillins are inherently resistant to cleavage by bacterial beta-lactamases. Instead, resistance develops when the target for these drug, PBPs, are altered such that the drug doesn't bind effectively.



14. **Answer: E.** HAART in the management of HIV infection is reported in many but not all patients to decrease viral load, increase CD4 cells, slow disease progression, and reduce opportunistic infections. However, in terms of the chemotherapy of AIDS, the word *cure* has little meaning. Discontinuance of HAART, after suppression of viral RNA copies below the sensitivity of the best current methods of analysis, is followed by the reemergence of detectable viral RNA in the blood within a few months.
15. **Answer: D.** Neuraminidase is an enzyme on the lipid envelope of influenza A and B virions that prevents their clumping together and also their binding to the surface of cells that have been already infected. Neuraminidase inhibitors interfere with this activity and reduce the availability of virions for entry into noninfected cells. Oseltamivir and zanamivir decrease the severity and duration of symptoms if given within a day or two of onset.
16. **Answer: E.** Vancomycin is usually considered to be a backup drug to metronidazole in colitis due to *Clostridium difficile* on the grounds that it is no more effective, is more costly, and should be reserved for treatment of resistant gram-positive coccal infections. None of the other drugs has activity in pseudomembranous colitis—indeed, they may cause it!
17. **Answer: B.** AIDS patients being treated with protease inhibitors (e.g., indinavir) have developed a syndrome involving derangement of lipid and CHO metabolism. Changes in lipid metabolism and distribution occur quite commonly, and type 2 diabetes has also been reported. Indinavir is also notable for its tendency to precipitate in the urinary tract, causing nephrolithiasis, unless the patient is maintained in a high state of hydration.
18. **Answer: C.** Fluconazole is distinctive in terms of its ability to penetrate into the cerebrospinal fluid, reaching levels similar to those in the blood. It is effective against *C. neoformans* and has become the most appropriate drug to use in both prophylaxis and suppression because of its oral efficacy and low toxicity compared with amphotericin B. Flucytosine is also active against *C. neoformans* but is not used alone because of rapid emergence of resistance. Nystatin is too toxic for systemic use.
19. **Answer: B.** Pancreatic dysfunction, heralded by large increases in serum amylase and lipase, is associated with the use of several reverse-transcriptase inhibitors (RTIs). Didanosine appears to be the worst offender, and pancreatitis is the most characteristic adverse effect of this particular NRTI. Conditions enhancing susceptibility to drug-induced pancreatic dysfunction include hypertriglyceridemia, hypercalcemia, and history of excessive ethanol use. Liver dysfunction including hepatitis may occur with the antitubercular drugs, isoniazid, and pyrazinamide. Cholestasis is associated with the estolate form of erythromycin.
20. **Answer: C.** To inhibit DNA polymerases in HSV, acyclovir must undergo initial monophosphorylation by a viral specific thymidine kinase (TK). Most HSV strains resistant to acyclovir lack this enzyme and are thus TK<sup>-</sup> strains. A few strains of HSV are resistant to acyclovir by structural changes in TK that lower substrate affinity or by mutations in the gene that encode viral DNA polymerases.

21. **Answer: B.** Indications for the use of penicillin G are currently limited for a number of reasons. The drug has a narrow spectrum, is susceptible to beta-lactamases, and may cause hypersensitivity. Also, alternative antibiotics are available. However, penicillin G remains the drug of choice in syphilis, usually given IM as benzathine penicillin G, but as the Na<sup>+</sup> or K<sup>+</sup> salt IV in neurosyphilis. What would you do for patients who are highly allergic to penicillins? (Consider tetracyclines, or possibly desensitization.)
22. **Answer: C.** In amebic dysentery caused by *Entamoeba histolytica* and gastrointestinal infections with diarrhea (“backpacker’s diarrhea”) due to *Giardia lamblia*, metronidazole is the drug of choice. Diloxanide is a backup drug for noninvasive intestinal amebiasis, but it has minimal activity in *Giardia* infections. Quinacrine has effectiveness in giardiasis but not amebiasis. TMP-SMX has antiprotozoal effectiveness in *Pneumocystis carinii* pneumonia. Ciprofloxacin is devoid of antiprotozoal activity.



SECTION VI

**Drugs for Inflammatory  
and Related Disorders**



# Histamine and Antihistamines

---

## HISTAMINE

- Histamine is an autacoid present at high levels in the lungs, skin, and the gastrointestinal tract and is released from mast cells and basophils by type I hypersensitivity reactions, drugs, venoms, and trauma.
- Histamine receptors are of the serpentine family, with seven transmembrane-spanning domains with G-protein-coupled second messenger effectors.
  - H<sub>1</sub> activation
    - ↑ capillary dilation (via NO) → ↓ BP
    - ↑ capillary permeability → ↑ edema
    - ↑ bronchiolar smooth muscle contraction (via IP<sub>3</sub> and DAG release)
    - ↑ activation of peripheral nociceptive receptors → ↑ pain and pruritus
    - ↓ AV nodal conduction
  - H<sub>2</sub> activation
    - ↑ gastric acid secretion → ↑ gastrointestinal ulcers
    - ↑ SA nodal rate, positive inotropism, and automaticity

## H<sub>1</sub> ANTAGONISTS

- Mechanism of action:
  - H<sub>1</sub> antagonists act as competitive antagonists of histamine and therefore may be ineffective at high levels of histamine.
  - Vary in terms of both pharmacologic and kinetic properties, but all require hepatic metabolism and most cross the placental barrier.

**Table VI-1-1. Properties of Major Antihistamines**

Drug	M Block	Sedation	Antimotion	Other Characteristics
Diphenhydramine	+++	+++	+++	Widely used OTC drug
Promethazine	+++	+++	++	Some $\alpha$ block and local anesthetic action
Chlorpheniramine	++	++	++	Possible CNS stimulation
Meclizine	++	++	++++	Highly effective in motion sickness
Cetirizine	+/-	+	0	
Loratadine	+/-	0	0	No CNS entry
Fexofenadine	+/-	0	0	No CNS entry

- Uses:
  - Allergic reactions: hay fever, rhinitis, urticaria
  - Motion sickness, vertigo
  - Nausea and vomiting with pregnancy
  - Preoperative sedation
  - OTC: sleep aids and cold medications
  - Parkinson disease
  - Acute EPSs
- Side effects:
  - Extensions of M block and sedation (additive with other CNS depressants)

**Chapter Summary**

- \* Histamine is an autacoid released from mast cells and basophils by type I hypersensitivity reactions or under the influence of drugs, venoms, or trauma. Histamine receptors are the G-protein-coupled, seven-transmembrane type. Three different receptors are recognized: the well-characterized H<sub>1</sub> and H<sub>2</sub> types and an H<sub>3</sub> variant.
- \* The sequence of reactions leading to H<sub>1</sub> and H<sub>2</sub> activation is presented.
- \* H<sub>1</sub> antagonists are competitive inhibitors with varying pharmacologic and kinetic properties. All require hepatic metabolism and cross the placental barrier.
- \* The H<sub>1</sub> antagonists are used to treat allergic reactions, motion sickness, vertigo, nausea and vomiting in pregnancy, and preoperative sedation, and are in over-the-counter sleeping pills.
- \* The adverse effects are excess M block and sedation. Table VI-1-1 summarizes the properties of some of the major type 1 antihistamines.

# Drugs Used in Gastrointestinal Dysfunction

## DRUGS USED IN PEPTIC ULCER DISEASE (PUD)




Figure VI-2-1. Drug Actions in PUD

### H<sub>2</sub> Antagonists (e.g., Cimetidine, Ranitidine)

- Mechanisms of action:
  - Suppress secretory responses to food stimulation and nocturnal secretion of gastric acid via their ability to decrease (indirectly) the activity of the proton pump.
  - Also partially antagonize HCl secretion caused by vagally or gastrin-induced release of histamine from ECL-like cells (GI mast cells)
  - No effects on gastric emptying time



- Uses:
  - PUD (overall less effective than proton pump inhibitors)
  - Gastroesophageal reflux disease (GERD)
  - Zollinger-Ellison syndrome
- Side effects:
  - Cimetidine is a major inhibitor of P450 isoforms → drug interaction via ↑ effects
  - Cimetidine → ↓ androgens → gynecomastia and ↓ libido

### Proton Pump Inhibitors

- Mechanism of action:
  - **Omeprazole** and related “-prazoles” are irreversible, direct inhibitors of the proton pump ( $K^+/H^+$  antiport) in the gastric parietal cell
- Uses:
  - More effective than  $H_2$  blockers in peptic ulcer disease (PUD)
  - Also effective in GERD and Zollinger-Ellison syndrome
  - Eradication regimen for *H. pylori*

### Misoprostol

- Mechanism of action:  $PGE_1$  analog, which is cytoprotective → ↑ mucus and bicarbonate secretion and ↓ HCl secretion
- Uses: Selective use in NSAID-induced gastrointestinal ulcers

### Sucralfate

- Mechanism of action: polymerizes on gastrointestinal luminal surface to form a protective gel-like coating of ulcer beds. Requires acid pH (antacids may interfere)
- Uses: ↑ healing and ↓ ulcer recurrence

### Bismuth Subsalicylate

- Mechanism of action: like sucralfate, binds selectively to ulcer, coating it, and protecting it from acid and pepsin
- Combined with metronidazole and tetracycline to eradicate *H. pylori* (BMT regimen)

**Antacids:  $Al(OH)_3$ ,  $Mg(OH)_2$ ,  $CaCO_3$**


- Mechanism of action: bases that neutralize protons in the gut lumen
- Side effects: Constipation ( $Al^{+++}$ ), diarrhea ( $Mg^{++}$ )

**Clinical Correlate**

**Antacids and Drug Absorption**

- ↑ Oral absorption of weak bases (e.g., quinidine)
- ↓ Oral absorption of weak acids (e.g., warfarin)
- ↓ Oral absorption of tetracyclines (via chelation)

**ANTIEMETICS**



**Figure VI-2-2. The Emetic Pathways and Drug Action**

Figure VI-2-2 shows the complexity of the emetic pathways with an impact on the vomiting center and reveals the multiplicity of receptor types involved, including those activated by ACh, DA, 5HT, histamine, and endogenous opiopeptides.

Drugs for nausea and vomiting include:

- $5HT_3$  (a serotonin receptor—see following chapter) antagonists: ondansetron (commonly used in cancer chemotherapy), granisetron
- DA antagonists: prochlorperazine, metoclopramide (also used in cancer chemotherapy; also prokinetic in GERD)
- $H_1$  antagonists: diphenhydramine, meclizine, promethazine
- Muscarinic antagonists: scopolamine
- Cannabinoids: dronabinol
- $NK_1$ -receptor antagonist: aprepitant ( $NK_1$  is a receptor to substance P)

**Clinical Correlate**

Opioid analgesics (e.g., morphine) have duality of action: ↓ emesis by activating receptors that decrease pain transmission and ↑ emesis by activating receptors in the CTZ.

**Chapter Summary**

- \* The H<sub>2</sub> histamine antagonists (e.g., cimetidine and ranitidine) are used to suppress the secretion of gastric acid. The mechanism of action is illustrated in Figure VI-2-1. The clinical uses and adverse effects are discussed.
- \* Omeprazole and the other “-prazole” proton-pump inhibitors are more powerful inhibitors of gastric secretion than are the antagonists. Their clinical uses and adverse reactions are considered.
- \* Misoprostol is a cytoprotective prostaglandin E<sub>1</sub> analog.
- \* Sucralfate forms a protective gel, covering gastrointestinal ulcers. Bismuth subsalicylate behaves similarly.
- \* Antacids neutralize preformed protons.
- \* Figure VI-2-2 illustrates the number of complex factors impinging upon the emetic (vomiting) center. The antiemetic drugs are listed.

# Drugs Acting on Serotonergic Systems

---

- Serotonin (5-hydroxytryptamine, 5HT) is an autacoid synthesized and stored in gastrointestinal cells, neurons, and platelets. Metabolized by MAO type A, its metabolite 5-hydroxyindolacetic acid (5HIAA) is a marker for carcinoid.
- Of the seven receptor subtype families, all are G-protein coupled, except 5HT<sub>3</sub>, which is coupled directly to an ion channel.

## DRUG ACTIONS ON 5HT RECEPTORS

### 5HT<sub>1(a-h)</sub>

- Found in the CNS (usually inhibitory) and smooth muscle (excitatory or inhibitory)
- Drug: **bupirone**
  - Partial agonist at 5HT<sub>1a</sub> receptors → anxiolytic (generalized anxiety disorder [GAD])
- Drug: **sumatriptan**
  - Agonist at 5HT<sub>1d</sub> receptors in cerebral vessels → ↓ migraine pain
  - Side effects of “-triptans”: possible asthenia, chest or throat pressure or pain

### 5HT<sub>2(a-c)</sub>

- Found in CNS (excitatory)
- In periphery, activation → vasodilation, contraction of gastrointestinal, bronchial, and uterine smooth muscle, and platelet aggregation
- Drugs:
  - **Olanzapine** and other atypical antipsychotics: antagonist at 5HT<sub>2a</sub> receptors in CNS → ↓ symptoms of psychosis
  - **Cyproheptadine**
    - 5HT<sub>2</sub> antagonist used in carcinoid, other gastrointestinal tumors, and postgastrorectomy; also used for anorexia nervosa
    - Has marked H<sub>1</sub>-blocking action: used in seasonal allergies

### 5HT<sub>3</sub>

- Found in area postrema, peripheral sensory and enteric nerves
- Mechanism of action: activation opens ion channels (no second messengers)
- Drugs: **ondansetron** and “-setrons”
  - Antagonists → ↓ emesis in chemotherapy and radiation and postoperatively

## DRUGS USED IN MIGRAINE HEADACHES

- Ergot alkaloids
  - **Ergotamine**
    - Mechanism of action:  
Ergotamine acts as partial agonists at both  $\alpha$  and 5HT<sub>2</sub> receptors in the vasculature and possibly in the CNS.  
Vasoconstrictive actions to decrease pulsation in cerebral vessels may be relevant to acute actions of ergotamine during migraine attack.
    - Uses: ergotamine used in acute attacks
    - Side effects: gastrointestinal distress, prolonged vasoconstriction → ischemia and gangrene, abortion near term
- In addition to the “-triptans” and ergots:
  - Analgesics: ASA (+/- caffeine, or butabarbital), other NSAIDs, acetaminophen (+/- caffeine), oral or injectable opioid-analgesics, and butorphanol (spray)
  - Prophylaxis: propranolol, verapamil, amitriptyline, valproic acid

### Note

#### Ergonovine

- Mechanism of action: uterine smooth muscle contraction
- Use: given intramuscularly after placental delivery

### Chapter Summary

- \* Serotonin (5HT) is an autacoid synthesized and stored in gastrointestinal cells, neurons, and platelets. Monoamine oxidase (MAO) type A degrades it, forming 5-hydroxyindoleacetic acid (5HIAA), a carcinoid marker.
- \* There are seven receptor subtypes, six of which are G-protein coupled. The seventh type, 5HT<sub>3</sub>, is directly coupled to an ion channel.
- \* The locations and normal functions of different types of 5HT receptors, as well as drugs acting on them, are described.
- \* There are approximately 20 natural ergot alkaloids. A few of these, plus some derivatives, are used pharmacologically. Several act via 5HT receptors, but  $\alpha$ - and D<sub>2</sub> receptors are also utilized. The clinical uses and properties of specific ergots are indicated.
- \* Drugs (in addition to the “-triptans” and ergots) used to treat migraines are mentioned, as are other drugs affecting serotonergic neurotransmission.

# Eicosanoid Pharmacology

## OVERVIEW

- Eicosanoids are cell-regulating polyunsaturated fatty acids primarily synthesized from arachidonic acid and released by the action of phospholipase A<sub>2</sub> from lipids in cell membranes.
- Eicosanoids are present in low concentrations in most cells but are synthesized and released “on demand” in response to stimuli, including IgE-mediated reactions, inflammatory mediators, trauma, heat, and toxins.
- Eicosanoids interact with specific receptors, which are G-proteins coupled to second messenger effector systems.




Figure VI-4-1. Drugs Acting on Eicosanoids

## Bridge to Physiology

Prostaglandins (PGs) are cytoprotective in the stomach, dilate renal vasculature, contract the uterus, and maintain the ductus arteriosus. Thromboxane (TxA<sub>2</sub>) causes platelet aggregation. GI PGs and platelets TxA<sub>2</sub>s are synthesized by COX 1 (constitutive). COX 2 (inducible) synthesizes PGs involved in inflammation, fever, and pain. Both enzymes synthesize renal PGs → ↑ RBF.

## LEUKOTRIENES (LTs)

- Formed (via hydroperoxides) from the action of lipoxygenases on arachidonic acid
  - $LTB_4$ :
    - Mechanism of action: inflammatory mediator → neutrophil chemoattractant; activates PMNs; ↑ free radical formation → cell damage
  - $LTA_4$ ,  $LTC_4$ , and  $LTD_4$ 
    - Cause anaphylaxis and bronchoconstriction (role in asthma)
- Leukotrienes are “targets” for the following:
  - Glucocorticoids: → ↓ phospholipase  $A_2$  activity → contributes to both antiinflammatory and immunosuppressive actions
  - Zileuton: inhibits lipoxygenase → ↓ LTs and is used in treatment of asthma
  - Zafirlukast and “-lukasts”: LT-receptor antagonists used in treatment of asthma

## PROSTAGLANDINS (PGs)

- PGs are formed (via endoperoxides) from the actions of cyclooxygenases (COXs).
- COX 1 is expressed in most tissues, including platelets and stomach, where it acts to synthesize thromboxane and cytoprotective prostaglandins, respectively.
- COX 2 is expressed in the brain and kidney and at sites of inflammation.

### Note

Indomethacin is used to close a patent ductus arteriosus.

### PGE<sub>1</sub>

- Drugs:
  - **Misoprostol** (analog) used in treatment of NSAID-induced ulcers (protective action on gastric mucosa)
  - **Alprostadil**
    - Maintains patency of ductus arteriosus
    - Vasodilation; used in male impotence
- Contraindicated in pregnancy, unless used as an abortifacient (misoprostol in combination with mifepristone)

### PGE<sub>2</sub>

- Mechanism of action: uterine smooth muscle contraction
- Uses: **dinoprostone** can be used for “cervical ripening” and as abortifacient

### PGF<sub>2α</sub>

- Mechanism of action: uterine and bronchiolar smooth muscle contraction
- Drugs:
  - **Carboprost** used as abortifacient
  - **Latanoprost** for treatment of glaucoma (↓ intraocular pressure)

**PGI<sub>2</sub> (Prostacyclin)**

- Platelet stabilizer and vasodilator
- Drug: **epoprostenol**
- Uses: pulmonary hypertension

**PGE<sub>2</sub> and PGE<sub>2α</sub>**

- Both ↑ in primary dysmenorrhea
- Therapeutic effects of NSAIDs may be due to inhibition of their synthesis

**THROMBOXANES (TXAs)****TXA<sub>2</sub>**

- Platelet aggregator (inhibition of synthesis underlies protective role of acetylsalicylic acid [ASA] post-MI)

**NONSTEROIDAL ANTIINFLAMMATORY DRUGS (NSAIDs)**

- Most NSAIDs are nonselective inhibitors of cyclooxygenases, acting on both COX 1 and COX 2 isoforms to decrease formation of PGs and thromboxanes.
- They are analgesic, antipyretic, and antiinflammatory and have antiplatelet effects.
- Acetylsalicylic acid (ASA) is the prototype of the group, which includes more than 20 individual drugs.

**Acetylsalicylic Acid (ASA; Aspirin)**

- Causes irreversible inhibition of COX
- Covalent bond via acetylation of a serine hydroxyl group near the active site
- Actions are dose-dependent:
  - *Antiplatelet aggregation.* Low dose, the basis for post-MI prophylaxis and to reduce the risk of recurrent TIAs
  - *Analgesia and antipyresis.* Moderate dose
  - *Antiinflammatory.* High doses
  - *Uric acid elimination*
    - Low to moderate doses: ↓ tubular secretion → hyperuricemia
    - High doses: ↓ tubular reabsorption → uricosuria
  - *Acid-base and electrolyte balance*
    - Dose-dependent actions
    - High therapeutic: mild uncoupling of oxidative phosphorylation → ↑ respiration → ↓ pCO<sub>2</sub> → *respiratory alkalosis* → renal compensation → ↑ HCO<sub>3</sub><sup>-</sup> elimination → *compensated respiratory alkalosis* (pH = normal, ↓ HCO<sub>3</sub><sup>-</sup>, ↓ pCO<sub>2</sub>)
    - In adults, this can be a stable condition; in children → ↑ toxicity.
    - Toxic doses: inhibits respiratory center → ↓ respiration → ↑ pCO<sub>2</sub> → *respiratory acidosis* (↓ pH, ↓ HCO<sub>3</sub><sup>-</sup>, normalization of pCO<sub>2</sub>) plus inhibition of Krebs cycle and severe uncoupling of oxidative phosphorylation (↓ ATP) → *metabolic acidosis*, hyperthermia, and hypokalemia (↓ K<sup>+</sup>).

**Bridge to Physiology and Biochemistry****Platelet Stability and Eicosanoids**

Activation of TxA<sub>2</sub> receptors → stimulation of phospholipase C → ↑ PIP<sub>2</sub> hydrolysis → ↑ IP<sub>3</sub> → mobilization of bound Ca<sup>2+</sup> → ↑ free Ca<sup>2+</sup> → platelet aggregation.

Activation of PGI<sub>2</sub> receptors → stimulation of adenylyl cyclase → ↑ cAMP → ↑ activity of internal Ca<sup>2+</sup> “pumps” → ↓ free Ca<sup>2+</sup> → platelet stabilization.



- Side effects:
  - Gastrointestinal irritation: gastritis, ulcers, bleeding
  - Salicylism: tinnitus, vertigo, ↓ hearing—often first signs of toxicity
  - Bronchoconstriction: exacerbation of asthma
  - Hypersensitivity, especially the “triad” of asthma, nasal polyps, rhinitis
  - Reye syndrome: encephalopathy
  - ↑ bleeding time (antiplatelet)
  - Chronic use: associated with renal dysfunction
  - Drug interactions: ethanol (↑ gastrointestinal bleeding), OSUs and warfarin (↑ effects), and uricosurics (↓ effects)
- Aspirin overdose and management:
  - Extensions of the toxic actions described above, plus at high doses vasomotor collapse occurs, with both respiratory and renal failure.
  - No specific antidote. Management includes gastric lavage (+/– activated charcoal) plus ventilatory support and symptomatic management of acid-base and electrolyte imbalance, and the hyperthermia and resulting dehydration. ↑ urine volume and its alkalization facilitate salicylate renal elimination. (Note: ASA follows zero-order elimination kinetics at toxic doses.)

### Other NSAIDs

#### Types

- Reversible inhibitors of COX 1 and COX 2, with analgesic, antipyretic, and antiinflammatory actions, include:
  - Ibuprofen
  - Naproxen
  - Indomethacin
  - Ketorolac
  - Sulindac
- Comparisons with ASA:
  - Analgesia: ketorolac > ibuprofen/naproxen > ASA
  - Gastrointestinal irritation: < ASA, but still occurs (consider misoprostol)
  - Minimal effects on acid-base balance; no effects on uric acid elimination
  - Allergy: common, possible cross-hypersensitivity with ASA
  - Renal: chronic use may cause nephritis, nephritic syndrome, acute failure (via ↓ formation of PGE<sub>2</sub> and PGI<sub>2</sub>, which normally maintain GFR and RBF)—does not occur with sulindac
- Specific toxicities:
  - Indomethacin: thrombocytopenia, agranulocytosis, and > CNS effects
  - Sulindac: Stevens-Johnson syndrome, hematotoxicity

### Selective COX 2 Inhibitors: Celecoxib

- Compared with conventional NSAIDs, it is no more effective as an antiinflammatory agent.
- Primary differences are:
  - Less gastrointestinal toxicity
  - Less antiplatelet action
- However, it may possibly exert prothrombotic effects via inhibition of endothelial cell function (MI and strokes).
- Cross-hypersensitivity between celecoxib and sulfonamides

## OTHER DRUGS

### Acetaminophen

- Mechanisms
  - No inhibition of COX in peripheral tissues and lacks significant antiinflammatory effects
  - Equivalent analgesic and antipyretic activity to ASA due to inhibition of cyclooxygenases in the CNS
- Comparisons with ASA:
  - No antiplatelet action
  - Not implicated in Reye syndrome
  - No effects on uric acid
  - Not bronchospastic (safe in NSAID hypersensitivity and asthmatics)
  - Gastrointestinal distress is minimal at low to moderate doses
- Overdose and management:
  - Hepatotoxicity—Acetaminophen is metabolized mainly by liver glucuronyl transferase to form the inactive conjugate. A minor pathway (via P450) results in formation of a reactive metabolite (*N*-acetylbenzoquinoneimine), which is inactivated by glutathione (GSH). In overdose situations, the finite stores of GSH are depleted. Once this happens, the metabolite reacts with hepatocytes, causing nausea and vomiting, abdominal pain, and ultimately liver failure due to centrilobular necrosis. Chronic use of ethanol enhances liver toxicity via induction of P450.
  - Management of the hepatotoxicity: *N*-acetylcysteine (supplies –SH groups), preferably within the first 12 hours (*N*-acetylcysteine is also used as a mucolytic for cystic fibrosis)

### Clinical Correlate

#### “Tot” Toxicity

Young children are gustatory explorers. Among the compounds responsible for toxicity in youngsters under the age of 3 years are three items commonly found in households with “tots”: aspirin, acetaminophen (people know about Reye syndrome!), and supplementary iron tablets.

**Chapter Summary**

**Eicosanoid Pharmacology**

- \* Eicosanoids are synthesized and released on demand to interact with specific G-protein-coupled receptors. They include the leukotrienes (LTs), prostaglandins (PGs), and thromboxanes (TxAs).
- \* Figure VI-4-1 presents the pathways for the synthesis of PGI<sub>2</sub>, PGE<sub>2</sub>, PGF<sub>2α</sub>, TXA<sub>2</sub>, and the leukotrienes from the membrane phospholipids. It also shows the sites of action of the glucocorticoids, NSAIDs, COX 2 inhibitors, zileuton, and zafirlukast.
- \* The physiologic functions of relevant eicosanoids interacting with specific receptor types and the clinical aspects of the drugs affecting these actions are considered.

**Nonsteroidal Antiinflammatory Drugs**

- \* There are more than 20 nonsteroidal antiinflammatory drugs (NSAIDs) in use. Acetylsalicylic acid (ASA), the prototype, like most other NSAIDs, is a nonselective inhibitor of the cyclooxygenases; however, it binds in an irreversible fashion, whereas the others do so in a reversible manner.
- \* Progressively higher doses of ASA cause antiplatelet aggregation, analgesia, antipyresis, and antiinflammation. The mechanisms responsible for each of these responses, modes of excretion, effects on the acid-base balance, and side effects are discussed.
- \* Aspirin overdoses can cause vasomotor collapse and renal failure. The management of such toxic overdose cases is considered, as are the doses required to elicit such dangerous effects in adults and children.
- \* Other NSAIDs, including ibuprofen, naproxen, indomethacin, ketorolac, and sulindac, also have analgesic, antipyretic, and antiinflammatory properties. The properties of these NSAIDs are compared with those of ASA.
- \* Celecoxib is a selective inhibitor of cyclooxygenase 2 (COX 2), providing less gastrointestinal and antiplatelet activity than are imparted by the nonselective COX inhibitors.
- \* Acetaminophen is not an NSAID but an analgesic and antipyretic. Its properties are compared with those of ASA. It has the potential for creating severe liver damage.

# Drugs Used for Treatment of Rheumatoid Arthritis

**Table VI-5-1. Disease-Modifying Antirheumatic Drugs (DMARDs)**

Drug	Mechanism(s)	Side Effects
Hydroxychloroquine	Stabilizes lysosomes and ↓ chemotaxis	GI distress and visual dysfunction (cinchonism), hemolysis in G6PD deficiency
Methotrexate	Cytotoxic to lymphocytes	Hematotoxicity
Sulfasalazine	Sulfapyridine → ↓ B-cell functions; 5-ASA possibly inhibits COX	Hemolysis in G6PD deficiency
Glucocorticoids	↓ LTs, ILs, and platelet-activating factor (PAF)	ACTH suppression, cushingoid state, osteoporosis, GI distress, glaucoma
Gold salts	↓ Lysosomal and macrophage functions	Dermatitis, hematotoxicity, nephrotoxicity
Penicillamine	Suppresses T cells and circulating rheumatoid factor	Proteinuria, hematotoxicity, autoimmune disease
Leflunomide	Inhibits dihydro-orotic acid dehydrogenase (DHOD) → ↓ UMP → ↓ ribonucleotides → arrests lymphocytes in G <sub>1</sub>	Alopecia, rash, diarrhea, hepatotoxicity
Etanercept	Binds tumor necrosis factor (TNF); is a recombinant form of TNF receptor	Infections
Infliximab	Monoclonal antibody to TNF	Infections
Anakinra	IL-1 receptor antagonist	Infections

- NSAIDs are commonly used for initial management of rheumatoid arthritis (RA), but the doses required generally result in marked adverse effects.
- NSAIDs decrease pain and swelling but have no beneficial effects on the course of the disease or on bone deterioration.
- DMARDs are thought to slow disease progression.
- DMARDs may be started with NSAIDs at the time of initial diagnosis if symptoms are severe because DMARDs take 2 weeks to 6 months to work.
- Hydroxychloroquine is often recommended for mild arthritis and methotrexate (MTX) for moderate to severe RA.
- Other DMARDs are used less frequently, sometimes in combination regimens for refractory cases.

### Chapter Summary

- \* NSAIDs are commonly used to help alleviate the pain and inflammation associated with rheumatoid arthritis. However, they have no effect on the progress of the disease. Disease-modifying antirheumatic drugs (DMARDs) are used with the hope of slowing the disease progress. Table VI-5-1 summarizes the mechanisms of action and the adverse effects of the DMARDs.

# Drugs Used for Treatment of Gout

## TREATMENT OF ACUTE INFLAMMATORY EPISODES

- **Colchicine**, indomethacin, other NSAIDs (naproxen, sulindac), and intra-articular steroids
- Colchicine
  - Mechanism of action: binds to tubulin  $\rightarrow$   $\downarrow$  microtubular polymerization,  $\downarrow$  LTB<sub>4</sub>, and  $\downarrow$  leukocyte and granulocyte migration
  - Side effects:
    - Acute: include diarrhea and gastrointestinal pain
    - Longer use: hematuria, alopecia, myelosuppression, gastritis, and peripheral neuropathy

## PROPHYLAXIS OF CHRONIC GOUT

- Drug strategy: reduction of the uric acid pool
- Drugs: **allopurinol** and **probenecid**
- Action of allopurinol:
  - Prodrug (a suicide substrate) converted by xanthine oxidase, forming alloxanthine, which inhibits the enzyme  $\rightarrow$   $\downarrow$  purine metabolism  $\rightarrow$   $\downarrow$  uric acid.
  - Also used in cancer chemotherapy and radiation therapy





Figure VI-6-1. Mechanism of Action of Allopurinol

- Side effects of allopurinol: rash, hypo [xanthine] stones
- Drug interaction: inhibits 6-mercaptopurine (6-MP) metabolism



**Figure VI-6-2. Drug Interaction Between Allopurinol and 6-Mercaptopurine**

- Action of probenecid:
  - Inhibits proximal tubular reabsorption of urate, but ineffective if GFR <50 mL/min
  - Also inhibits secretion of many acidic drugs, e.g., cephalosporins, fluoroquinolones
- Side effects of probenecid:
  - Crystallization if high excretion of uric acid
  - ASA may ↓ effects

### Chapter Summary

- \* Acute inflammatory episodes are treated with colchicine, NSAIDs, and intraarticular steroids. The mode of colchicine's action and its adverse effects are considered.
- \* Chronic gout is treated with allopurinol, a suicide inhibitor of xanthine oxidase. The goal is to reduce the uric acid pool by inhibiting its formation from purines. The adverse effects of allopurinol are considered.
- \* Probenecid decreases the uric acid pool by inhibiting the proximal tubular reabsorption of urate. Its use and side effects are also discussed.

# Glucocorticoids

Table VI-7-1. Synthetic Derivatives of Cortisol

Drugs	Glucocorticoid Activity	Mineralocorticoid Activity	Duration
Cortisol	1	1	Short
Prednisone	4	0.3	Medium
Triamcinolone	5	0	Intermediate
Betamethasone	25	0	Long-acting
Dexamethasone	30	0	Long-acting

- Mechanisms of action:
  - Cellular effects
    - ↓ leukocyte migration
    - ↑ lysosomal membrane stability → ↓ phagocytosis
    - ↓ capillary permeability
  - Biochemical actions
    - Inhibit  $PLA_2$  (via lipocortin expression) → ↓ PGs and ↓  $LTs$
    - ↓ expression of COX 2
    - ↓ platelet-activating factor
    - ↓ interleukins (e.g., IL-2)
- Uses: antiinflammatory and immunosuppressive
- Side effects:
  - Suppression of ACTH: cortical atrophy, malaise, myalgia, arthralgia, and fever—may result in a shock state with abrupt withdrawal
  - Iatrogenic cushingoid syndrome → fat deposition, muscle weakness/atrophy, bruising, acne
  - Hyperglycemia due to ↑ gluconeogenesis → increased insulin demand and other adverse effects
  - Osteoporosis: vertebral fractures—aseptic hip necrosis
  - ↑ gastrointestinal acid and pepsin release → ulcers, gastrointestinal bleeding
  - Electrolyte imbalance:  $Na^+$ /water retention → edema and hypertension, hypokalemic alkalosis, hypocalcemia

## Clinical Correlate

### Minimize Steroidal Toxicity

- Alternate-day therapy; local application (e.g., aerosols)
- Dose-tapering to avoid cortical suppression



- ↓ skeletal growth in children
- ↓ wound healing, ↑ infections (e.g., thrush)
- ↑ glaucoma, ↑ cataracts (via ↑ sorbitol)
- ↑ mental dysfunction

### Chapter Summary

- \* Synthetic derivatives of cortisol are often used to manage inflammatory conditions or to promote immunosuppression. This chapter discusses the duration of action of several antiinflammatory steroids, their cellular effects and biochemical actions, as well as the many and severe adverse effects.

# Drugs Used for Treatment of Asthma

- Asthma is an inflammatory disease associated with bronchial hyperactivity (BHR), bronchospasm, ↑ mucus secretion, edema, and cellular infiltration.
- Early asthmatic responses (EAR) lasting from 30 to 60 minutes are associated with bronchospasm from the actions of released histamine and leukotrienes.
- Late asthmatic responses (LAR) involve infiltration of eosinophils and lymphocytes into airways → bronchoconstriction and inflammation with mucous plugging.
- Management of asthma includes bronchodilators to provide short-term relief and antiinflammatory agents that reduce bronchial hyperactivity and protect against cellular infiltration.




Figure VI-8-1. Drug Actions on Bronchiolar Smooth Muscle

## BETA-RECEPTOR AGONISTS

- Beta-2 selective drugs (albuterol, metaproterenol, terbutaline) are widely used for relief of acute bronchoconstriction and in prophylaxis of exercise-induced asthma (*see* Figure VI-8-1).
- Longer-acting drugs (e.g., salmeterol) may decrease nighttime attacks (prophylaxis only) and permit dosage reduction of other agents.
- Aerosolic forms have low potential for systemic toxicity but may cause anxiety, muscle tremors, and cardiovascular toxicity with overuse.

## MUSCARINIC-RECEPTOR BLOCKERS

- Ipratropium and other M blockers used via inhalation cause bronchodilation in acute asthma, especially in COPD patients, and they may be safer than  $\beta$  agonists are in patients with cardiovascular disease.
- They are the drugs of choice in bronchospasm caused by  $\beta$  blockers.
- There are minor atropine-like effects.

## THEOPHYLLINE

- Bronchodilates via inhibition of phosphodiesterase (PDE)  $\rightarrow$   $\uparrow$  cAMP and also by antagonism of adenosine (a bronchoconstrictor)
- Mainly adjunctive; regular use may decrease symptoms, but narrow therapeutic window predisposes to toxicity  $\rightarrow$  nausea, diarrhea, CV ( $\uparrow$  IIR, arrhythmias) and CNS excitation
- Many drug interactions; toxicity  $\uparrow$  by erythromycin, cimetidine, and fluoroquinolones
- Aminophylline IV sometimes used in bronchospasm or status asthmaticus

## CROMOLYN AND NEDOCROMIL

- Prevent degranulation of pulmonary mast cells and  $\downarrow$  release of histamine, PAF, and LTC<sub>4</sub> from inflammatory cells
- Prophylactic use:
  - $\downarrow$  symptoms and bronchial hyperactivity (BHR), especially responses to allergens
  - Minimal systemic toxicity but may cause throat irritation and cough
  - Relieved by a  $\beta_2$  agonist

## GLUCOCORTICOIDS

- Block mediator release and  $\downarrow$  BHR via  $\downarrow$  PGs, LTs, and inflammatory interleukins (ILs)
- Surface-active drugs (beclomethasone, flunisolide) used via inhalation for both acute attacks and for prophylaxis
- May cause oropharyngeal candidiasis (prevented with spacers and gargling)
- Low dosage may also prevent the desensitization of  $\beta$  receptors that can occur with overuse of  $\beta_2$  agonist
- Prednisone (oral) and IV steroids generally reserved for severe acute attacks

## ANTILEUKOTRIENES

- Zafirlukast and montelukast are antagonists at LTD<sub>4</sub> receptors with slow onset of activity used prophylactically for many forms of asthma, including antigen, exercise, or drug-induced (e.g., ASA).
- Zileuton is a selective inhibitor of lipoxygenases (LOX),  $\downarrow$  formation of all LTs. It has a more rapid onset (1–3 hours) and is adjunctive to steroids.

### Chapter Summary

- \* The management of asthma involves the use of bronchodilators to relieve short-term effects and antiinflammatories to reduce bronchial hyperactivity and protect against cellular infiltration.
- \*  $\beta_2$ -selective agonists are used for the relief of acute bronchoconstriction and as a prophylaxis in exercise-induced asthma. Longer-acting  $\beta$ -adrenoceptor agonists can be used prophylactically to decrease nighttime attacks. The mechanisms responsible for their effects are shown in Figure VI-8-1, which illustrates the action of antiasthmatic drugs.
- \* The roles of muscarinic receptor blockers, theophylline, cromolyn, nedocromil, glucocorticoids, and antileukotrienes in the treatment of asthma are discussed. Their modes of action are also illustrated in Figure VI-8-1.



# List of Drugs for Inflammatory Disorders and Practice Questions

---

## Histamine and Antihistamines

- H<sub>1</sub> antagonists: diphenhydramine, promethazine, meclizine, hydroxyzine, loratadine
- H<sub>2</sub> antagonists: cimetidine, ranitidine

## Drugs Used in Gastrointestinal Dysfunction

- Proton pump inhibitor: omeprazole
- PGE<sub>1</sub> analog: misoprostol
- Polymer: sucralfate

## Drugs Acting on Serotonergic Systems

- 5HT<sub>1a</sub> partial agonist: buspirone
- 5HT<sub>1d</sub> agonist: sumatriptan
- 5HT<sub>2</sub> antagonist: cyproheptadine, atypical antipsychotics
- 5HT<sub>3</sub> antagonist: ondansetron
- 5HT<sub>4</sub> agonist: tegaserod

## Antiemetics

- DA antagonist: metoclopramide, prochlorperazine
- H<sub>1</sub> antagonist: meclizine, promethazine
- Muscarinic antagonist: scopolamine
- Cannabinoid: dronabinol
- 5HT<sub>3</sub> antagonist: ondansetron
- NK<sub>1</sub> antagonist: aprepitant

## NSAIDs

- Aspirin, indomethacin, ibuprofen, naproxen, sulindac
- COX 2 inhibitor: celecoxib
- Acetaminophen

### Glucocorticoids

- Prednisone, triamcinolone, dexamethasone

### Drugs Used for Treatment of Gout

- Acute: colchicine, indomethacin
- Chronic: allopurinol, probenecid

### Drugs Used for Treatment of RA

- NSAIDs
- DMARDs: methotrexate, etanercept, infliximab, anakinra, and others

### Drugs Used for Treatment of Asthma

- $\beta_2$  agonists: albuterol, terbutaline
- M-blocker: ipratropium
- Methylxanthine: theophylline
- Mast-cell stabilizer: cromolyn
- Steroids: flunisolide
- LT modifiers: zafirlukast, zileuton

## Review Questions

1. A patient using NSAIDs for chronic pain develops a bleeding ulcer. What drug is designed to selectively treat ulcers of this type?
  - A. Famotidine
  - B. Bismuth
  - C. Aluminum hydroxide
  - D. Misoprostol
  - E. Muscarinic antagonists
2. Acute poisoning with acetaminophen often requires the use of a specific antidote. The beneficial property of this antidote is that it
  - A. supplies sulfhydryl groups to detoxify a reactive metabolite
  - B. induces P450 enzymes to enhance elimination
  - C. blocks the metabolism of acetaminophen
  - D. enhances renal clearance of acetaminophen
  - E. chelates acetaminophen
3. Which glucocorticoid is most likely to cause sodium and water retention?
  - A. dexamethasone
  - B. betamethasone
  - C. cortisol
  - D. celecoxib
  - E. desmopressin
4. A patient with RA is being treated with ibuprofen, but joint pain and stiffness are increasing. His physician prescribes another drug to be used with ibuprofen that may slow progression of the disease. Unfortunately, side effects develop, including dizziness, tinnitus, blurred vision, and pruritus. Ocular examination reveals corneal deposits and slight retinal pigmentation. What is the drug?
  - A. Gold salts
  - B. Etanercept
  - C. Hydroxychloroquine
  - D. Methotrexate
  - E. Thioridazine
5. A patient suffers from troublesome allergic rhinitis due to pollen, and you want to prescribe a drug for her that is least likely to cause sedation. What would your best choice be?
  - A. Betamethasone
  - B. Cimetidine
  - C. Hydroxyzine
  - D. Loratadine
  - E. Metoclopramide



6. The widely used anticoagulant warfarin is often implicated in drug interactions. If a patient takes warfarin but later begins self-medicating for ulcer pain, what drug useful for ulcers would increase the risk for bleeding?
  - A. Ranitidine
  - B. Sucralfate
  - C. Misoprostol
  - D. Cimetidine
  - E. Metoclopramide
  
7. A patient with a migraine headache is treated with sumatriptan. This drug is beneficial because it
  - A. blocks 5HT<sub>3</sub> receptors
  - B. stimulates 5HT<sub>1D</sub> receptors
  - C. blocks 5HT<sub>4</sub> receptors
  - D. stimulates 5HT<sub>2</sub> receptors
  - E. blocks muscarinic receptors
  
8. A child suffering from asthma is to be treated with a drug that blocks the synthesis of leukotrienes. What drug would be an appropriate choice?
  - A. Cromolyn
  - B. Montelukast
  - C. Ipratropium
  - D. Zileuton
  - E. Theophylline
  
9. Which one of the following is likely to be used in motion sickness, and nausea and vomiting of pregnancy?
  - A. Loratidine
  - B. Ondansetron
  - C. Meclizine
  - D. Fexofenadine
  - E. Cimetidine
  
10. For temporary maintenance of a patent ductus arteriosus prior to cardiac surgery in an infant, what is the drug of choice?
  - A. Alprostadil
  - B. Indomethacin
  - C. Epoprostenol
  - D. Celecoxib
  - E. Zileuton

11. Following an overdose of an over-the-counter drug, a young college student has marked gastrointestinal distress and is lethargic and confused, with an elevated body temperature. Lab analysis of blood reveals:  $p\text{CO}_2$ ,  $\downarrow \text{HCO}_3^-$ ,  $\downarrow \text{K}^+$ , and an anion gap acidosis. The most likely cause of these signs and symptoms is a toxic dose of
- A. acetaminophen
  - B. acetylsalicylic acid
  - C. diphenhydramine
  - D. pseudoephedrine
  - E. naproxen
12. Which statement below is accurate regarding aspirin overdose?
- A. N-acetylcysteine should be given immediately
  - B. The metabolism rate of aspirin is first-order
  - C. Elimination rate is directly proportional to plasma concentration.
  - D. Increasing urinary pH would be beneficial
  - E. Plasma concentrations decrease exponentially with time.
13. Which one of the following antiinflammatory drugs used in rheumatoid arthritis can bind directly tumor necrosis factor?
- A. Etanercept
  - B. Sulfasalazine
  - C. Prednisone
  - D. Celecoxib
  - E. Penicillamine
14. When used in the management of asthma, glucocorticoids are likely to cause
- A. hypoglycemia
  - B. decreases in blood pressure
  - C. anabolic actions in wound healing
  - D. oral thrush
  - E. sedation
15. A reasonable explanation for the therapeutic effects of ibuprofen or naproxen in primary dysmenorrhea is that these drugs
- A.  $\downarrow \text{PGE}_2$  and  $\text{PGF}_{2\alpha}$
  - B. selectively inhibit COX 2
  - C.  $\downarrow \text{LTB}_4$
  - D. inhibit  $\text{PLA}_2$
  - E.  $\text{PI}_2$

16. When a patient is started on an appropriate drug for chronic gout it is observed that the plasma levels of uric acid decrease while the urine levels of uric acid increase. What drug was the patient treated with?
- A. Allopurinol
  - B. Acetylsalicylic acid
  - C. Indomethacin
  - D. Colchicine
  - E. Probenecid
17. The plasma levels of ketoconazole are lower than normal following its oral absorption in patients treated with lansoprazole. What is the reason for this?
- A. The induction of enzymes that metabolize ketoconazole
  - B. Ketoconazole requires an acid environment for its oral absorption
  - C. Lansoprazole binds acidic drugs in the gastrointestinal tract
  - D. Increased gastrointestinal transit time because of the prokinetic effects of lansoprazole
  - E. Competition for transport mechanisms in the gastrointestinal tract
18. Cromolyn useful in many patients with asthma because it
- A. inhibits COX 2
  - B. blocks adenosine receptors in bronchiolar smooth muscle
  - C. prevents antigen-induced degranulation of mast cells
  - D. inhibits phosphodiesterase
  - E. ↓ mRNA for IL-2
19. Which one of the following is able to effectively lower intraocular pressure?
- A. Latanoprost
  - B. Ergonovine
  - C. Atropine
  - D. Terbutaline
  - E. Morphine
20. Cancer patients being treated with 6-MP may require a dosage adjustment if they are concurrently treated for which of the following?
- A. Constipation
  - B. Malaria
  - C. Chronic gout
  - D. Arthritis
  - E. Headache

21. Constipation is highly unlikely to occur with the use of which of the following?
- A. Diphenhydramine
  - B. Docusate
  - C. Promethazine
  - D. Loperamide
  - E. Scopolamine
22. A 2-year-old child is brought into the emergency department in convulsions. According to her mother, she had ingested most of a bottle of "sleeping pills," an over-the-counter preparation. What do the sleeping pills she ingested probably contain?
- A. Caffeine
  - B. Chlorpromazine
  - C. Diphenhydramine
  - D. Meperidine
  - E. Temazepam
23. In a person who regularly consumes ethanol daily, the potential for hepatotoxicity due to acetaminophen is greater than normal. What is the most likely explanation for this?
- A. Cirrhosis of the liver
  - B. Ethanol inhibits the metabolism of acetaminophen
  - C. Most beer drinkers are smokers, and nicotine sensitizes the liver to toxins
  - D. Nutritional deficiency
  - E. Ethanol induces P450 enzymes that form a toxic metabolite

### Answers

1. **Answer: D.** Misoprostol is a prostaglandin analog indicated for specific use in NSAID-induced ulcers since NSAIDs inhibit the synthesis of protective GI prostaglandins. Other answer choices may be of benefit in this type of ulcer but none are selectively used for NSAIDs.
2. **Answer: A.** Acetaminophen is metabolized primarily by glucuronidation to an inactive metabolite. A minor pathway for metabolism involves cytochrome P450 conversion of acetaminophen to a reactive metabolite that damages the liver. The reactive metabolite is rapidly inactivated normally by glutathione. Prompt administration of N-acetylcysteine is useful because, like glutathione, it supplies sulfhydryl groups to bind the reactive species.
3. **Answer: C.** Various glucocorticoids have different abilities to affect the mineralocorticoid receptor to cause sodium and water retention (an aldosterone-like effect). Generally, the more potent the glucocorticoid, the less likely it is to have an aldosterone effect. Cortisol is a weak glucocorticoid that is equally effective at stimulating mineralocorticoid receptors and thus has sodium and water retention as a property.
4. **Answer: C.** Ocular toxicity is characteristic of chloroquine and hydroxychloroquine. Corneal deposits are reversible, but retinal pigmentation can ultimately lead to blindness. Patients will complain about gastrointestinal distress, visual dysfunction, ringing in the ears (note that tinnitus also occurs in salicylism), and “itchy skin.” Hydroxychloroquine also promotes oxidative stress that can lead to hemolysis in G6PD deficiency. DMARDs include gold salts (e.g., auranofin), methotrexate, and etanercept, but thioridazine is a phenothiazine used as an antipsychotic; it lacks an antiinflammatory effect, but does cause retinal pigmentation.
5. **Answer: D.** The usual choice for pollen-induced allergies would be an H<sub>1</sub> antagonist. Of the two listed, loratadine would be the best choice in this case because it does not cross the blood–brain barrier and is non-sedating; hydroxyzine is an effective CNS depressant used for preoperative sedation. Cromolyn (not listed) can also be used in allergic rhinitis and is also non-sedating. Betamethasone, a potent antiinflammatory steroid, is less effective than antihistamines in this situation and would cause more serious side effects. Metoclopramide is a DA-receptor antagonist and prokinetic used as an antiemetic and in GERD. Cimetidine is the prototype H<sub>2</sub> antagonist used in gastrointestinal ulcers.
6. **Answer: D.** Cimetidine is an inhibitor of the hepatic cytochrome P450 isoform that metabolizes warfarin, consequently decreasing its clearance and thus increasing its elimination half-life. The hepatic metabolism of many other drugs can be inhibited by cimetidine, possibly necessitating dose reductions to avoid toxicity, including beta blockers, isoniazid, procainamide, metronidazole, tricyclic antidepressants, and phenytoin.
7. **Answer: B.** It is important to be able to match the serotonin drugs with their respective receptors. The “triptans” used in migraine headaches are agonists at the 5HT<sub>1D</sub> receptor.
8. **Answer: D.** Zileuton blocks the enzyme 5-lipoxygenase which prevents the formation of leukotrienes. This drug is one of many adjuncts available in asthma. Montelukast blocks leukotriene receptors but has no effect on the synthesis of leukotrienes.

9. **Answer: C.** Meclizine is a first-generation antihistamine that effectively penetrates the CNS. Like all first-generation drugs it also blocks muscarinic receptors. Blocking the  $H_1$  and muscarinic receptors are beneficial in motion sickness and nausea and vomiting in pregnancy. Second-generation drugs like loratidine and fexofenadine don't effectively penetrate the CNS and are of no benefit in these conditions.
  
10. **Answer: A.** During fetal development, the ductus arteriosus is kept open by prostaglandins. For temporary maintenance of patency in the infant, the  $PGE_1$  analog alprostadil is used. Closure of the ductus in the infant can often be accomplished by intravenous indomethacin, which  $\downarrow$  PG synthesis by inhibiting COX. Epoprostenol is a prostacyclin analog used in primary pulmonary hypertension.
  
11. **Answer: B.** If the patient had been able to mention tinnitus, this would be a classic case of aspirin poisoning. At high salicylate blood levels, the combination of effects leading to respiratory depression (respiratory acidosis) and metabolic acidosis results in the observed pH and electrolyte changes, the anion gap (a marker for acidosis), and hyperthermia.
  
12. **Answer: D.** Back to basic principles. Zero-order elimination means that plasma levels of a drug decrease linearly with time. This occurs with ASA at toxic doses, with phenytoin at high therapeutic doses, and with ethanol at all doses. Enzymes that metabolize ASA are saturated at high plasma levels  $\rightarrow$  constant rate of metabolism = zero-order kinetics. Remember that application of the Henderson-Hasselbalch principle can be important in drug overdose situations. In the case of aspirin, a weak acid, urinary alkalization favors ionization of the drug  $\rightarrow$   $\downarrow$  tubular reabsorption  $\rightarrow$  renal elimination. N-acetylcysteine is the antidote for acetaminophen.
  
13. **Answer: A.** Etanercept binds directly to tumor necrosis factor (TNF), resulting in the inactivation of this cytokine, which plays a major role in a number of inflammatory disorders, including Crohn disease and rheumatoid arthritis. In the synovium, TNF recruits inflammatory cells and leads to angiogenesis and joint destruction. Infliximab, a monoclonal antibody, also inactivates TNF.
  
14. **Answer: D.** Most often glucocorticoids are used in the treatment of asthma not controlled by a beta-2 agonist inhaler alone. The glucocorticoid is often given by metered dose inhaler when enhances the risk of oral candidiasis (thrush). This can be avoided by rinsing the mouth thoroughly and by using spacers. All of the other effects listed are "opposites," so anticipate possible hyperglycemia, hypertension, decreased wound healing, and CNS excitatory effects that have been interpreted as psychosis.
  
15. **Answer: A.**  $PGE_2$  and  $PGF_{2\alpha}$  both increase in primary dysmenorrhea, and the therapeutic effects of NSAIDs appear to be due to inhibition of the synthesis of these prostaglandins. Both ibuprofen and naproxen are nonselective COX inhibitors that can inhibit the synthesis of prostacyclin ( $PGI_2$ ). NSAIDs do not inhibit phospholipase  $A_2$ , and they do not decrease leukotrienes.

16. **Answer: E.** In chronic gout, the strategy is to decrease uric acid formation from purines by inhibiting xanthine oxidase with allopurinol or increasing urate elimination with uricosurics such as probenecid. Probenecid blocks the tubular reabsorption of uric acid which lowers blood levels of uric acid but results in uricosuria. Colchicine and NSAIDs are less effective and cause more side effects when used in chronic gout. There are preferred in acute gout attacks. Although ASA is uricosuric at antiinflammatory doses, its toxicity makes the drug a poor choice.
17. **Answer: B.** Several drugs, including ketoconazole and fluoroquinolones, require an acidic environment in the gastrointestinal tract for effective absorption into the systemic circulation. Drugs used in treatment of gastrointestinal ulcers such as proton pump inhibitors (lansoprazole) commonly increase gastric pH, leading to the ↓ absorption of such drugs and, consequently, a ↓ in their effects.
18. **Answer: C.** Cromolyn is a mast-cell stabilizer used in asthma (especially antigen-induced) and in food allergies. Inhibition of degranulation with decreased release of histamine and eicosanoids contributes to its antiinflammatory effectiveness in asthma, where it is used for prophylaxis. Methylxanthines, such as theophylline, exert bronchodilating effects via their inhibition of phosphodiesterases and their antagonism of adenosine receptors. Steroids used in asthma ↓ bronchial hyperactivity by several mechanisms, including inhibition of interleukin synthesis. COX 2 inhibitors have no established role in asthma management.
19. **Answer: A.** Latanoprost is a prostaglandin  $F_{2\alpha}$  analog that is used in glaucoma to lower intraocular pressure. Ergonovine causes smooth muscle contraction (both uterine and vascular) and is used for control of postpartum hemorrhage. Atropine has the potential to raise intraocular pressure and precipitate glaucoma. Neither a  $\beta_2$  agonist (terbutaline) nor opioids (morphine) is useful in glaucoma.
20. **Answer: C.** Allopurinol is a uricosuric drug used in chronic gout that prevents formation of uric acid from purines by acting as a suicide substrate of xanthine oxidase. The drug is commonly used in patients undergoing treatment of cancer to slow down formation of uric acid derived from purines released by the cytotoxic action of drugs or radiation. The metabolism of 6-mercaptopurine (6-MP), a substrate for xanthine oxidase, is also inhibited by allopurinol, necessitating a major dose reduction to avoid its toxic effects.
21. **Answer: B.** Docusate is a stool-softening laxative that facilitates mixing of oil and water via its surfactant properties. Drugs that have muscarinic blocking effects, such as scopolamine and the antihistamines diphenhydramine and promethazine, tend to cause constipation by decreasing gastrointestinal motility. Loperamide is an opioid derivative, with no analgesic activity, used in the treatment of diarrheal states.
22. **Answer: C.** Over-the-counter (OTC) sleep aids invariably contain sedating antihistamines such as diphenhydramine. Sometimes called sedative-autonomics, overdoses of such drugs are dangerous, especially in small children. They usually have muscarinic-blocking (atropine-like) effects causing hyperthermia, and they lower the seizure threshold, leading to convulsions. Chlorpromazine is very similar in its pharmacology but is not available OTC and would not be appropriate as a sleeping aid because of its autonomic side effects. Temazepam, a benzodiazepine, is used as a sleeping pill but requires a prescription and raises the seizure threshold. Meperidine is an opioid-analgesic that can cause seizures in OD, but it is not used as a sleeping aid or available OTC. Caffeine is a CNS stimulant.

23. **Answer: E.** Ethanol has mixed effects on liver metabolism of drugs. Acutely, it can act as an enzyme inhibitor, but chronic use may lead to enzyme induction. Acetaminophen is metabolized mainly via conjugation reactions, but a minor pathway involving P-450 (probably the CYP2E1 isoform) results in formation of small amounts of the reactive metabolite, which is (normally) rapidly inactivated by GSH. The chronic ingestion of more than average amounts of ethanol induces the formation of the P-450 isozyme that converts acetaminophen to its reactive metabolite. Thus, more-than-normal amounts of *N*-acetyl-benzoquinoneimine would be formed in an overdose situation, resulting in enhanced hepatotoxicity.





**SECTION VII**

**Drugs Used in  
Blood Disorders**



# Anticoagulants

## OVERVIEW

Blood coagulates by transformation of soluble fibrinogen into insoluble fibrin. Circulating proteins interact in a “cascade,” where clotting factors undergo limited proteolysis to become active serine proteases. Anticoagulants are drugs that decrease the formation of fibrin clots. Oral anticoagulants (e.g., warfarin) inhibit the hepatic synthesis of clotting factors II, VII, IX, and X. Heparin inhibits the activity of several activated clotting factors (especially factors IIa and Xa) via its activation of antithrombin III. The endogenous anticoagulants, protein C and protein S, cause proteolysis of factors Va and VIIIa.




Figure VII-1-1. Actions of Blood Drugs

## COMPARATIVE PROPERTIES OF HEPARIN AND WARFARIN

Table VII-1-1. Properties of Heparin and Warfarin (Coumarins)

Feature	Heparin(s)	Warfarin (Coumarins)
Chemical nature	Large polysaccharide, water-soluble	Small molecule, lipid-soluble derivatives of vitamin K
Kinetics	Given parenterally (IV, SC), hepatic and reticuloendothelial elimination, half-life = 2 h, no placental access	Given orally, 98% protein bound, PO, liver metabolism, half-life = 30+ h, placental access
Mechanism	Heparin catalyzes the binding of antithrombin III (a serine protease inhibitor) to factors IIa, IXa, Xa, XIa, and XIIa, resulting in their rapid inactivation	↓ Hepatic synthesis of vitamin K-dependent factors II, VII, IX, X—coumarins prevent $\gamma$ -carboxylation by inhibiting vitamin K epoxide reductase; no effect on factors already present. <i>In vivo</i> effects only
Monitoring	Partial thromboplastin time (PTT)	Prothrombin time (PT); INR
Antagonist	Protamine sulfate—chemical antagonism, fast onset	Vitamin K—↑ cofactor synthesis, slow onset; fresh frozen plasma (fast)
Uses	Rapid anticoagulation (intensive) for thromboses, emboli, unstable angina, disseminated intravascular coagulation (DIC), open-heart surgery, etc.	Longer-term anticoagulation (controlled) for thromboses, emboli, post-MI, heart valve damage, atrial arrhythmias, etc.
Toxicity	Bleeding, osteoporosis, heparin-induced thrombocytopenia (HIT), hypersensitivity	Bleeding, skin necrosis (if low protein C), drug interactions, teratogenic (bone dysmorphism)

### Note

- Argatroban is a direct thrombin inhibitor.
- It does not require ATIII.
- It does not interact with heparin-induced antibodies.
- It can be used in HIT.

### Heparins

- Heparin is a mixture of sulfated polysaccharides with molecular weights of 15–20,000 daltons.
- Low-molecular-weight (LMW) heparins (e.g., enoxaparin) have potential advantage of longer half-lives, less thrombocytopenia, and possibly enhanced activity against factor Xa.
- Danaparoid, a heparin of different structure, may be safer in hypersensitivity to heparin.

### Warfarin

- Drug interactions:
  - Acidic molecule: oral absorption ↓ by cholestyramine
  - Extensive (but weak) plasma protein binding: displacement by other drugs may increase free fraction → ↑ PT (e.g., ASA, sulfonamides, phenytoins)

- Slow hepatic metabolism via P450:
  - Inducers (barbiturates, carbamazepine, rifampin) → ↓ PT
  - Inhibitors (cimetidine, macrolides, azole antifungals) → ↑ PT
- Protein C deficiency:




Figure VII-1-2. Activation and Role of Protein C

Table VII-1-2. Coagulation Factor Half-Lives

Factor	IIa	VIIa	IXa	Xa	C
Half-life (h)	60	8	24	40	14

- Transient protein C deficiency can be induced when initiating treatment with warfarin because factors VII and protein C have the shortest half-lives of the coagulation factors (Table VII-1-2).
- Consequently, the extrinsic pathway and protein C system are inactivated, whereas the intrinsic system remains active for a few days. Hypercoagulability occurs (Figure VII-1-2), which may result in dermal vascular thrombosis and skin necrosis.

### Bivalirudin

- Thrombin-specific anticoagulant
- Used with aspirin in unstable angina when undergoing percutaneous transluminal coronary angioplasty (PTCA)

**Chapter Summary**

- \* Table VII-1-1 summarizes the properties of heparin and warfarin (a coumarin).
- \* Low-molecular-weight heparin derivatives (e.g., enoxaparin) and danaparoid, a heparan with heparin-like properties, have potential advantages over heparin itself.
- \* The drug interactions of warfarin are given.
- \* The activation and role of protein C in the clotting cascade are illustrated in Figure VII-1-2. Transient protein C deficiency can be induced by treatment with warfarin, which promotes hypercoagulation through the action of the intrinsic pathway.

# Thrombolytics




Also called fibrinolytics, these agents lyse thrombi by catalyzing the formation of the endogenous fibrinolytic plasmin (a serine protease) from its precursor, plasminogen.

Thrombolytics include tissue plasminogen activator (tPA, recombinant) and streptokinase (bacterial). They are used intravenously for short-term emergency management of coronary thromboses in myocardial infarction (MI), deep venous thrombosis, pulmonary embolism, and ischemic stroke (tPA).

Drugs:

- Streptokinase
  - Acts on both bound and free plasminogen (not clot specific), depleting circulating plasminogen and factors V and VIII
  - Is antigenic (foreign protein derived from  $\beta$ -hemolytic streptococci); may cause a problem if recent past use or infection—strep antibodies may  $\downarrow$  activity
- Alteplase (tPA)
  - Clot specific, acting mainly on fibrin-bound plasminogen, the natural activator, so no allergy problems



**Figure VII-2-1. Actions of Streptokinase and Alteplase**

Clinical features:

- Overriding factor in effectiveness is early administration, e.g.,  $>60\%$  decrease in mortality post-MI if used within 3 hours
- ASA, beta blockers, and nitrates further  $\downarrow$  mortality, and adenosine  $\downarrow$  infarct size
- Complications include bleeding, possible intracerebral hemorrhage
- Streptokinase may cause hypersensitivity reactions and hypotension
- Antifibrinolytics (aminocaproic and tranexamic acids)—possible antidotes in excessive bleeding



**Chapter Summary**

- \* Thrombolytics (also referred to as fibrinolytics) are of clinical value in the early treatment of fibrin-clot-induced ischemia (e.g., >60% decrease in post-MI mortality if used within 3 hours).

# Antiplatelet Drugs




Figure VII-3-1. Platelet Activation

## In A Nutshell

### Platelet Aggregation

*Increased by:*

ADP, 5HT, TxA<sub>2</sub>, thrombin, α<sub>2</sub> agonists

*Decreased by:*

PGI<sub>2</sub>, cAMP, ASA, ticlopidine, clopidogrel, gp IIb/IIIa blockers

Thrombus (clot) formation involves:

- Platelet adhesion to site of vascular injury
- Activation of platelets by factors that include TxA<sub>2</sub>, ADP, collagen, 5HT, and thrombin → ↑ expression of glycoprotein IIb/IIIa receptors
- Aggregation of platelets by a cross-linking reaction due to fibrinogen binding to glycoprotein IIb/IIIa receptors

Drugs:

- Aspirin
  - Irreversibly inhibits COX in platelets → ↓ activation
  - Low doses prevent MI and recurrence; prophylaxis in atrial arrhythmias and TIAs
  - Adverse effects (see Section VI, Drugs for Inflammatory and Related Disorders)

- Ticlopidine and clopidogrel
  - Block ADP receptors on platelets → ↓ activation
  - Alternatives to ASA in TIAs, post-MI, and unstable angina
  - Hemorrhage, leukopenia, and thrombocytopenic purpura
- Abciximab, eptifibatide, and tirofiban
  - Antagonists that bind to glycoprotein IIb/IIIa receptors → ↓ aggregation by preventing the cross-linking reaction
  - Used mainly in acute coronary syndromes and postangioplasty

### **Chapter Summary**

- \* Platelets adhere to sites of vascular injury, where they are activated by various factors to express a glycoprotein to which fibrinogen binds, resulting in platelet aggregation and formation of a platelet plug. Antiplatelet drugs inhibit this process, thus reducing the chances of thrombi formation. The major drugs are aspirin, ticlopidine, clopidogrel, abciximab, eptifibatide, and tirofiban.

# List of Drugs Used in Blood Disorders and Practice Questions

---

## Anticoagulants

- Heparin
- Warfarin
- Argatroban
- Bivalirudin

## Thrombolytics

- Alteplase (tPA)
- Streptokinase

## Antiplatelet

- Aspirin
- Ticlopidine
- Clopidogrel
- Abciximab

## Review Questions

1. Which of the following compounds is most likely to block ADP receptors and prevent platelet aggregation?
  - A. Clopidogrel
  - B. Aspirin
  - C. Prostacyclin
  - D. Abciximab
  - E. Montelukast
2. A woman who has a mechanical heart valve and who is taking warfarin informs you that she hopes to get pregnant in the near future. What advice should she receive regarding her antithrombotic medication during the anticipated pregnancy?
  - A. Warfarin should be continued until the third trimester.
  - B. Warfarin should be replaced with aspirin at analgesic doses.
  - C. All medications that affect the blood should be discontinued.
  - D. Warfarin should be replaced with heparin.
  - E. Warfarin should be discontinued, and supplementary vitamin K taken throughout the pregnancy.

3. The primary advantage of enoxaparin over heparin is that it
  - A. is unlikely to cause bleeding
  - B. more effectively inhibits the synthesis of clotting factors
  - C. has a more rapid onset
  - D. does not cause thrombocytopenia
  - E. has a longer half-life
  
4. Which of the following statements regarding warfarin is true?
  - A. It is a prodrug converted to its active metabolite spontaneously in the blood.
  - B. It has low lipophilicity and does not cross the placental barrier.
  - C. It causes a depletion in protein C before it decreases prothrombin.
  - D. It inhibits release of vitamin K–dependent clotting factors from hepatocytes.
  - E. It is inactivated by protamine.
  
5. Which of the following statements is true regarding the parenteral administration of alteplase?
  - A. It increases the formation of plasminogen.
  - B. It is less effective than streptokinase when given after a myocardial infarction.
  - C. It causes a high incidence of thrombocytopenia.
  - D. It may cause bleeding reversible by aminocaproic acid.
  - E. It activates free plasminogen.
  
6. Following a myocardial infarction, a patient is stabilized on warfarin, the dose being adjusted to give a prothrombin time of 22 seconds. Which of the following statements regarding potential drug interactions in this patient is accurate?
  - A. Cholestyramine will increase prothrombin time.
  - B. Cimetidine is likely to decrease prothrombin time.
  - C. Antibacterial sulfonamides may enhance the effects of warfarin.
  - D. Vitamin K would restore prothrombin time to normal within 30 minutes.
  - E. If this patient takes half an aspirin tablet daily, the dose of warfarin will need to be increased.

## Answers

- Answer: A.** Platelet aggregation is stimulated by many compounds, including ADP, thromboxane A<sub>2</sub>, fibrin, and serotonin. Clopidogrel, along with ticlopidine, blocks ADP receptors and prevent platelet activation. Prostacyclin (PGI<sub>2</sub>) from endothelial cells is a naturally occurring compound that inhibits platelet aggregation by stimulating PGI<sub>2</sub> receptors. Aspirin inhibits the synthesis of thromboxane A<sub>2</sub>. Abciximab is a monoclonal antibody targeted to the glycoprotein IIb/IIIa receptor which inhibits aggregation. Montelukast blocks leukotriene receptors and is used in asthma.
- Answer: D.** Discontinuing warfarin is appropriate during pregnancy because it is a known teratogen that causes bone dysmorphogenesis. The patient will need continued protection against thrombus formation, and heparin (or a related low molecular weight compound) is usually advised, despite the fact that the drug will require parenteral administration and can cause thrombocytopenia.
- Answer: E.** Enoxaparin is a low-molecular weight heparin. As such, it is smaller and will have a longer half-life compared to heparin. It still has a risk of causing bleeding and thrombocytopenia, but is not more rapid in onset. Heparins do not affect the synthesis of clotting factors, but rather rapidly inactivate existing factors.
- Answer: C.** Warfarin inhibits the hepatic synthesis of factors II (prothrombin), VII, IX, and X. Its onset of anticoagulation activity is slow, and its impact on individual coagulation factors depends on their half-lives. Factor VII and protein C have much shorter half-lives than prothrombin, and so the extrinsic pathway and protein C system are the first to be affected by warfarin. The intrinsic pathway continues to function for 2 to 3 days, causing a state of hypercoagulability and possible vascular thrombosis.
- Answer: D.** Alteplase is thrombolytic (or “fibrinolytic”) because it activates plasminogen, resulting in the increased formation of plasmin. Its efficacy is equivalent to that of streptokinase, but alteplase has the advantage of only activating plasminogen bound to fibrin (clot specific) but not free plasminogen. All thrombolytics can cause bleeding, which may be counteracted to some extent by administration of antifibrinolytics, such as aminocaproic acid.
- Answer: C.** Warfarin binds extensively (98%) but weakly to plasma proteins and can be displaced by other drugs (e.g., ASA, chloral hydrate, phenytoin, sulfinpyrazone, and sulfonamides), resulting in an increase in its anticoagulant effects. Bile acid sequestrants bind acidic drugs such as warfarin, preventing their gastrointestinal absorption (↓ prothrombin time [PT]), and cimetidine, which inhibits the metabolism of warfarin, causing an increase in PT. Vitamin K restores levels of prothrombin and several other coagulation factors, but the action is slow (24 to 48 hours). Due to antiplatelet effects, even low doses of ASA may enhance bleeding in patients on warfarin.



**SECTION VIII**

**Endocrine  
Pharmacology**





# Drugs Used in Diabetes

## INSULINS

### Diabetes Mellitus

- Type 1 (IDDM):
  - Early onset
  - Loss of pancreatic B cells → absolute dependence on insulin (diet + insulin ± oral agents)
  - Ketoacidosis-prone
- Type 2 (NIDDM)
  - Usually adult onset
  - ↓ response to insulin → (diet → oral hypoglycemics ± insulin)
  - Not ketoacidosis-prone

### Insulin Forms

Table VIII-1-1. Kinetics (in Hours) of Insulin Forms with Subcutaneous Injection

Form	Onset	Peak Effect	Duration
Lispro*	0.3–0.5	1–2	3–4
Regular*	0.5–1	2–4	5–7
Glargine	1	no peak	≥24

\*Only forms that can be used intravenously; peak action in 2 to 4 min.

- Glargine:
  - Insulin analog with no peak (“peakless,” i.e., broad plasma concentration plateau)
  - Ultralong duration of action
  - Used to supply a constant background level
- Mechanism: insulin binds to transmembrane receptors which activate tyrosine kinase to phosphorylate tissue-specific substrates

### In A Nutshell

#### Insulin Release

Increased by:

- Glucose
- Sulfonylureas
- M-agonists
- β<sub>2</sub>-agonists

Decreased by:

- α<sub>2</sub>-agonists

### Clinical Correlate

#### Diabetic Ketoacidosis

- Symptoms: polyuria, polydipsia, nausea, fatigue, dehydration, Kussmaul breathing, “fruity” breath
- Treatment: regular insulin IV, fluid and electrolyte replacement

## ORAL HYPOGLYCEMICS: SULFONYLUREAS

### Note

#### Hypoglycemic Reactions

- Symptoms: lip/tongue tingling, lethargy, confusion, sweats, tremors, tachycardia, coma, seizures
- Treatment: oral glucose, IV dextrose if unconscious, or glucagon (IM or inhalation)




Figure VIII-1-1. Mode of Action of Sulfonylureas

- Mechanisms:
  - Normally, K<sup>+</sup> efflux in pancreatic  $\beta$  cells maintains hyperpolarization of membranes, and insulin is released only when depolarization occurs.
  - Glucose acts as an insulinogen by increasing intracellular ATP  $\rightarrow$  closure of K<sup>+</sup> channels  $\rightarrow$  membrane depolarization  $\rightarrow$   $\uparrow$  Ca<sup>2+</sup> influx  $\rightarrow$  insulin release.
  - The acute action of sulfonylureas is to block K<sup>+</sup> channels  $\rightarrow$  depolarization  $\rightarrow$  insulin release.
- Effects of increased insulin:
  - $\rightarrow$   $\downarrow$  glucagon release from pancreatic  $\alpha$  cells
  - Continued use of sulfonylureas  $\uparrow$  tissue responses to insulin (especially muscle and liver) via changes in receptor function
- Drugs:
  - First generation:
    - Acetohexamide (active metabolite,  $\downarrow$  dose in renal dysfunction)
    - Tolbutamide (appropriate in renal dysfunction)
    - **Chlorpropamide** (long-acting, SIADH/disulfiram reactions)
  - Second generation:
    - Glipizide ( $\downarrow$  dose in hepatic dysfunction)
    - Glyburide (active metabolite,  $\downarrow$  dose in renal dysfunction)
- Side effects:
  - Hypoglycemia
  - Weight gain
  - Drug interactions mainly with first-generation drugs  $\rightarrow$   $\uparrow$  hypoglycemia with cimetidine, insulin, salicylates, sulfonamides

### Note

#### Repaglinide

- Mechanisms: stimulates insulin release from pancreatic beta cells
- Use: adjunctive use in type 2 diabetes—administer just before meals due to short half-life

## METFORMIN


- “Euglycemic,” ↓ postprandial glucose levels, but does not cause hypoglycemia or weight gain
- Mechanisms: may involve ↑ tissue sensitivity to insulin and/or ↓ hepatic gluconeogenesis (Figure VIII-1-2)
- Use: monotherapy or combinations (synergistic with sulfonylureas)
- Side effects: possible lactic acidosis; gastrointestinal distress is common

## ACARBOSE

- No hypoglycemia
- Mechanisms: inhibits α-glucosidase in brush borders of small intestine → ↓ formation of absorbable carbohydrate → ↓ postprandial glucose → ↓ demand for insulin (Figure VIII-1-2).
- Side effects: gastrointestinal discomfort, flatulence, and diarrhea—recent concern over potential hepatotoxicity

## THIAZOLIDINEDIONES: PIOGLITAZONE AND ROSIGLITAZONE

- Mechanisms: bind to nuclear peroxisome proliferator-activating receptors (PPARs) involved in transcription of insulin-responsive genes → sensitization of tissues to insulin, plus ↓ hepatic gluconeogenesis and triglycerides and ↑ insulin receptor numbers (Figure VIII-1-2).
- Side effects: less hypoglycemia than sulfonylureas, but weight gain and edema reported



\* osu = oral sulfonylureas

Figure VIII-1-2. Modes of Action of Drugs Used to Treat Diabetes

## AGENTS AFFECTING GLUCAGON-LIKE PEPTIDE-1 (GLP-1)

### Exenatide

- Mechanism: GLP-1 is an incretin released from the small intestine. It augments glucose-dependent insulin secretion. Exenatide is a long-acting GLP-1 receptor full agonist used in combination with other agents in type 2 diabetes.
- Side effects: nausea, hypoglycemia when used with oral sulfonylureas

### Note

Pramlintide is a synthetic version of amylin that slows the rate at which food is absorbed from the intestine, decreases glucose production, and decreases appetite. It is used in type 1 and type 2 diabetes.

### Sitagliptin

- Mechanism: inhibits dipeptidyl peptidase (DPP-4) thereby inhibiting the inactivation of GLP-1




Figure VIII-1-3. Action of Drugs Affecting GLP-1

## Chapter Summary

- \* Type 1 (IDDM) and type 2 (NIDDM) diabetes mellitus are defined at the beginning of the chapter.
- \* The times of activity onset, peak activity, and duration of activity for lispro, regular, lente, and ultralente insulins are summarized in Table VIII-1-1.
- \* The oral antidiabetic drugs are the sulfonylureas, metformin, acarbose, thiazolidinediones, and repaglinide.
- \* By blocking  $K^+$  channels in the pancreatic cells, the sulfonylureas stimulate insulin release. The extra insulin in turn inhibits glucagon release from the cells and increases peripheral tissue sensitivity to insulin (Figure VIII-1-1). The first- and second-generation drugs are listed. The adverse effects include weight gain and potential hypoglycemia.
- \* Metformin enhances tissue sensitivity to insulin and inhibits liver gluconeogenesis. The potential side effect is lactic acidosis.
- \* Acarbose inhibits intestinal-glucosidase, thereby slowing glucose absorption and decreasing insulin demand. The side effect is gastrointestinal distress.
- \* The thiazolidinediones (glitazones) act via peroxisome proliferation activating receptors that control insulin-responsive genes. They are less hypoglycemic than the sulfonylureas, but they still induce weight gain and edema and have potential liver toxicity.
- \* Repaglinide, like the sulfonylureas, stimulates-cell secretion of insulin. Figure VIII-1-2 summarizes the modes of action of these drugs.
- \* Both exenatide and sitagliptin increase glucose-dependent insulin secretion.

# Steroid Hormones

---

## ADRENAL STEROIDS

- Nonendocrine uses: For use in inflammatory disorders (and accompanying adverse effects), *see* Section VI, Drugs for Inflammatory and Related Disorders.
- Endocrine uses of glucocorticoids (e.g., prednisone, dexamethasone) and the mineralocorticoid (fludrocortisone) include:
  - Addison disease—replacement therapy
  - Adrenal insufficiency states (infection, shock, trauma)—supplementation
  - Premature delivery to prevent respiratory distress syndrome—supplementation
  - Adrenal hyperplasia—feedback inhibition of ACTH
- Adrenal steroid antagonists:
  - Spironolactone
  - Blocks aldosterone and androgen receptors (*see* Section III, Cardiac and Renal Pharmacology)
- Mifepristone:
  - Blocks glucocorticoid and progestin receptors
- Synthesis inhibitors:
  - Metyrapone (blocks 11-hydroxylation)
  - Ketoconazole

## ESTROGENS

- Pharmacology: Estradiol is the major natural estrogen. Rationale for synthetics is to ↑ oral bioavailability, ↑ half-life, and ↑ feedback inhibition of FSH and LH.
- Drugs:
  - Conjugated equine estrogens (Premarin)—natural
  - Ethinyl estradiol and mestranol—steroidal
  - Diethylstilbestrol (DES)—nonsteroidal
- Clinical uses:
  - Female hypogonadism
  - Hormone replacement therapy (HRT) in menopause → ↓ bone resorption (↓ PTH)
  - Contraception—feedback ↓ of gonadotropins
  - Dysmenorrhea
  - Uterine bleeding
  - Acne
  - Prostate cancer (palliative)

- Side effects:
  - General
    - Nausea
    - Breast tenderness
    - Endometrial hyperplasia
    - ↑ gallbladder disease, cholestasis
    - Migraine
  - ↑ blood coagulation—via ↓ antithrombin III and ↑ factors II, VII, IX, and X (only at high dose)
  - Cancer risk
    - ↑ endometrial cancer (unless progestins are added)
    - ↑ breast cancer—questionable, but caution if other risk factors are present
    - DES given during breast feeding → ↑ vaginal adenocarcinoma cancer in offspring
- Other drugs:
  - Anastrozole
    - Mode of action: aromatase inhibitor → ↓ estrogen synthesis
    - Use: estrogen-dependent, postmenopausal breast cancer




Figure VIII-2-1. Mechanism of Action of Anastrozole

- **Clomiphene** (fertility pill)
  - Mode of action: ↓ feedback inhibition → ↑ FSH and LH → ↑ ovulation → pregnancy
  - Use: fertility drug
  - Adverse effect: ↑ multiple births
- Selective estrogen-receptor modulators (SERMs):
  - **Tamoxifen**
    - Variable actions depending on “target” tissue
    - E-receptor agonist (bone), antagonist (breast), and partial agonist (endometrium)
    - Possible ↑ risk of endometrial cancer
    - Used in estrogen-dependent breast cancer and for prophylaxis in high-risk patients
  - Raloxifene
    - E-receptor agonist (bone), antagonist breast and uterus
    - When used in menopause, there is no ↑ cancer risk
    - Use: prophylaxis of postmenopausal osteoporosis

Table VIII-2-1. Comparison of Tamoxifen and Raloxifene in Various Tissues

Drug	Bone	Breast	Endometrium
Tamoxifen	Agonist	Antagonist	Agonist
Raloxifene	Agonist	Antagonist	Antagonist

## PROGESTIN

- Pharmacology: Progesterone is the major natural progestin. Rationale for synthetics is ↑ oral bioavailability and ↑ feedback inhibition of gonadotropins, especially luteinizing hormone (LH).
- Drugs:
  - Medroxyprogesterone
  - Norethindrone
  - Desogestrel is a synthetic progestin devoid of androgenic and antiestrogenic activities, common to other derivatives
- Clinical uses:
  - Contraception (oral with estrogens)—depot contraception (medroxyprogesterone IM every 3 months)
  - Hormone replacement therapy (HRT)—with estrogens to ↓ endometrial cancer
- Side effects:
  - ↓ HDL and ↑ LDL
  - Glucose intolerance
  - Breakthrough bleeding
  - Androgenic (hirsutism and acne)
  - Antiestrogenic (block lipid changes)
- Antagonist: mifepristone—abortifacient (use with prostaglandins [PGs])

## ORAL CONTRACEPTIVES

- Pharmacology:
  - Combinations of estrogens (ethinyl estradiol, mestranol) with progestins (norgestrel, norethindrone) in varied dose, with mono-, bi-, and triphasic variants
  - Suppress gonadotropins, especially midcycle LH surge
- Side effects:
  - Estrogens
    - Nausea
    - Bloating
    - Headache
    - Mastalgia
  - Progestins
    - Weight gain
    - Hirsutism
    - Acne



- Tiredness
- Depression
- ↓ HDL and ↑ LDL (high progestins)
- Interactions: ↓ contraceptive effectiveness when used with antimicrobials and enzyme inducers
- Benefits:
  - ↓ risk of endometrial and ovarian cancer
  - ↓ dysmenorrhea
  - ↓ endometriosis
  - ↓ pelvic inflammatory disease (PID)
  - ↓ osteoporosis

## ANDROGENS

- Pharmacology: include methyltestosterone and 17-alkyl derivatives with increased anabolic actions, e.g., oxandrolone, nandrolone
- Uses:
  - Male hypogonadism and for anabolic actions → ↑ muscle mass, ↑ RBCs, ↓ nitrogen excretion
  - Illicit use in athletics
- Side effects:
  - Excessive masculinization
  - Premature closure of epiphysis
  - Cholestatic jaundice
  - Aggression
  - Dependence
- Antagonists:
  - **Flutamide:** androgen receptor blocker—used for androgen-receptor-positive prostate cancer
  - **Leuprolide:** GnRH analog—repository form used for androgen-receptor-positive prostate cancer
  - **Finasteride**
    - 5-Alpha reductase inhibitor, preventing conversion of testosterone to dihydrotestosterone (DHT)
    - DHT is responsible for hair loss and prostate enlargement
    - Uses: BPH, male pattern baldness
    - Caution: teratogenicity




Figure VIII-2-2. Mechanism of Action of Finasteride

## Chapter Summary

### Adrenal Steroids

- \* The nonendocrine uses in inflammatory disorders were discussed in the previous chapter.
- \* The glucocorticoids are used to treat Addison disease and adrenal insufficiency states, as a supplement in infantile respiratory distress syndrome, and in adrenal hyperplasia.

### Estrogens

- \* Synthetic estrogens are used to increase the oral bioavailability and half-life relative to that obtained with estradiol and to induce feedback inhibition of FSH and LH.
- \* The uses and adverse effects of estrogens are listed.
- \* The clinical uses of anastrozole (decreases estrogen synthesis), danazol (decreases ovarian steroid synthesis), clomiphene (decreases feedback inhibition), and the selective estrogen-receptor modulators tamoxifen and raloxifene are considered.

### Progestin

- \* Synthetic progestins are used to increase oral bioavailability and half-life relative to progesterone and to induce feedback inhibition of gonadotropins, especially LH.
- \* The progestin-like drugs, their use in contraception and in hormonal replacement therapy, and their adverse effects are considered.
- \* Mifepristone is an antagonist used with PG as an abortifacient.
- \* The pharmacology of oral contraceptives and their adverse effects, drug interactions, and benefits are pointed out.

### Androgens

- \* Clinically useful androgen analogs include methyltestosterone and 17-alkyl derivatives. Their clinical and illicit uses and side effects are presented.
- \* Clinically useful drug antagonists are flutamide (an androgen-receptor blocker used to treat prostate cancer), leuprolide (a GnRH analog used to treat prostate cancer), and finasteride (a 5- $\alpha$ -reductase inhibitor used to treat benign prostatic hyperplasia and male pattern baldness).




# Antithyroid Agents

**Table VIII-3-1. The Synthesis of Thyroid Hormone and Action and Effects of Antithyroid Agents**

Thyroid Hormone Synthesis	Effects of Antithyroid Agents
1. Active accumulation of iodide into gland	Basis for selective cell destruction by $^{131}\text{I}$
2. Oxidation of iodide to iodine by peroxidase	Inhibited by thioamides
3. Iodination of tyrosyl residues (organification) on thyroglobulin to form MIT and DIT	Inhibited by thioamides
4. Coupling of MIT and DIT to form $\text{T}_3$ and $\text{T}_4$	Inhibited by thioamides
5. Proteolytic release of $\text{T}_3$ and $\text{T}_4$ from thyroglobulin	Inhibited by high doses of iodide*
6. Conversion of $\text{T}_4$ to $\text{T}_3$ via 5' deiodinase in peripheral tissues	Inhibited by propranolol* and propylthiouracil*

*Definition of abbreviations:* MIT, monoiodotyrosine; DIT, diiodotyrosine;  $\text{T}_3$ , triiodothyronine;  $\text{T}_4$ , thyroxine.

\*Thyroid storm management may include the use of any or all of these agents.



**Figure VIII-3-1. Thyroid Hormone Synthesis**

- Thioamides: propylthiouracil and methimazole
  - Use: uncomplicated hyperthyroid conditions; slow in onset
  - High-dose propylthiouracil inhibits 5' deiodinase
  - Common maculopapular rash
  - Both drugs cross the placental barrier, but propylthiouracil is safer in pregnancy because it is extensively protein bound
- Iodide
  - Potassium iodide plus iodine (Lugol's solution) possible use in thyrotoxicosis: used pre operatively → ↓ gland size, fragility, and vascularity
  - No long-term use because thyroid gland "escapes" from effects after 10 to 14 days

### **Chapter Summary**

- \* The steps in thyroid hormone synthesis and the antithyroid agents' effects upon them are summarized in Table VIII-3-1. The clinical uses and their potential complications are presented in greater detail for the thioamides (propylthiouracil and methimazole) and iodine.

# Drugs Related to Hypothalamic and Pituitary Hormones

Table VIII-4-1. Drugs Related to Hypothalamic and Pituitary Hormones

Hormone	Pharmacologic Agent	Clinical Uses
GH	Somatrem or somatropin	Pituitary dwarfism, osteoporosis
Somatostatin	Ocreotide	Acromegaly, carcinoid and secretory-GI tumors
ACTH	Cosyntropin	Infantile spasms
GnRH	Leuprolide, nafarelin	Endometriosis, prostate carcinoma (repository form)
FSH and LH	Urofollitropin (FSH), placental HCG (LH), menotropins (FSH and LH)	Hypogonadal states
PIH (DA)	Bromocriptine	Hyperprolactinemia
Oxytocin	Oxytocin	Labor induction
Vasopressin	Desmopressin (V2 selective)	<ul style="list-style-type: none"> <li>• Neurogenic (pituitary) diabetes insipidus</li> <li>• Hemophilia A (↑ factor VIII from liver)</li> <li>• von Willebrand disease (↑ vW factor from endothelium)</li> <li>• Primary nocturnal enuresis</li> </ul>

*Definition of abbreviations:* ACTH, adrenocorticotropin hormone; DA, dopamine; FSH, follicle-stimulating hormone; GH, growth hormone; GnRH, gonadotropin-releasing hormone; LH, luteinizing hormone; PIH, prolactin-inhibiting hormone.

## Chapter Summary

- \* The clinical uses of drugs used to treat functions associated with hypothalamic or pituitary hormones are summarized in Table VIII-4-1.



# Drugs Used for Bone and Mineral Disorders

---

## BISPHOSPHONATES

- Mechanisms: stabilize hydroxyapatite bone structure and also induce osteoblasts to secrete inhibitors of osteoclasts → ↓ bone resorption → ↓ progression of osteoporosis
- Clinical uses:
  - Established use in Paget disease
  - Efficacy in postmenopausal osteoporosis depends on individual drug, but alendronate is effective and with HRT causes ↑ bone mineral density (BMD).
  - Alendronate is drug of choice for glucocorticoid-induced osteoporosis
- Side effects:
  - Etidronate and pamidronate → bone mineralization defects
  - Gastrointestinal distress, including esophageal ulcers (alendronate)

## TERIPARATIDE

- Mechanism: recombinant DNA PTH analog
- Clinical use: once daily to stimulate osteoblasts and new bone formation
- Continuous infusion would stimulate osteoclast activity
- Used for less than 2 years; may ↑ risk of osteosarcoma

### Chapter Summary

- \* The bisphosphonates decrease bone resorption and slow the progress of osteoporosis. Alendronate is effective for treatment of postmenopausal and steroid-induced osteoporosis. The principal potential side effects are gastrointestinal distress and esophageal ulcers.





# Endocrine Drug List and Practice Questions

Table VIII-6-1. Endocrine Drug List

Drugs Used in Diabetes	Steroid Hormones ( <i>Cont'd</i> )
Insulins, glargine	<i>Androgens</i> Methyltestosterone Oxandrolone Flutamide (antagonist) Finasteride (5- $\alpha$ -reductase inhibitor)
Sulfonylureas—chlorpropamide, tolbutamide, acetohexamide, glipizide, glyburide	
Metformin	
Acarbose	
Thiazolidinediones—pioglitazone, rosiglitazone	
GLP-1 drugs—exenatide, sitagliptin	<b>Antithyroid Drugs</b> Propylthiouracil Methimazole KI and I (Lugol's)
<b>Steroid Hormones</b>	
<i>Adrenosteroids</i>	<b>Hypothalamic/Pituitary Drugs</b> Somatropin Octreotide Leuprolide Oxytocin, vasopressin
Cortisol	
Triamcinolone	
Fludrocortisone	
Prednisone	
Dexamethasone	
<i>Estrogens</i>	<b>Drugs Used in Bone and Mineral Disorders</b> Alendronate Teriparatide
Ethinyl estradiol	
Mestranol	
Diethylstilbestrol	
Tamoxifen (SERM)	
Raloxifene (SERM)	
<i>Progestins</i>	
Medroxyprogesterone	
Norgestrel	
Norethindrone	
Desogestrel	
Mifepristone (antagonist)	

### Review Questions

1. A 70-year-old man is diagnosed with benign prostatic hyperplasia (BPH), and his physician is considering drug treatment of the condition. It was decided that the drug finasteride will be used. The effects of finasteride will result in a decrease in the synthesis of what substance?
  - A. Epinephrine
  - B. Norepinephrine
  - C. Dihydrotestosterone
  - D. Testosterone
  - E. GnRH
2. Which of the following statements is accurate regarding drug management of hyperthyroidism?
  - A. The actions of thyroid peroxidase are inhibited by  $I^{131}$
  - B. Propylthiouracil inhibits the conversion of thyroxine to triiodothyronine
  - C. Methimazole is unable to cross the placental barrier
  - D. Iodide salts can be used for long-term management
  - E. The iodination of tyrosyl residues to form MIT and DIT are inhibited by beta blockers
3. What drug is useful to distinguish neurogenic from nephrogenic diabetes insipidus?
  - A. Amiloride
  - B. Demeclocycline
  - C. Desmopressin
  - D. Hydrochlorothiazide
  - E. Lithium
4. The release of insulin from the pancreatic B cell would most likely be stimulated by which of the following?
  - A. Clonidine
  - B. Norepinephrine
  - C. Diazoxide
  - D. Glipizide
  - E. Hypoglycemia
5. When used at higher doses than commonly employed for other purposes, what drug can effectively inhibit steroidogenesis in a variety of tissues?
  - A. Flutamide
  - B. Misoprostol
  - C. Clomiphene
  - D. Tamoxifen
  - E. Ketoconazole

6. In a patient with type 2 diabetes, which drug mimics the action of incretins to augment glucose-dependent insulin secretion?
  - A. Acarbose
  - B. Glucagon
  - C. Exenatide
  - D. Metformin
  - E. Rosiglitazone
  
7. To supplement other oral type 2 diabetes medication, a patient is prescribed a drug to inhibit the intestinal absorption of carbohydrates. What would be an appropriate drug?
  - A. Metformin
  - B. Acarbose
  - C. Repaglinide
  - D. Insulin lispro
  - E. Pioglitazone
  
8. What is the drug of choice for management of adrenal glucocorticoid-induced osteoporosis?
  - A. Alendronate
  - B. Calcitonin
  - C. Estrogen
  - D. Ketoconazole
  - E. Vitamin D
  
9. Which drug has utility in inhibiting the severe secretory diarrhea of hormone-secreting tumors of the pancreas and GI tract, as well as in the treatment of acromegaly?
  - A. Octreotide
  - B. Leuprolide
  - C. Bromocriptine
  - D. Sertraline
  - E. Anastrozole

### Answers

- Answer: C.** The drug finasteride inhibits the enzyme 5- $\alpha$  reductase, the enzyme that convert testosterone to dihydrotestosterone (DHT). DHT is responsible for prostate enlargement in BPH. The other commonly used drugs in BPH are alpha-1 antagonists such as prazosin and tamsulosin.
- Answer: B.** Thioamides used at conventional doses in Graves disease are slow to act; they inhibit iodination and the coupling reactions in hormone synthesis and do not affect the release of stored thyroxine. At high doses, propylthiouracil may act more rapidly because of its inhibition of 5'-deiodinase, preventing the conversion of T<sub>4</sub> to T<sub>3</sub>. Thioamides are not teratogenic, and they do not decrease glandular size or vascularity; KI plus iodine (Lugol's solution) is used preoperatively to this end. Use of iodide in hyperthyroidism is only temporary because the thyroid gland "escapes" from its actions within a week or two.
- Answer: C.** Neurogenic diabetes insipidus is treated with desmopressin, a drug that is similar to vasopressin (ADH), yet is a selective activator of V<sub>2</sub> receptors in the kidney. Remember that V<sub>1</sub> receptors are present in smooth muscle, and their activation leads to vasoconstriction and bronchoconstriction. Nephrogenic diabetes insipidus (decreased response of vasopressin receptors) is treated with thiazides, except in the case of that induced by lithium, when amiloride is preferred (because thiazides increase blood levels of lithium). In cases where it is necessary to distinguish between neurogenic and nephrogenic diabetes insipidus, desmopressin is used. Desmopressin would alleviate the symptoms of neurogenic but have no effect in nephrogenic diabetes insipidus.
- Answer: D.** The release of insulin from the pancreas is stimulated by insulinogens (glucose), sulfonylurea hypoglycemics (glipizide), activators of beta-2 adrenoceptors (e.g., albuterol) and activators of muscarinic receptors (e.g., pilocarpine). Activation of alpha-2 receptors inhibits insulin release (clonidine and norepinephrine). Hypokalemia, and diazoxide, keep potassium channels on beta cells open resulting in decreased insulin release.
- Answer: E.** Ketoconazole is an antifungal drug that decreases the synthesis of various steroids including cortisol and testosterone by inhibiting cytochrome P450 enzymes. Flutamide is an androgen-receptor antagonist, and tamoxifen is a partial agonist (or mixed agonist-antagonist) at estrogen receptors. Misoprostol is a prostaglandin analog used in NSAID-induced ulcers. Clomiphene is used to induce ovulation.
- Answer: C.** Exenatide is a glucagon-like peptide-1 (GLP-1) analog. GLP-1 is an incretin released from the small intestine that enhances glucose-dependent insulin secretion. Metformin is "euglycemic," lowering elevated glucose levels to the normal range, and acarbose simply prevents postprandial hyperglycemia. Glucagon causes hyperglycemia, an effect that is sometimes employed in management of hypoglycemia. Rosiglitazone increases the sensitivity to insulin by increasing insulin receptor numbers.
- Answer: B.** Acarbose inhibits the enzyme  $\alpha$ -glucosidase in the brush borders of the small intestine. This decreases the formation of absorbable carbohydrate and thereby decreases glucose absorption. The net effect is that glucose levels after a meal don't rise as significantly and therefore the insulin demand is reduced.

8. **Answer: A.** Alendronate is currently the drug of choice to prevent osteoporosis in patients who must be maintained on steroids for their antiinflammatory and immunosuppressive effects. The drug also decreases bone resorption during menopause and is sometimes favored in patients who are at risk for neoplasias if treated with sex hormones. Care must be taken with alendronate to avoid esophageal ulceration. Estrogen hormone replacement therapy  $\pm$  vitamin D also has proven valuable for slowing bone resorption in menopause, and increases in bone mass have been reported for combinations of estrogens with alendronate.
  
9. **Answer: A.** Octreotide is a somatostatin analog that is effective for carcinoid and other secretory GI tumors. It has been used to varying degrees in other forms of secretory diarrhea such as chemotherapy-induced and the diarrhea of HIV and diabetes. Octreotide has proven to be of major importance in the management of acromegaly and can significantly decrease the levels of growth hormone.



**SECTION IX**

# **Anticancer Drugs**





# Anticancer Drugs

## PRINCIPLES AND DEFINITIONS

### Log-Kill hypothesis

Cytotoxic actions of anticancer drugs follow first-order kinetics: They kill a fixed percentage of tumor cells, not a fixed number → one rationale for drug combinations.

### Growth Fraction

Cytotoxic drugs are more effective against tumors that have a high growth fraction (large percentage actively dividing). Normal cells with high growth fraction (e.g., bone marrow) are also more sensitive to anticancer drugs.

### Cell-Cycle Specificity

- Drugs that act specifically on phases of the cell cycle are called cell-cycle specific (CCS) and are more effective in tumors with high-growth fraction (leukemias, lymphomas).
- Drugs that are cell-cycle nonspecific (many bind to and damage DNA) can be used in tumors with low-growth fraction, as well as tumors with high-growth fraction.




Figure IX-1-1. Cell-Cycle Specificity of Anticancer Drugs

## DRUGS

Table IX-1-1. Characteristics of Important Anticancer Drugs

Drug	Mechanism	Uses	Side Effects
Cyclophosphamide	Alkylating agent—attacks guanine N7—dysfunctional DNA	Non-Hodgkin, ovarian, breast cancer, neuroblastoma	BMS, hemorrhagic cystitis ( <b>mesna</b> , traps acrolein and is protective)
Cisplatin	Alkylating agent—cross-links DNA strands	Testicular, ovarian, bladder, lung cancer	Nephrotoxicity (use amifostine); neurotoxicity (deafness)
Procarbazine	Alkylating agent	Hodgkin	BMS, leukemogenic
Doxorubicin	Intercalator, forms free radicals, inhibits topoisomerase	Hodgkin, breast, endometrial, lung, and ovarian cancers	BMS—delayed CHF ( <b>dexrazoxane</b> is an iron-chelating agent preventing the formation of free radicals; it is not a free radical “trapper”)
Methotrexate (CCS)	Antimetabolite—inhibits DHF reductase (S phase)	Leukemias, lymphomas, breast cancer; rheumatoid arthritis, psoriasis	BMS, <b>leucovorin</b> (folinic acid) rescue
5-Fluorouracil (CCS)	Pyrimidine antimetabolite (S phase) bioactivated to inhibit thymidylate synthetase	Breast, ovarian, head, and neck cancer—topical for basal cell cancer and keratoses	BMS
6-Mercaptopurine (CCS)	Purine antimetabolite (S phase) bioactivated by HGP transferase	Acute lymphocytic leukemia; immunosuppression	BMS
Bleomycin (CCS)	Complexes with Fe and O <sub>2</sub> → DNA strand scission (G <sub>2</sub> phase)	Hodgkin, testicular, head, neck, skin cancer	Pneumonitis, pulmonary fibrosis
Vinblastine (CCS) Vincristine	↓ Microtubular polymerization—spindle poisons (M phase)	Vinblastine—Hodgkin, testicular cancer, Kaposi Vincristine—Hodgkin, leukemias, Wilms	BMS Neurotoxicity

Definition of abbreviations: BMS, bone marrow suppression; CCS, cell-cycle specific; CHF, congestive heart failure; GI, gastrointestinal.

## Clinical Correlate

## Thymineless Death of Cells

Flucytosine (FC) and 5-fluorouracil (5-FU) are bioactivated to 5-fluorodeoxyuridine (5-FdUMP), which inhibits thymidylate synthetase → “thymineless death” of fungal cells (FC) or neoplastic cells (5-FU).

## TOXICITY OF ANTICANCER DRUGS

Table IX-1-2. Other Dose-Limiting or Distinctive Toxicities

Toxicity	Drug(s)
Renal	Cisplatin,* methotrexate
Pulmonary	Bleomycin,* busulfan, procarbazine
Cardiac	Doxorubicin, daunorubicin
Neurologic	Vincristine,* cisplatin
Immunosuppressive	Cyclophosphamide, methotrexate
Other	Cyclophosphamide (hemorrhagic cystitis); procarbazine (leukemia); asparaginase* (pancreatitis)

\*Less BMS—"marrow sparing"

- Rapidly proliferating cells, such as the bone marrow, gastrointestinal tract mucosa, hair follicles, and gonads are the most sensitive to cytotoxic drugs.
- Most often *bone marrow suppression* (BMS) is dose limiting.
- Anticancer drug dosage is usually carefully titrated to avoid excessive neutropenia (granulocytes  $<500/\text{mm}^3$ ) and thrombocytopenia (platelets  $<20,000/\text{mm}^3$ ).
- Colony-stimulating factors, erythropoietin, and thrombopoietin can be supportive →↓ infections and need for antibiotics (see table below).

Table IX-1-3. Clinical Uses of Cytokines

Cytokine	Clinical Uses
Aldesleukin (IL-2)	↑ Lymphocyte differentiation and ↑ NKs—use in renal cell cancer and metastatic melanoma
Interleukin-11	↑ Platelet formation—used in thrombocytopenia
Filgrastim (G-CSF)	↑ Granulocytes—used for marrow recovery
Sargramostim (GM-CSF)	↑ Granulocytes and macrophages—used for marrow recovery
Erythropoietin	Anemias, especially associated with renal failure
Thrombopoietin	Thrombocytopenia

## RESISTANCE TO ANTICANCER DRUGS

Another rationale for combination drug regimens in cancer is to prevent or delay the emergence of resistance.

**Table IX-1-4. Modes of Resistance to Anticancer Drugs**

Mechanism	Drugs or Drug Groups
Change in sensitivity (or ↑ level) or ↓ binding affinity of target enzymes or receptors	Etoposide, methotrexate, vinca alkaloids, estrogen and androgen receptors
Decreased drug accumulation via ↑ expression of glycoprotein transporters, or ↓ permeability	Methotrexate, alkylating agents, dactinomycin
Formation of drug-inactivating enzymes	Purine and pyrimidine antimetabolites
Production of reactive chemicals that “trap” the anticancer drug	Alkylators, bleomycin, cisplatin, doxorubicin
Increased nucleic acid repair mechanisms	Alkylating agents, cisplatin
Reduced activation of prodrugs	Purine and pyrimidine antimetabolites

### Chapter Summary

- \* The “log-kill” hypothesis states that cytotoxic anticancer agents kill a certain percentage, not a fixed number, of cells.
- \* Cytotoxic drugs are most effective against rapidly dividing cells.
- \* Drugs that act on proliferating cells are cell-cycle specific and are usually also cycle-phase specific. Figure IX-1-1 illustrates the cell cycle and the drugs acting in each cycle phase.
- \* Drugs that act on nonproliferating cells are dose-dependent and cell-cycle-independent.
- \* Rationales for combination drug usage are that each drug will independently kill a fixed percentage and that one drug will still kill a cancer cell that has developed resistance to a different drug in the cocktail.
- \* Rapidly proliferating normal cells are more sensitive to cytotoxic drugs. Bone marrow suppression often determines the upper limit of tolerable chemotherapy. Table IX-1-1 lists mechanisms of action, selected clinical uses, and side effects of major anticancer drugs. Table IX-1-2 shows the dose-limiting and distinctive toxicities of anticancer drugs.
- \* Table IX-1-4 summarizes the modes of resistance developed by cancers toward specific anticancer drugs.

# Anticancer Drug Practice Questions

---

## Review Questions

1. Which of the following chemotherapeutic drugs inhibits the polymerization of microtubules but is not associated with causing bone marrow suppression?
  - A. Cyclophosphamide
  - B. Cisplatin
  - C. 5-Fluorouracil
  - D. Vinblastine
  - E. Vincristine
2. A patient with non-Hodgkin lymphoma is to be started on the CHOP regimen, which consists of cyclophosphamide, doxorubicin, vincristine, and prednisone. Which one of the following agents is most likely to be protective against the toxicity of doxorubicin?
  - A. Amifostine
  - B. Dexrazoxane
  - C. Leucovorin
  - D. Mesna
  - E. Vitamin C
3. A drug used in a chemotherapy regimen works by complexing with iron and oxygen to promote DNA strand breaks. While on this drug the patient must be monitored closely due to pulmonary side effects. In what phase of the cell cycle does this drug work?
  - A. G<sub>1</sub>
  - B. S
  - C. G<sub>2</sub>
  - D. M
  - E. This drug is not cell-cycle dependent.
4. Resistance to which anticancer drug, used in leukemias, lymphomas, and breast cancer, is associated with increased production of dihydrofolate reductase?
  - A. Doxorubicin
  - B. Vinblastine
  - C. 6-MP
  - D. Cytarabine
  - E. Methotrexate

5. A patient undergoing cancer chemotherapy has an increase in urinary frequency with much discomfort. No specific findings are apparent on physical examination. Laboratory results include hematuria and mild leukopenia, but no bacteria or crystalluria. If the symptoms experienced by the patient are drug related, what is the most likely cause?
- A. Cyclophosphamide
  - B. 5-FU
  - C. Methotrexate
  - D. Prednisone
  - E. Tamoxifen

### Answers

1. **Answer: E.** Only two of the drugs listed do not cause bone marrow suppression: cisplatin and vincristine. Only two of the drugs listed inhibit microtubule polymerization: vinblastine and vincristine. The drug that fits both categories is vincristine. Patients on vincristine should be monitored for neurotoxicity, especially peripheral neuropathies.
2. **Answer: B.** Dexrazoxane is an iron-chelating agent that prevents the formation of free radicals and reduces the cardiotoxicity of anthracyclines such as doxorubicin. Amifostine is protective of nephrotoxicity caused by cisplatin. Folinic acid (leucovorin) reduces the toxicity of methotrexate because it provides an active form of folate to normal (nonneoplastic) cells, resulting in "leucovorin rescue." Mesna, which inactivates acrolein, is available for protection against hemorrhagic cystitis in patients treated with cyclophosphamide.
3. **Answer: A.** It helps to know which anticancer drugs are cell-cycle specific and which have characteristic toxicities. Bleomycin forms a complex with iron and oxygen and promotes DNA strand breaks. Its major side effects are pulmonary toxicities including pneumonitis and fibrosis. It acts acting mainly in the G<sub>2</sub> phase of the cell-cycle.
4. **Answer: E.** Methotrexate is a widely-used chemotherapy drug that is also commonly used in moderate to severe rheumatoid arthritis. It inhibits the enzyme dihydrofolate reductase (DHFR) thereby reducing the synthesis of tetrahydrofolate and thus inhibiting DNA synthesis. Resistance occurs when cancer cells upregulate DHFR or alter the binding of methotrexate to DHFR.
5. **Answer: A.** These symptoms are those of a mild case of hemorrhagic cystitis. Bladder irritation with hematuria is a fairly common complaint of patients treated with cyclophosphamide. It appears to be due to acrolein, a product formed when cyclophosphamide is bioactivated by liver P450 to form cytotoxic metabolites. Mesna is the antidote used to detoxify acrolein and protect against hemorrhagic cystitis.





**SECTION X**

# **Immunopharmacology**



# Immunopharmacology

---

## CYCLOSPORINE AND TACROLIMUS

- Mechanism of action:
  - Bind to cyclophilin (cyclosporin) or FK-binding protein (tacrolimus) → ↓ calcineurin (cytoplasmic phosphatase) → ↓ activation of T-cell transcription factors → ↓ IL-2, IL-3, and interferon- $\gamma$
- Uses:
  - Cyclosporine is DOC for organ or tissue transplantation (+/- mycophenolate, +/- steroids, +/- cytotoxic drugs)
  - Tacrolimus used alternatively to cyclosporine in renal and liver transplants
  - Mycophenolate, an inhibitor of de novo synthesis of purines, has adjunctive immunosuppressant actions, permitting dose reductions of cyclosporine to limit toxicity.
- Side effects: nephrotoxicity (both), gingival overgrowth (cyclosporine)

## AZATHIOPRINE

Immunosuppressant converted to 6-mercaptopurine—same properties as 6-MP.

## ANTI-D IMMUNOGLOBULIN

- Human IgG antibodies to red cell D antigen (rhesus antigen)
- Uses: Administer to Rh-negative mother within 72 hours of Rh-positive delivery to prevent hemolytic disease of newborn in subsequent pregnancy

## MONOCLONAL ANTIBODIES

Table X-1-1. Clinical Uses of Monoclonal Antibodies

Mab	Clinical Uses
Abciximab	Antiplatelet—antagonist of IIb/IIIa receptors
Infliximab	Rheumatoid arthritis and Crohn disease—binds TNF
Trastuzumab	Breast cancer—antagonist to ERB-B2 (Her 2/neu)
Dacliximab	Kidney transplants—blocks IL-2 receptors
Muromonab	Kidney transplant—blocks allograft rejection
Palivizumab	Respiratory syncytial virus—blocks RSV protein
Rituximab	Non-Hodgkin lymphoma—binds to surface protein

## CYTOKINES (RECOMBINANT FORMS)

Table X-1-2. Clinical Uses of Interferons

Interferon- $\alpha$	Hepatitis B and C, leukemias, melanoma
Interferon- $\beta$	Multiple sclerosis
Interferon- $\gamma$	Chronic granulomatous disease $\rightarrow \uparrow$ TNF

### Chapter Summary

- \* The mechanism of action, uses, and toxicities associated with cyclosporine are presented. Azathioprine converts to 6-mercaptopurine, making it a useful immunosuppressant.
- \* Anti-D immunoglobulin is given to Rh-negative mothers shortly after parturition to prevent hemolytic disease in future births.
- \* Table X-1-1 summarizes the clinical uses of monoclonal antibodies. Table X-1-2 summarizes the clinical uses of recombinant cytokines.

# Immunopharmacology Practice Questions

---

## Review Questions

1. A patient is treated with an immunosuppressant drug following a liver transplant. The drug is known to bind to cyclophilin and inhibit the actions of calcineurin. For what drug toxicity should this patient be monitored?
  - A. Pulmonary fibrosis
  - B. Hypotension
  - C. Hypoglycemia
  - D. Nephrotoxicity
  - E. CHF
2. Which one of the following agents has utility in the management of acute coronary syndromes such as unstable angina?
  - A. Abciximab
  - B. Interferon- $\alpha$
  - C. Aldesleukin
  - D. Filgrastim
  - E. Trastuzumab

## Answers

1. **Answer: D.** This patient is being treated with cyclosporin, a drug that binds to cyclophilin and inhibits calcineurin. As a result, the transcription of various T-cells factors such as IL-2, IL-3, and Interferon- $\gamma$  are inhibited. Cyclosporin is associated with nephrotoxicity, gingival hyperplasia, hyperglycemia, hypertension, and hirsutism.
2. **Answer: A.** Abciximab is an antibody-based drug that targets glycoprotein IIb/IIIa receptors. Binding of the drug to these receptors results in decreased platelet aggregation by preventing the cross-linking reaction. It is useful in acute coronary syndromes such as unstable angina and post-angioplasty.



**SECTION XI**

# **Toxicology**





# Toxicology

## COMMON TOXIC SYNDROMES

Table XI-1-1. Signs, Symptoms, and Interventions or Antidotes for Common Toxic Syndromes

Compound(s)	Signs and Symptoms	Interventions and Antidotes
AChE inhibitors	Miosis, salivation, sweats, GI cramps, diarrhea, muscle twitches → seizures, coma, respiration failure	Respiratory support; atropine + pralidoxime (for irreversible AChE inhibitors)
Atropine and muscarinic blockers	↑ HR, ↑ BP, hyperthermia (hot, dry skin), delirium, hallucinations, mydriasis	Control cardiovascular symptoms and hyperthermia + physostigmine (crosses blood–brain barrier)
Carbon monoxide (>10% carboxyHb)	Nausea and vomiting, dyspnea with hyperventilation, mydriasis, vertigo; cardiovascular signs prominent, ↓ BP, syncope, ↑ HR, arrhythmias	Hyperbaric O <sub>2</sub> and decontamination (humidified 100% O <sub>2</sub> okay in mild overdose)
CNS stimulants	Anxiety/agitation, hyperthermia (warm, sweaty skin), mydriasis, ↑ HR, ↑ BP, psychosis, seizures	Control cardiovascular symptoms, hyperthermia, and seizures— +/- BZs or antipsychotics
Opioid analgesics	Lethargy, sedation, ↓ HR, ↓ BP, hypoventilation, miosis, coma, respiration failure	Ventilatory support; naloxone at frequent intervals
Salicylates (ASA)*	Confusion, lethargy, hyperventilation, hyperthermia, dehydration, hypokalemia, acidosis, seizures, coma	Correct acidosis and electrolytes—urinary alkalization, possible hemodialysis
Sedative-hypnotics and ethanol	Disinhibition (initial), lethargy, ataxia, nystagmus, stupor, coma, hypothermia, respiratory failure	Ventilatory support—flumazenil if BZs implicated
SSRIs	Agitation, confusion, hallucination, muscle rigidity, hyperthermia, ↑ HR, ↑ BP, seizures	Control hyperthermia and seizures—possible use of cyproheptadine, antipsychotics, and BZs
Tricyclic antidepressants	Mydriasis, hyperthermia (hot, dry skin), 3 Cs (convulsions, coma, and cardiotoxicity) → arrhythmias	Control seizures and hyperthermia, correct acidosis and possible arrhythmias

\*More details in antiinflammatory section

## HEAVY METAL POISONING

Signs and symptoms distinctive but usually result from inhibition of –SH groups on enzymes and regulatory proteins.

Table XI-1-2. Signs, Symptoms, and Interventions or Antidotes for Heavy Metal Poisoning

Metals and Source	Signs and Symptoms	Interventions and Antidotes
Arsenic (wood preservatives, pesticides, ant poisons)	<i>Acute:</i> gastroenteritis, hypotension, metabolic acidosis, garlic breath, “rice water” stools, torsades, seizures  <i>Chronic:</i> pallor, skin pigmentation (raindrop pattern), alopecia, stocking glove neuropathy, myelosuppression	Activated charcoal, <b>dimercaprol</b>  <b>Penicillamine</b> or succimer
Iron (medicinal for anemias and prenatal supplements)	<i>Acute</i> (mainly children): severe GI distress → necrotizing gastroenteritis with hematemesis and bloody diarrhea, dyspnea, shock, coma	Gastric aspiration + carbonate lavage, <b>deferoxamine IV</b>
Lead (tap water, leaded paint chips, herbal remedies, gas sniffing, glazed kitchenware, etc.)	<i>Acute:</i> nausea and vomiting, GI distress and pain, malaise, tremor, tinnitus, paresthesias, encephalopathy (red or black feces)  <i>Chronic:</i> multisystem effects— <i>anemia</i> (↓ heme synthesis), neuropathy (wrist drop), nephropathy (proteinuria, failure), hepatitis, mental retardation (from pica), ↓ fertility and ↑ stillbirths	Decontamination—gastric lavage + <b>dimercaprol</b> (severe) or EDTA or succimer (penicillamine if unable to use dimercaprol or succimer)  Children: <b>succimer PO</b>
Mercury (elemental in instruments); salts used in amalgams, batteries, dyes, electroplating, fireworks, photography	<i>Acute:</i> vapor inhalation—chest pain, dyspnea, pneumonitis  <i>Acute:</i> inorganic salt ingestion—hemorrhagic gastroenteritis, acute tubular necrosis, shock  <i>Chronic:</i> organic Hg—CNS effects, ataxia, paresthesias, auditory and visual loss, loosening of teeth	<b>Succimer PO</b> or <b>dimercaprol (IM)</b>  Activated charcoal for oral ingestion, then support with succimer PO or dimercaprol ( <i>not IV</i> ) → causes redistribution of Hg to the CNS → ↑ neurotoxicity

## ANTIDOTES

Table XI-1-3. Summary of Antidotes

Antidote	Type of Poisoning
Acetylcysteine	Acetaminophen
Atropine + pralidoxime (for irreversible AChE inhibitors)	AChE inhibitors—physostigmine, neostigmine, and pyridostigmine; organophosphates, including insecticides, such as malathion and parathion
Deferoxamine	Iron and iron salts
Digoxin immune F(ab)	Digoxin
Dimercaprol (BAL)	Arsenic, gold, mercury, lead; oral succimer for milder lead and mercury toxicity
EDTA	Backup in lead poisoning, then for rarer toxicities (Cd, Cr, Co, Mn, Zn)
Esmolol	Theophylline, beta agonists
Ethanol, fomepizole	Methanol or ethylene glycol
Flumazenil	Benzodiazepines, zolpidem, zaleplon
Naloxone	Opioid analgesics
Oxygen	Carbon monoxide
Penicillamine	Copper (e.g., Wilson's disease), iron, lead, mercury
Physostigmine	Anticholinergics: atropine, antihistamine, antiparkinsonian— <i>not</i> tricyclics
Protamine	Heparins
Vitamin K	Warfarin and coumarin anticoagulants
Activated charcoal	Nonspecific: all oral poisonings except Fe, CN, Li, solvents, mineral acids, or corrosives

## NATURAL MEDICINALS

“Natural” medicinals are available without prescription and are considered to be nutritional supplements rather than drugs. Herbal (botanic) products are marketed without FDA review of safety and efficacy, and there are no requirements governing the purity or the chemical identities of constituents. Evidence supporting the clinical effectiveness of herbal products is commonly incomplete.

Table XI-1-4. Characteristics of Selected Herbs

Name	Medicinal Use(s)	Possible Mechanism(s)	Side Effects
Echinacea	↓ Cold symptoms	↑ ILs and TNF	GI distress, dizziness, headache
Garlic	Hyperlipidemias, cancer (evidence is weak)	Inhibits HMG-CoA reductase and ACE	Allergies, hypotension, antiplatelet actions; use caution when used with anticoagulants
Ginkgo	Intermittent claudication; Alzheimer disease (evidence is weak)	Antioxidant, free radical scavenger, ↑ NO	Anxiety, GI distress, insomnia, antiplatelet actions; <b>use caution when used with anticoagulants</b>
Ginseng	Possible ↑ in mental and physical performance (evidence is weak)	Unknown	Insomnia, nervousness, hypertension, mastalgia, vaginal bleeding
Saw palmetto	Symptomatic treatment of BPH	5 $\alpha$ -reductase inhibitor and androgen receptor antagonist	GI pain, decreased libido, headache, hypertension
St. John's wort	Depressive disorder (variable evidence for clinical efficacy)	May enhance brain 5HT functions	Major drug interactions: <b>serotonin syndrome</b> with SSRIs; induces P450, leading to ↓ effects of multiple drugs

Table XI-1-5. Purified Nutritional Supplements

Name	Pharmacology	Side Effects
Dehydroepiandrosterone (DHEA)	Androgen precursor advocated for treatment of AIDS ( $\uparrow$ CD4 in females), Alzheimer disease and "aging," diabetes, hypercholesterolemia, and SLE ( $\downarrow$ in symptoms and "flare-ups" in females)	<i>Females:</i> androgenization and concern regarding CV disease and breast cancer <i>Males:</i> feminization in young and concern in elderly regarding BPH and cancer
Melatonin	Serotonin derivative used for "jet-lag" and sleep disorders	Drowsiness, sedation, headache. Contraindicated in pregnancy, in women trying to conceive ( $\downarrow$ LH), and in nursing mothers ( $\downarrow$ prolactin)

### Chapter Summary

- \* Table XI-1-1 lists the common toxic syndromes with their signs and symptoms and potential modes of intervention and/or antidotes.
- \* Table XI-1-2 lists the common heavy metal poisons with their most common sources, signs, and symptoms and potential modes of intervention and/or antidotes.
- \* Table XI-1-3 lists antidotes and the type of poisoning against which they act.
- \* Table XI-1-4 lists the characteristics of selected herbals, and Table XI-1-5 lists the relevant purified nutritional supplements.



# Toxicology Practice Questions

---

## Review Questions

1. Chronic ingestion of lead-based paint chips will result in which of the following?
  - A. Garlic breath
  - B. Changes in skin pigmentation
  - C. Accumulation of  $\delta$ -aminolevulinate and inhibition of heme synthesis
  - D. Auditory and visual loss
  - E. Interstitial pneumonitis and neurological effects
2. A 3-year-old child was brought to the ER following the ingestion of several pills. The child is suffering from severe gastrointestinal discomfort and has thrown up twice, each time producing a bloody vomitus. Questioning of the mother reveals the child got into the mother's old prenatal vitamins. What antidote should be given?
  - A. Dimercaprol
  - B. Deferoxamine
  - C. EDTA
  - D. Penicillamine
  - E. Succimer

## Answers

1. **Answer: C.** Chronic poisoning with lead will result in a multitude of effects including inhibition of heme synthesis and accumulation of  $\delta$ -aminolevulinate in the plasma. Arsenic poisoning is associated with garlic breath and changes in skin pigmentation. Organic mercury causes auditory and visual loss and loosening of the teeth, while inhaled mercury vapor can cause interstitial pneumonitis and neurological effects.
2. **Answer: B.** The child is suffering from iron poisoning. Deferoxamine chelates iron and is the antidote in iron poisoning. The other choices are all metal chelators with utility in other types of heavy metal poisoning. Dimercaprol is useful for a variety of metals including lead, arsenic, and mercury. EDTA is a back-up in lead poisoning. Penicillamine is useful in copper poisoning, and succimer is preferred for lead poisoning in kids.





# Index

Note: Page numbers followed by f indicate figures.

## A

### Abciximab

- characteristics/uses, 264
- clinical uses, 304
- platelet activation, 263f, 264

### Absorption, in pharmacokinetics, 3f, 6–7

- antacid impact, 89, 223
- bioavailability, 7, 7f, 10
- first-pass effect, 7, 7f
- nature of, 6
- plasma level curves, 6, 6f

### Abuse. *See* Drugs of abuse

### Acarbose

- characteristics/uses, 273
- modes of action, 273f

### Acebutolol, characteristics/uses, 59

### Acetaldehyde toxicity, 133f

### Acetalsalicylic acid (ASA)

- characteristics/uses, 229–230
- common toxic syndromes, 309

### Acetaminophen

- antidote for, 311
- characteristics/uses, 231
- cytochrome P450 isozymes, 10
- metabolism of, 11
- for migraine headaches, 226

### Acetazolamide

- as carbonic anhydrase inhibitor, 108
- modes of action/effects, 112

### Acetohexamide, 272

### Acetylation, 11

### Acetylcholine (ACh), 40. *See also* Cholinergic pharmacology

- muscarinic agonist, 47
- receptors for, 23, 24

### Acetylcholinesterase (AChE), 25, 45f, 46

- inhibition. *See* AChE inhibitors

### Acetylcysteine, antidote for, 311

### AChE inhibitor poisoning, 48–49, 309, 311

### AChE inhibitors, 45f, 48–49

- drug lists, 48, 69
- mechanism of action, 62

### Acidification of urine, 5

### Action potential, 83–85

- fast-response fibers, 83f, 83–84
- phases, 83–84
- slow-response fibers, 84f, 84–85

### Activated charcoal, as antidote, 311

### Active metabolites, 9

### Active transport, 5

### Acute mountain sickness, drugs for treating, 108

### Acyclovir

- characteristics/uses, 196
- mechanism of action, 195, 196

### Adenosine, characteristics/uses, 91

### ADHD (attention deficit hyperactivity disorder), drugs for treating, 156, 163

### Adrenergic pharmacology

- adrenergic receptor activators, 54, 55–58, 69
- adrenergic receptor antagonists, 58–60, 69
- catecholamine synthesis/action/degradation, 53f, 53–55

### Adrenergic receptor activators, 53f, 54–58

- $\alpha_1$  agonists, 54, 55, 55f, 69
- $\alpha_2$  agonists, 54, 55, 69
- $\beta$  agonists, 54, 56, 69
- direct-acting, 54, 55–58, 69
- drug list, 69
- indirect-acting, 58, 69
- mixed, 54, 56–58, 69

### Adrenergic receptor antagonists

- $\alpha_1$  antagonists, 58–59, 69
- $\alpha_2$  antagonists, 69
- $\beta$  receptor antagonists, 59–60, 69, 94
- drug list, 69
- indirect-acting, 58, 69

### Adrenoceptor sensitivity, 54

### Adrenoceptors, 24

### Adrenosteroids

- characteristics/uses, 275
- drug list, 287

### Affinity, defined, 19

### Ag sulfadiazine, characteristics/uses, 184

### Agonists

- defined, 19
- in dose-response (D-R) curves, 19–22
- full, 20, 20f
- partial, 20, 20f, 21, 21f

### Albuterol, as beta adrenergic agonist, 56

- Alcohols
  - antidotes for poisoning, 311
  - cytochrome P450 isozymes, 10
  - as drugs of abuse, 159–160
  - metabolism and pharmacologic actions, 10, 11, 133, 133f
  - metabolism of, 133f
  - and pregnancy, 133
  - withdrawal signs, 131
- Aldesleukin (IL-2), clinical uses, 297
- Aldosterone
  - as adrenal steroid antagonist, 275
  - in hormonal feedback loop, 41–42
- Aliskiren, 96, 96f
- Alkalosis, drugs for treating, 107, 108
- Alkylating agents
  - cell-cycle specificity, 295f
  - resistance to androgen receptors, 298
- Allergic reactions
  - drugs for treating, 219–220
  - to sulfonamide-containing drugs, 109
- Allopurinol
  - characteristics/uses, 235–236
  - drug interaction with 6-Mercaptopurine, 236f
  - mechanism of action, 235f
- Alpha adrenergic receptors
  - activators, 54, 55, 55f, 69, 93
  - blockers. *See* Alpha blockers
- Alpha blockers, 58–59, 69
  - as antihypertensives, 93, 94, 97
- Alprazolam
  - characteristics/uses, 130
  - grapefruit juice and, 10
- Alprostadil, 228
- Alteplase (tPA)
  - actions, 261f
  - characteristics/uses, 261
- Alzheimer disease, drugs for treating
  - acetylcholinesterase inhibitors, 48
  - characteristics of illness, 48
- Amantadine
  - as antiparkinsonian drug, 151
  - characteristics/uses, 199
  - as M blockers, 50
  - mechanism of action, 195, 199
- Amebiasis, metronidazole for, 203
- Amides, 141
- Amikacin, 181, 186
  - mechanism of resistance, 176
- Amiloride
  - as diuretic, 111
  - modes of action/effects, 112
- Amine hypothesis, 155
- Aminocaproic acid, 261
- Aminoglycosides
  - characteristics/uses, 181–182
  - daily dosage, 181
  - drug list, 205
  - mechanism of action, 175
  - mechanism of resistance, 176
  - mechanism summary, 181
  - pregnancy risk, 183
- Aminophylline VI, in treating asthma, 240
- Amiodarone, characteristics/uses, 90–91
- Amitriptyline, 155
  - for migraine headaches, 226
- Amoxicillin, 177
  - for treating *H. pylori* ulcers, 186
- Amphetamines
  - as drugs of abuse, 159
  - as indirect-acting adrenergic agonist, 58
- Amphotericin B (Amp B), characteristics/uses, 191–192
- Ampicillin, 177
- Amyl nitrite, for cyanide poisoning, 95
- Anakinra, characteristics/uses, 233
- Analgesics
  - for migraine headache, 226
  - nonsteroidal antiinflammatory drugs (NSAIDs), 229–231
  - opioid. *See* Opioid analgesics
- Anastrozole, 276
  - mechanism of action, 276f
- Androgens
  - characteristics/uses, 278
  - drug list, 287
  - resistance to androgen receptors, 298
- Androstenedione, mechanism of action of anastrozole, 276f
- Anesthetics, 139–143
  - compartmentalization in body, 139f
  - drug list, 163
  - general, 139–140
  - inhaled, 139–140, 163
  - intravenous, 140, 163
  - local, 141, 163
  - skeletal muscle relaxants, 142–143, 163
- Angiotensin-converting enzyme inhibitors (ACEIs)
  - for heart failure, 100
  - for hypertension, 25, 96f, 96–97
- Angiotensin I, 96f
- Angiotensin II, 96f
  - in hormonal feedback loop, 41–42
  - receptors for, 24
- Angiotensin-receptor blockers (ARBs)
  - for heart failure, 100
  - for hypertension, 96f, 96–97
- Anion overdose, drugs for treating, 109
- ANS. *See* Autonomic nervous system (ANS)
- Antacids
  - absorption and, 89, 223
  - for peptic ulcer disease, 223
- Antagonists
  - antagonism and potentiation, 21f, 21–22
  - chemical, 22
  - competitive, 21
  - defined, 19
  - noncompetitive, 21, 22
  - partial agonists as, 21
  - physiologic, 22

- Anti-D immunoglobulin, characteristics/uses, 303
- Anti-HIV agents, 197–199
  - drug list, 205
  - HIV prophylaxis, 199
  - protease inhibitors (PIs), 198–199
  - reverse transcriptase inhibitors (RTIs), 197–198
  - side effects, 198
  - zidovudine, 198
- Anti-motion sickness drugs
  - antihistamines, 220
  - M blockers, 50
- Antianginal drugs, 103–106
  - beta-blockers, 59, 104
  - calcium channel blockers, 104
  - carvedilol, 104
  - drug list, 117
  - nitrates, 103f, 103–104
  - rationale for use, 103
  - smooth muscle contraction/relaxation, 105f
  - for stable and vasospastic angina, 103
- Antiarrhythmic drugs, 89–92
  - beta-blockers, 59, 90
  - calcium-channel blockers, 91
  - and cardiac action potential, 83–84, 85
  - drug list, 117
  - K<sup>+</sup> channel blockers, 60, 90–91
  - Na<sup>+</sup> channel blockers, 89–90
  - unclassified, 91
- Antiasthmatic drugs
  - characteristics/uses, 239–241
  - drug actions on bronchiolar smooth muscle, 239f
  - drug list, 244
  - M blockers, 50
- Antibacterial agents, 175–189
  - antitubercular drugs, 186–187
  - drug list, 205
  - inhibitors of bacterial protein synthesis, 175, 180–184
  - inhibitors of cell-wall synthesis, 175, 176–180
  - inhibitors of nucleic acid synthesis, 175, 184–186
  - metronidazole, 186
  - pregnancy risk, 183
  - principles of antimicrobial chemotherapy, 175–176
- Anticancer drugs, 295–298
  - cell-cycle specificity, 295, 295f
  - characteristics/uses, 296
  - clinical uses of cytokines, 297
  - principles and definitions, 295–296
  - resistance to, 298
  - “thymineless” death of cells, 296
  - toxicity, 297
- Anticholinergics
  - antidote for, 311
  - as drugs of abuse, 160
- Anticoagulants, 257–260
  - drug list, 265
  - heparins, 258
  - warfarin, 258–259
- Anticonvulsants, 135–137
  - characteristics/uses, 135–137
  - drug list, 163
  - seizure states/drugs of choice, 135
- Antidepressants, 155–157
  - characteristics/uses, 155–156
  - drug list, 163
- Antidiarrheal drugs, M blockers, 50
- Antidotes
  - for heavy metal poisoning, 310
  - summary table, 311
- Antiemetics
  - characteristics/uses, 223
  - drug list, 243
  - emetic pathways, 223f
- Antifibrinolytics, 261
- Antifolates, drug list, 205
- Antifungal agents, 191–193
  - azoles, 192
  - drug list, 205
  - mechanism of action, 191f
  - other, 193
  - polyenes, 191–192
- Antiherpetics, 196–197
  - acyclovir, 196
  - drug list, 205
  - foscarnet, 197
  - ganciclovir, 197
- Antihistamines, 219–220
  - drug list, 243
  - as M blockers, 50
  - physostigmine as antidote for, 311
  - properties of major, 220
- Antihyperlipidemics, 113–115
  - bile acid sequestrants, 114, 117
  - drug list, 117
  - ezetimibe, 115, 117
  - gemfibrozil, 114–115, 117
  - HMG-CoA reductase inhibitors, 113–114
  - nicotinic acid, 114, 117
  - orlistat, 115, 117
  - summary of uses, 115
- Antihypertensive drugs, 93–98
  - adrenoceptor agonists, 54
  - angiotensin-converting inhibitors (ACEIs), 25, 96f, 96–97
  - angiotensin-receptor blockers (ARBs), 96f, 96–97
  - ANS feedback loop, 40–41, 42
  - beta-blockers, 59
  - calcium-channel blockers, 95
  - in comorbid conditions, 97
  - direct-acting vasodilators, 94–95
  - diuretics, 96, 109. *See also* Diuretics
  - drug list, 117
  - drug strategy, 93
  - drugs altering sympathetic activity, 93–94
  - endocrine feedback loop, 41–42
  - for pulmonary hypertension, 97, 229
  - thiazides, 110–111

- Antileukotrienes, in treating asthma, 240
  - Antimalarial drugs, 203–204
  - Antimetabolites, 184
  - Antimicrobial agents
    - antibacterial agents, 175–189, 205
    - antifungal agents, 191–193, 205
    - antiprotozoal agents, 203–204, 205
    - antiviral agents, 195–201, 205
    - mechanism of action, 175
    - mechanism of resistance, 176
  - Antimycobacterials, drug list, 205
  - Antiparkinsonian drugs, 149–151
    - characteristics/uses, 150–151
    - CNS targets, 150f
    - dopaminergic neural pathways and, 149
    - drug list, 163
    - M blockers, 50
    - physostigmine as antidote for, 311
  - Antiplatelet drugs
    - characteristics/uses, 263–264
    - drug list, 265
    - platelet activation, 263f
    - platelet aggregation, 263
  - Antiprotozoal agents, 203–204
    - antimalarial, 203–204
    - drug list, 205
    - helminthic infections, 204
    - infections/drugs of choice, 203
  - Antipsychotic drugs, 152–153
    - characteristic properties, 153
    - drug list, 163
    - M blockers, 50
    - parenteral formulations, 153
  - Antisecretor drugs, M blockers, 50
  - Antispasmodic drugs, M blockers, 50
  - Antithyroid agents
    - characteristics/uses, 281–282
    - drug list, 287
    - effects, 281
    - thyroid hormone synthesis, 281, 281f
  - Antitubercular drugs, 186–187
    - characteristics/uses, 186
    - summary table, 187
  - Antitumor antibiotics, 295f
  - Antiviral agents, 195–201
    - anti-HIV, 197–199, 205
    - antiherpetics, 196–197, 205
    - drug list, 205
    - fusion inhibitors, 199
    - mechanism of action, 195
    - other, 199–200
    - sites of drug actions, 195f
  - Apparent volume of distribution ( $V_d$ ), 8, 17
  - Aprepitant, 223
  - Argatroban, 258
  - Aripiprazole, characteristic properties, 153
  - Aromatase, 276f
  - Arsenic poisoning, 310, 311
  - Arsenicals, as antiprotozoal agent, 203
  - Arteriodilators, for heart failure, 100
  - Aspartate protease, 25, 198
  - Aspirin
    - characteristics/uses, 229–230
    - platelet activation, 263, 263f
  - Asthma, drugs for treating, 50, 239–241, 244
  - Astreonam, mechanism of action, 175
  - Atenolol, characteristics, 59
  - Atomoxetine, 156
  - Atorvastatin, grapefruit juice and, 10
  - Atovaquone-proguanil, 204
  - ATP (adenosine triphosphate), 23
  - Atracurium, 142
  - Atropine
    - as antidote for AChE inhibitor poisoning, 311
    - characteristics/uses, 49–50
    - common toxic syndromes, 309
    - physostigmine as antidote for, 311
  - Automaticity, cardiac action potentials, 85
  - Autonomic feedback loop, 40–41, 41f
  - Autonomic nervous system (ANS), 39–44
    - anatomy, 39–40, 40f
    - blood pressure control mechanisms, 40–42
    - pupillary size and accommodation mechanisms, 43f
    - regulation of heart rate, 87
  - Autonomic pharmacology
    - adrenergic receptor activators/antagonists, 53–60, 69
    - autonomic nervous system (ANS), 39–44
    - cholinergic receptor activators/antagonists, 45–52, 69
    - drug list, 69
    - glaucoma, 61–67
  - Azathioprine, characteristics/uses, 303
  - Azidothymidine, 198
  - Azithromycin, mechanism of resistance, 176
  - Azlocillin, 177
  - Azoles, 192
  - AZT, 195, 198
  - Aztronam
    - characteristics/uses, 179
    - mechanism of action, 179
- ## B
- Baclofen, as centrally acting skeletal muscle relaxant, 143
  - Bacterial protein synthesis inhibitors, 180–184
    - drug list, 205
    - mechanism of action, 175
    - mechanism summary, 181
  - Baldness, drugs for treating, 95
  - Barbiturates
    - characteristics/uses, 129–131
    - CNS effects based on dosage, 129f
    - cytochrome P450 isozymes, 10
    - as drugs of abuse, 159–160
    - withdrawal signs, 131
  - Barriers to distribution, 8
  - Batrachotoxin, 141
  - Benzathine penicillin G, 177
  - Benzocaine, as local anesthetic, 141

- Benzodiazepines (BZs)  
 as centrally acting skeletal muscle relaxants, 143  
 characteristics/uses, 129–131, 140  
 CNS effects based on dosage, 129f  
 as drugs of abuse, 159–160  
 flumazenil as antidote, 311  
 formation of nordiazepam, 9  
 withdrawal signs, 131
- Benztropine  
 as antiparkinsonian drug, 151  
 characteristics/uses, 50
- Beta-blockers, 59–60  
 as antianginal, 104  
 as antiarrhythmics, 59, 90  
 as antihypertensives, 94, 97  
 characteristics/uses, 59–60, 62, 90, 94  
 chronic use, 59  
 in glaucoma treatment, 62
- Beta-receptor agonists, 239  
 activators, 54, 56, 69  
 blockers versus. *See* Beta-blockers  
 esmolol as antidote, 311  
 for heart failure, 100
- Beta-receptor antagonists. *See* Beta-blockers
- Betamethasone, 237
- Bethanechol, muscarinic agonist, 47
- Bile acid sequestrants, 114, 117
- Bilirubin, protein-binding capacity, 8
- Bioavailability (f), 7, 7f, 10
- Biotransformation, 9–11  
 nature of, 9  
 process of, 9f  
 types, 10–11
- Biotransformation, in pharmacokinetics, 3f, 9–11  
 classification, 10–11  
 process, 9f
- Bipolar disorders, lithium for treating, 156, 163
- Bismuth subsalicylate, 222
- Bleomycin  
 cell-cycle specificity, 295f, 296  
 characteristics/uses, 296  
 resistance to androgen receptors, 298  
 toxicity, 297
- Blood-brain barrier (BBB) to distribution, 8, 9
- Blood disorders, drugs for treating, 257–267  
 actions of blood drugs, 257f  
 anticoagulants, 257–260, 265  
 antiplatelet drugs, 263–264, 265  
 drug list, 265  
 thrombolytics, 261–262, 265
- Blood pressure control mechanisms, 40–42  
 autonomic feedback loop, 40–41, 41f  
 blood pressure/heart rate tracings, 42f  
 hormonal feedback loop, 41–42, 42f
- Blood pressure tracings, 42f
- Bone and mineral disorders, drugs for treating  
 characteristics/uses, 287  
 drug list, 287
- Bosentan, for pulmonary hypertension, 97
- Botulinum toxin, 45, 45f
- Bromocriptine, as antiparkinsonian drug, 151
- Bupivacaine, as local anesthetic, 141
- Buprenorphine, characteristics/uses, 146
- Bupropion, 156
- Buspiroline, characteristics/uses, 131, 225
- Busulfan, toxicity, 297
- Butobarbital, for migraine headaches, 226
- Butorphanol, for migraine headaches, 226
- C**
- Ca<sup>+</sup> channels, slow-response fibers, 84–85
- Caffeine, for migraine headaches, 226
- Calcium-channel blockers (CCBs)  
 as antianginal, 104, 105f  
 as antiarrhythmic, 91  
 as antihypertensive, 95, 97  
 cardiac/vascular selectivity, 95f
- Calculations, in pharmacokinetics, 17
- cAMP (cyclic-AMP), 23–24, 24f  
 ANS regulation of heart rate, 87
- Cancer, drugs for treating, 295–298
- Capreomycin, 186
- Captopril, 96
- Carbamazepine  
 characteristics/uses, 136  
 cytochrome P450 isozymes, 10  
 seizure states/drugs of choice, 135
- Carbon monoxide poisoning, 309, 311
- Carbonic anhydrase, 25
- Carbonic anhydrase inhibitors, 108f, 108–109
- Carboprost, 228
- Cardiac output (CO), 40, 41f
- Cardiac pharmacology  
 action potential, 83–85  
 ANS regulation of heart rate, 87  
 antianginal drugs, 103–106, 117  
 antiarrhythmic drugs, 59, 60, 89–92, 117  
 antihyperlipidemics, 113–115, 117  
 antihypertensive drugs, 40–42, 54, 59, 93–98, 117  
 cytochrome P450 isozymes, 10  
 diuretics, 96, 101, 107–112, 117  
 heart failure drugs, 99–102, 117  
 Na<sup>+</sup> channels, 86
- Carmustine, 295f
- Carvedilol, 60  
 as antianginal, 104  
 for heart failure, 101
- Catechol-O-methyltransferase (COMT), 54
- Catecholamine synthesis/action/degradation, 53f, 53–55
- Catecholamines, receptors for, 24
- Cefaclor, 178
- Cefazolin, 178
- Cefdinir, 178
- Cefepime, 178
- Cefixime, 178
- Cefoperazone, 133, 178
- Cefotaxime, 178

- Cefotetan, 133, 178
- Ceftriaxone, 178
- Cefuroxime, 178
- Celecoxib, 109
  - characteristics/uses, 231
- Cell-cycle specificity (CCS)
  - of anticancer drugs, 295f, 296
  - defined, 295
- Cell-wall synthesis inhibitors, 176–180
  - drug list, 205
  - mechanism of action, 175
- Cephalexin, 178
- Cephalosporins
  - characteristics/uses, 178
  - drug list, 205
  - mechanism of action, 175, 176f, 178
  - mechanism of resistance, 176, 178
  - organisms not covered by (LAME), 178
- Cetirizine, properties, 220
- cGMP (cyclic GMP) signaling, 25
- Chloramphenicol
  - characteristics/uses, 182
  - cytochrome P450 isozymes, 10
  - mechanism of action, 175
  - mechanism of resistance, 176
  - mechanism summary, 181
  - metabolism of, 11
- Chloroquine, 204
- Chlorpheniramine, properties, 220
- Chlorpromazine, characteristic properties, 153
- Chlorpropamide, 133, 272
- Cholestyramine
  - as bile acid sequestrant, 114, 117
  - summary of antihyperlipidemic uses, 115
- Choline, 45
- Cholinergic neuroeffector junctions, 45f, 45–47
- Cholinergic pharmacology, 45–52
  - cholinergic neuroeffector junctions, 45f, 45–47
  - muscarinic receptor activators, 46, 47–49, 69
  - muscarinic receptor antagonists, 49–50, 69
  - nicotinic receptor antagonists, 50–51, 69
- Cholinergic receptor activators
  - AChE inhibitors, 45f, 48–49, 69
  - direct, 45–47, 69
  - drug list, 69
- Cholinergic receptor antagonists
  - drug list, 69
  - ganglionic blockers, 50–51, 69
  - M blockers, 49–50, 69
- Cholinomimetics
  - irreversibly acting, 49f
  - mechanism of action, 62
- Ciguatoxin, 141
- Cimetidine
  - characteristics/uses, 221–222
  - cytochrome P450 isozymes, 10, 222
- Ciprofloxacin, characteristics/uses, 185–186
- Cisplatin
  - cell-cycle specificity, 295f
  - characteristics/uses, 296
  - resistance to androgen receptors, 298
  - toxicity, 297
- Clarithromycin
  - mechanism of resistance, 176
  - for treating *H. pylori* ulcers, 186
- Class IA antiarrhythmic drugs, 89, 117
- Class IB antiarrhythmic drugs, 89–90, 117
- Class IC antiarrhythmic drugs, 90, 117
- Class II antiarrhythmic drugs, 90, 117
- Class III antiarrhythmic drugs, 90–91, 117
- Class IV antiarrhythmic drugs, 91, 117
- Clavulanic acid, 177
- Clearance (Cl), 14
- Clindamycin
  - characteristics/uses, 183
  - mechanism of resistance, 176
  - mechanism summary, 181
- Clomiphene, 276
- Clomipramine, 155
- Clonidine
  - as  $\alpha_2$  adrenergic agonist, 55, 93
  - characteristics/uses, 93
- Clopidogrel
  - characteristics/uses, 264
  - platelet activation, 263f
- Clotrimazole, 192
- Clozapine, characteristic properties, 153
- CNS pharmacology
  - ADHD, 156, 163
  - alcohols, 129, 133, 163
  - anesthetics, 139–143, 163
  - anticonvulsants, 135–137, 163
  - antidepressants, 155–157, 163
  - antiparkinsonian drugs, 149–151, 163
  - antipsychotics, 152–153, 163
  - bipolar disorder, 156, 163
  - CNS stimulants, common toxic syndromes, 309
  - cytochrome P450 isozymes, 10
  - drug list, 163
  - drugs of abuse, 159–161
  - opioid analgesics, 145–147, 163, 309
  - redistribution, f, 9
  - sedative-hypnotics, 129–133, 163, 309
- Cocaine
  - in blocking norepinephrine uptake, 141
  - as drug of abuse, 159
  - as indirect-acting adrenergic agonist, 58
  - as local anesthetic, 141
- Codeine, characteristics/uses, 146
- Colchicine, characteristics/uses, 235
- Colestipol
  - as bile acid sequestrant, 114, 117
  - summary of antihyperlipidemic uses, 115
- Comorbid conditions, indications for use of antihypertensive drugs, 97

- Competitive antagonists, 21  
 Concentration gradient, 4  
 Conductance, cardiac action potentials, 85  
 Congestive heart failure, diuretics for treating, 111  
 COPD drugs, M blockers, 50  
 Copper poisoning, 311  
 Cortisol, synthetic derivatives of, 237  
 Coumarin. *See* Warfarin (coumarin)  
 COX 1, 228  
 COX 2, 228  
 CREB, 23  
 Cromolyn, in treating asthma, 240  
 Cumulative D-R curves. *See* Quantal (cumulative) dose-response (D-R) curves  
 Cyanide poisoning, 95  
 Cyanomethemoglobin, 95  
 Cyclooxygenases, 25  
 Cyclophosphamide  
   characteristics/uses, 296  
   toxicity, 297  
 Cycloserine, 186  
 Cyclosporine  
   characteristics/uses, 303  
   grapefruit juice and, 10  
 Cyproheptadine, 225  
 Cytarabine, cell-cycle specificity, 295f  
 Cytochrome P450 isozymes, 10, 222  
 Cytokines  
   clinical uses, 297, 304  
   receptors for, 25
- D**  
 D<sub>1</sub> receptors, activation, 54  
 D-Tubocurarine prototype, 142  
 DA. *See* Dopamine (DA)  
 Daciximab, clinical uses, 304  
 Dactinomycin, resistance to androgen receptors, 298  
 Danaparoid, 258  
 Dantrolene, 143  
 Daunorubicin, toxicity, 297  
 Deferoxamine, for heavy metal poisoning, 310, 311  
 Dehydroepiandrosterone (DHEA), characteristics/uses, 313  
 Dehydrogenases, 11  
 Demeclocycline, 182  
 DES (diethylstilbestrol), 275  
 Desogestrel, 277  
 Dexamethasone, 237  
 Dextromethorphan, 147  
 DHEA (dehydroepiandrosterone), characteristics/uses, 313  
 Diabetes, drugs for treating, 271–274  
   acarbose, 273, 273f  
   agents affecting glucagon-like peptide-1 (GLP-1), 274, 287  
   drug list, 287  
   insulins, 271–272, 287  
   metformin, 273  
   modes of action, 273f  
   sulfonylureas, 272, 287  
   thiazolidinediones, 273, 287
- Diabetic ketoacidosis, 271  
 Diacylglycerol (DAG), 24, 24f  
 Diastolic pressure, 42f  
 Diazepam  
   characteristics/uses, 130  
   formation of nordiazepam, 9  
   seizure states/drugs of choice, 135  
 Diazoxide, as direct-acting vasodilator, 95  
 Didanosine (DDI)  
   mechanism of action, 195  
   side effects, 198  
 Diethylstilbestrol (DES), 275  
 Diffusion, 5  
 Digoxin  
   antidote for, 311  
   characteristics/uses, 100  
   for heart failure, 100, 100f  
 Digoxin immune, 311  
 Dihydrofolate reductase, 25  
 Dihydropyridines, as calcium-channel blocker, 95  
 Dihydroxyphenylalanine (DOPA), 53  
 Diloxanide, 203  
 Diltiazem, as calcium-channel blocker, 91, 95  
 Dimercaprol, for heavy metal poisoning, 310, 311  
 Dinoprostone, 228  
 Diphenhydramine  
   as antiemetic, 223  
   as antiparkinsonian drug, 151  
   properties, 220  
 Disease-modifying antirheumatic drugs (DMARDs), 233–234, 244  
 Distribution, in pharmacokinetics, 3f, 8–9  
   apparent volume of distribution ( $V_d$ ), 8, 17  
   conditions affecting, 8  
   special barriers to, 8  
 Diuretics, 107–112  
   actions, 107f  
   alkalosis, 107  
   carbonic anhydrase inhibitors, 108f, 108–109  
   drug list, 117  
   for heart failure, 100, 101  
   for hypertension, 96  
   hypokalemia, 107  
   K<sup>+</sup>-sparing agents, 111–112, 117  
   loop, 109f, 109–110  
   mode of action/effects, 112  
   osmotic, 108  
   thiazides, 109, 110–111, 117  
 DMARDs (disease-modifying antirheumatic drugs), 233–234, 244  
 DNA/RNA polymerase, 25  
 Dobutamine  
   as beta adrenergic agonist, 56  
   for heart failure, 101  
 Donepezil, characteristics/uses, 48  
 Dopamine (DA), 40, 53  
   blood-brain barrier to distribution, 8  
   drugs of abuse and, 159  
   for heart failure, 101  
   as precursor to norepinephrine, 149  
   receptors for, 24, 149, 150f

- reuptake inhibitors, 25
- use in shock, 54
- Dopamine hypothesis, 152
- Dopamine receptors, 24, 149, 150f
- Dopaminergic neural pathways, 149
- Dorzolamide, as carbonic anhydrase inhibitor, 108
- Dosage
  - CNS effects of sedative-hypnotic-anxiolytic drugs, 129f
  - epinephrine (E), dosage effects on heart rate and blood pressure, 57f
  - infusion rate equations, 17
  - multiple-dose equations, 17
  - single-dose equations, 17
- Dose-response (D-R) curves
  - dosage calculations, 17
  - in drug development and testing, 26
  - graded (quantitative), 19–22
  - quantal (cumulative), 22–23
  - signaling mechanisms, 23–25
- Doxazosin, as alpha blocker, 59, 94
- Doxorubicin
  - characteristics/uses, 296
  - resistance to androgen receptors, 298
  - toxicity, 297
- Doxycycline, 182, 204
- Dronabinol, 223
- Drugs of abuse, 159–161
  - opioid analgesics, 146–147, 159
  - properties, 159–160
- E**
- Echinacea, characteristics/uses, 312
- Echothiophate, 46
- Echothiophate, mechanism of action, 62
- Ecstasy (MDMA), as drug of abuse, 160
- Edema, diuretics for treating, 109
- Edrophonium, characteristics/uses, 48
- EDTA, as antidote, 311
- Efavirenz, mechanism of action, 195
- Effective refractory period (ERP), 85
- Efficacy
  - defined, 19
  - of full and partial agonists, 20f
- Eicosanoids, 227–229
  - drugs acting on, 227f
  - leukotrienes (LTs), 228
  - nonsteroidal antiinflammatory drugs, 226, 229–231, 234, 243
  - platelet stability, 229
  - prostaglandins (PGs), 227, 228–229
  - thromboxanes (TXAs), 229
- Elimination, in pharmacokinetics, 3f, 12–14
  - elimination kinetics, 13
  - first-order elimination rate, 12–13, 13f
  - graphic analysis, 13, 13f
  - renal, 14
  - zero-order elimination rate, 12, 12f, 13
- Endocrine pharmacology
  - antithyroid agents, 281–282, 287
  - bone and mineral disorders, 287
  - diabetes, 271–274, 287
  - drug list, 287
  - hypothalamic hormones in, 283, 287
  - pituitary hormones in, 283, 287
  - steroid hormones, 275–279, 287
- Endothelin (ET), for pulmonary hypertension, 97
- Enfuvirtide, mechanism of action, 195, 199
- Entacapone, as antiparkinsonian drug, 151
- Enzymes
  - See also* Angiotensin-converting enzyme inhibitors (ACEIs)
- Enzymes, receptors that function as, 25
- Ephedrine, as indirect-acting adrenergic agonist, 58
- Epidermal growth factor (EGF), 25
- Epinephrine (E), 40
  - clinical uses, 58
  - dosage effects on heart rate and blood pressure, 57f
  - as mixed-acting adrenergic agonist, 57–58
  - nicotinic receptor activation, 47
- Eplerenone
  - as aldosterone receptor blocker, 111
  - modes of action/effects, 112
- Epoprostenol, 229
  - for pulmonary hypertension, 97
- Eptifibatid, characteristics/uses, 264
- Ergonovine, characteristics/uses, 226
- Ergosterol, 191
- Ergotamine, characteristics/uses, 226
- Erythromycin, mechanism of resistance, 176
- Erythropoietin, clinical uses, 297
- Esmolol
  - as antidote, 311
  - as beta-blocker, 90
  - characteristics/uses, 59, 90
- Esters, 141
- Estone, 276f
- Estradiol, 275
  - mechanism of action of anastrozole, 276f
- Estrogens
  - characteristics/uses, 275–277
  - drug list, 287
  - in oral contraceptives, 277–278
  - resistance to estrogen receptors, 298
- Etanercept, characteristics/uses, 233
- Ethacrynic acid
  - as loop diuretic, 109
  - modes of action/effects, 112
- Ethambutol, 186, 187
- Ethanol. *See* Alcohols
- Ethinyl estradiol, 275, 277
- Ethosuximide
  - characteristics/uses, 137
  - seizure states/drugs of choice, 135
- Etoposide
  - cell-cycle specificity, 295f
  - resistance to, 298
- Exenatide, 274
- Ezetimibe
  - as antihyperlipidemic, 115, 117
  - summary of antihyperlipidemic uses, 115



**F**

- Famciclovir, 196
- Felbamate, characteristics/uses, 137
- Fenoldopam, 54
- Fentanyl
  - characteristics/uses, 140
  - as drug of abuse, 160
- Fexofenadine, properties, 220
- Fibrinogen, platelet activation, 263f
- Fibrinolytics. *See* Thrombolytics
- Filgrastim (G-CSF), clinical uses, 297
- Finasteride, 278
- First-order elimination rates, 12–13, 13f
- First-pass effect, 7, 7f
- 5-Alpha reductase inhibitor, 278
- 5-Fluorouracil
  - cell-cycle specificity, 296
  - characteristics/uses, 296
- 5HT<sub>1</sub>, 225
- 5HT<sub>2</sub>, 225
- 5HT<sub>3</sub>, 223, 225
- 5HT receptors
  - drug actions on, 225
  - drugs of abuse and, 159
- Flecainide, as Na<sup>+</sup> channel blocker, 90
- Fluconazole, characteristics/uses, 192
- Flucytosine, characteristics/uses, 193
- Flumazenil
  - as antidote, 311
  - in reversing CNS depression, 130, 131, 160
- Fluoroquinolones, 186
  - drug list, 205
  - mechanism of action, 175
  - mechanism of resistance, 176
  - pregnancy risk, 183
- Fluoxetine, 156
- Fluphenazine, characteristic properties, 153
- Flutamide, 278
- Folic acid synthesis inhibitors
  - characteristics/uses, 184–185
  - mechanism of action, 175
- Fomepizole, 311
  - for alcohol overdose, 133
- Food and Drug Administration (FDA)
  - drug classification, 26
  - drug development and testing, 26
  - pregnancy risk, 26
- Foscarnet
  - characteristics/uses, 197
  - mechanism of action, 195, 197
- Fosphenytoin, seizure states/drugs of choice, 135
- Full agonists, 20, 20f
- Furanocoumarins, 10
- Furosemide
  - as loop diuretic, 109
  - modes of action/effects, 112
- Fusion inhibitors, 199

**G**

- GABA
  - coupled to ion channel, 23
  - drugs of abuse and, 159–160
  - reuptake inhibitors, 25
  - site of action of drugs on GABA complex, 129f
- Gabapentin, characteristics/uses, 137
- Ganciclovir
  - characteristics/uses, 197
  - mechanism of action, 195, 197
- Ganglia, location of ANS, 39
- Ganglion blocking agents, 50–51
  - characteristics, 51
  - drug list, 69
  - reflex control of heart rate, 52f
- Garlic, characteristics/uses, 312
- Gastrointestinal dysfunction, drugs for treating, 186, 221–224, 243
- Gemfibrozil
  - as antihyperlipidemic, 114–115, 117
  - summary of antihyperlipidemic uses, 115
- General anesthetics, 139–140
- Gentamicin, 181
  - mechanism of resistance, 176
- G<sub>i</sub> proteins, 24
- Giardiasis, metronidazole for, 203
- Ginkgo, characteristics/uses, 312
- Ginseng, characteristics/uses, 312
- Glargine, 271
- Glaucoma, drugs for treating, 61–67, 228
  - acetylcholinesterase inhibitors, 48, 62
  - anatomy of eye, 61f
  - beta-blockers, 60, 62
  - diuretics, 108
  - mechanism of action, 62
  - types of glaucoma, 61–62
- Glipizide, 272
- Glomerular filtration rate (GFR), 14
- Glucagon
  - agents affecting glucagon-like peptide-1 (GLP-1), 274
  - receptors for, 24, 59
- Glucagon-like peptide-1 (GLP-1)
  - characteristics/uses, 274
  - modes of action, 274f
- Glucocorticoids
  - characteristics/uses, 233, 237–238
  - cytochrome P450 isozymes, 10
  - drug list, 244
  - endocrine uses of, 275
  - leukotrienes and, 228
  - minimizing steroid toxicity, 237
  - synthetic derivatives of cortisol, 237f
  - in treating asthma, 240
- Glucuronidation, 11
- Glutathione (GSH) conjugation, 11
- Glyburide, 272
- Gold poisoning, 311
- Gold salts, characteristics/uses, 233
- Gonadal steroids, intracellular receptors, 23

**Gout**

- drugs for treating, 235, 244
- prophylaxis, 235–236

**G<sub>q</sub> proteins, 24****Graded (quantitative) dose-response (D-R) curves, 19–22**

- antagonism, 21f, 21–22
- duality of partial agonists, 21, 21f
- full and partial agonists, 20, 20f
- parallel and nonparallel, 19f, 19–20
- potentiation, 21f, 21–22

**Grapefruit juice, in metabolism of drugs, 10****Griseofulvin, 133**

- characteristics/uses, 193

**Growth factors, 25****Growth fraction, 295****G<sub>s</sub> proteins, 24****GTP-binding proteins (G-proteins), 23–24****Guanethidine, characteristics/uses, 94****Guanyl cyclase, 25****H****H<sub>1</sub> agonists, 219–220****H<sub>2</sub> antagonists, 221–222****Half-life, 12–14**

- defined, 12
- effect of loading dose, 16
- equation, 14, 17

**Hallucinogens, as drugs of abuse, 160****Haloperidol**

- characteristic properties, 153
- cytochrome P450 isozymes, 10

**Halothane, characteristics, 139****Heart failure**

- drugs for treating, 99–102, 117
  - diuretics, 100, 101, 107–112, 117
  - inotropes, 100–101, 117
  - other drugs, 101, 117
- process of, 99f

**Heart rate**

- ANS regulation of, 87
- heart rate tracings, 42f
- reflex control of, 52f

**Heavy metal poisoning**

- antidotes, 310, 311
- signs/symptoms/interventions, 310

**Helminthic infections, drugs for treating, 204****Hemicholinium, 45, 45f****Heparin**

- characteristics/uses, 258
- comparison with warfarin, 258
- protamine as antidote, 311

**Hepatic encephalopathy, lactulose in, 4****Heroin, as drug of abuse, 160****Hexamethonium, as ganglion blocking agent, 50–51****Hirsutism, drugs for treating, 111****His-Purkinje system, 83f, 83–84****Histamine**

- characteristics/uses, 219

**drug list, 243**

- receptors for, 24

**HIV. See Anti-HIV agents****HMG-CoA reductase inhibitors, 113–114, 117****Hormonal feedback loop, 41–42, 42f****Human interferons, 200****Hydralazine**

- as direct-acting vasodilator, 94–95
- metabolism of, 11

**Hydrochlorothiazide, 110, 117**

- modes of action/effects, 112

**Hydrolysis, 11****Hydroxychloroquine**

- characteristics/uses, 233
- for rheumatoid arthritis, 234

**Hydroxyurea, cell-cycle specificity, 295f****Hypercalcemia, drugs for treating, 109****Hypersensitivity reactions, 177****Hyperthermia, malignant, 143****Hypoglycemia, drugs for treating, 272, 287****Hypothalamic disorders, drugs for treating**

- characteristics/uses, 287
- drug list, 287

**I****Iamotrigine, characteristics/uses, 137****Ibuprofen, 230****Imipenem/meropenem**

- characteristics/uses, 179
- mechanism of action, 175, 179

**Imipramine, 155****Immunopharmacology, 303–305****Inamrinone, for heart failure, 100f, 101****Indapamide, 110, 117**

- modes of action/effects, 112

**Indinavir**

- mechanism of action, 195
- side effects, 199

**Indomethacin, 230****Inflammatory and related disorders, drugs for treating, 219–254**

- antiemetics, 223, 243
- asthma, 50, 239–241, 244
- DMARDs, 233–234, 244
- drug list, 243–244
- eicosanoid pharmacology, 227–229
- gastrointestinal dysfunction, 221–224, 243
- glucocorticoids, 237–238, 244
- gout, 235–236, 244
- histamine and antihistamines, 219–220, 243
- migraine headaches, 226
- NSAIDs, 229–231, 243
- rheumatoid arthritis, 233–234, 244
- serotonergic systems, 225–226, 243

**Infliximab**

- characteristics/uses, 233
- clinical uses, 304

**Infusion rate, 15**

- effect on plasma level, 15f
- equation, 17

- Inhalants, as drugs of abuse, 160
- Inhaled anesthetics, 139–140, 163  
 MAC (minimal alveolar anesthetic concentration), 139  
 properties, 139f  
 rates of onset/recovery, 140
- Inositol triphosphate (IP), 24, 24f
- Inotropes, 100–101, 117
- Insecticides, 46, 48
- Insulins  
 characteristics/uses, 271–272  
 forms, 271
- Interferon- $\alpha$ , clinical uses, 304
- Interferon- $\beta$ , clinical uses, 304
- Interferon- $\gamma$ , clinical uses, 304
- Interferons, human, 200
- Interleukin-II, clinical uses, 297
- Intracellular receptors, 23
- Intravenous anesthetics, 140, 163
- Inulin clearance, 14
- Ion channels, membrane receptors coupled to, 23
- Ionization, 4–5  
 renal clearance of drugs, 4f, 5, 5f
- Ipratropium, characteristics/uses, 50, 240
- Iron poisoning, 310, 311
- Isoniazid (INH), 11, 186  
 prophylaxis, 187
- Isoproterenol, as beta adrenergic agonist, 56
- Isosorbide, as antianginal, 104
- Itraconazole, characteristics/uses, 192
- J**
- Janus kinases (JAKs), 25
- K**
- K<sup>+</sup> channel blockers, 60, 90–91
- K<sup>+</sup> channels, direct-acting vasodilators and, 95
- K<sup>+</sup> current, fast-response fibers, 83, 84
- K<sup>+</sup>-sparing agents, 111–112, 117
- Kanamycin, 186
- Ketamine  
 characteristics/uses, 140  
 as drug of abuse, 160
- Ketoconazole, 275  
 characteristics/uses, 192  
 cytochrome P450 isozymes, 10
- Ketorolac, 230
- L**
- L-aromatic amino acid decarboxylase (DOPA decarboxylase), 53
- L-DOPA. *See* Levodopa (L-DOPA)
- Labetalol, 60
- Lactic acid, 4
- Lactulose, 4
- Lamivudine  
 mechanism of action, 195  
 side effects, 198
- Latanoprost, 228
- Lead poisoning, 310, 311
- Leflunomide, characteristics/uses, 233
- Leishmaniasis, stibogluconate for, 203
- Leukotrienes (LTs), characteristics/uses, 228
- Leuprolide, 278
- Levodopa (L-DOPA)  
 as antiparkinsonian drug, 150, 150f  
 blood-brain barrier to distribution, 8  
 inhibitors of metabolism, 151f
- Levofloxacin, characteristics/uses, 185–186
- Lidocaine  
 characteristics/uses, 90  
 first-pass effect, 7  
 as local anesthetic, 141  
 as Na<sup>+</sup> channel blocker, 90
- Linezolid  
 characteristics/uses, 183, 184  
 mechanism of action, 175  
 mechanism summary, 181
- Lipid-soluble drugs, redistribution of, 9, 9f
- Lispro, 271
- Lithium, characteristics/uses, 156
- Loading doses (LD), 16, 16f  
 equation, 17
- Local anesthetics, 141, 163  
 mode of action, 141f  
 nonmicrosomal metabolism, 11
- Log-kill hypothesis, 295
- Lomustine, 295f
- Long QT syndrome, drugs for treating, 91
- Loop diuretics, 109f, 109–110
- Loperamide, 147
- Loratadine, properties, 220
- Lorazepam  
 characteristics/uses, 130  
 seizure states/drugs of choice, 135
- Losartan, 96, 96f
- Lovastatin, 113–114
- M**
- M blockers, 49–50  
 atropine, 49–50, 309, 311  
 characteristics/uses, 50  
 common toxic syndromes, 309  
 drug lists, 59, 69  
 other M blockers, 50  
 in treating asthma, 240, 244
- Macrolides  
 characteristics/uses, 183  
 community-acquired pneumonia, 183  
 cytochrome P450 isozymes, 10  
 drug list, 205  
 mechanism of action, 175  
 mechanism of resistance, 176  
 mechanism summary, 181
- Magnesium, as antiarrhythmic, 91
- Maintenance dose (MD), 17
- Malaoxon, 48

- Malaria  
antiprotozoal agents for, 203–204  
chloroquine-sensitive, 204
- Malathion, 46, 48
- Malignant hyperthermia, 143
- MAO inhibitors. *See also* Monoamine oxidase (MAO)
- MAO inhibitors, characteristics/uses, 155
- Maraviroc, mechanism of action, 195, 199
- Marijuana, as drug of abuse, 160
- MDEA, as drug of abuse, 160
- MDMA (ecstasy), as drug of abuse, 160
- Mebendazole, 204
- Mecamylamine, as ganglion blocking agent, 50–51
- Meclizine  
as antiemetic, 223  
properties, 220
- Medroxyprogesterone, 277
- Mefloquine, 204
- Melatonin, characteristics/uses, 313
- Membrane receptors, coupled to ion channels, 23
- Meperidine  
characteristics/uses, 146  
as M blockers, 50
- Mepivacaine, as local anesthetic, 141
- Mercury poisoning, 310, 311
- Mestranol, 275, 277
- Metabolism, in pharmacokinetics, 3f, 9–11, 133f
- Metformin  
characteristics/uses, 273  
modes of action, 273f
- Methacholine, muscarinic agonist, 47
- Methadone  
characteristics/uses, 146  
as drug of abuse, 160
- Methimazole, placental barrier to distribution, 8
- Methotrexate  
cell-cycle specificity, 295f, 296  
characteristics/uses, 233, 296  
resistance to, 298  
toxicity, 297
- Methoxamine, as  $\alpha_1$  adrenergic agonist, 55
- Methyldopa  
as  $\alpha_2$  adrenergic agonist, 55, 93  
characteristics/uses, 93
- Methylnaltrexone, characteristics/uses, 146
- Methylphenidate, 156
- Metoclopramide, 223
- Metolazone, modes of action/effects, 112
- Metoprolol  
characteristics, 59  
for heart failure, 101
- Metronidazole, 133  
as antiprotozoal agent, 203  
characteristics/uses, 186
- Metyrapone, 275
- Mexiletine, as  $\text{Na}^+$  channel blocker, 90
- Miconazole, 192
- Microsomal metabolism, 10
- Midazolam  
characteristics/uses, 130, 140  
grapefruit juice and, 10
- Mifepristone, 228, 275
- Migraine headaches, drugs for treating, 60, 226
- Milrinone, for heart failure, 100f, 101
- Minocycline, 182
- Minoxidil, as direct-acting vasodilator, 95
- Mirtazapine, 156  
as alpha blocker, 59
- Misoprostol, 222, 228
- Mivacurium, 142
- Monoamine oxidase (MAO), 25, 53, 54  
forms of, 58  
metabolism of, 11
- Monoclonal antibodies, clinical uses, 304
- Montelukast, 240
- Morphine  
characteristics/uses, 145  
contraindications/cautions, 145  
as drug of abuse, 160  
duality of action, 223  
metabolism of, 11
- Motion sickness, drugs for treating, 50, 220
- Muromonab, clinical uses, 304
- Muscarinic agonists, 47, 69, 223
- Muscarinic receptors, 39  
activators, 46, 47–49  
mechanism of action, 47
- Myasthenia, drugs for treating, acetylcholinesterase inhibitors, 48
- Mycobacterium avium-intracellulare** (MAC), 186
- Mycobacterium tuberculosis**, 186
- Mycophenolate, 303
- Mydriasis, 43f
- N**
- N-acetylcysteine, 231
- $\text{Na}^+$  channel blockers, 89–90, 117
- $\text{Na}^+$  channel toxins, 141
- $\text{Na}^+$  channels  
activation, 86, 86f  
fast-response fibers, 83f, 83–84  
recovery, 86  
slow-response fibers, 84–85
- Na/K-ATPase, 25
- Nafcillin, 177
- Nalbuphine, characteristics/uses, 146
- Naloxone  
as antidote, 311  
characteristics/uses, 146
- Naltrexone, characteristics/uses, 146
- Naproxen, 230
- Natural medicinals, 312–313
- Nausea, drugs for treating, 220, 223, 243
- Nedocromil, in treating asthma, 240
- Nelfinavir, mechanism of action, 195
- Neostigmine, characteristics/uses, 48

- Nephrogenic diabetes insipidus, thiazides for, 110  
 Nephrolithiasis, thiazides for, 110  
 Nephrotoxicity, 133f  
 Nesiritide, for heart failure, 101  
 Neuraminidase, 25  
 Neurotransmitters  
   autonomic nervous system, 40  
   drugs of abuse and, 159–160  
 Nevirapine, mechanism of action, 195  
 Niacin  
   as antihyperlipidemic, 114  
   summary of antihyperlipidemic uses, 115  
 Nicotinic acid, 117  
   as antihyperlipidemic, 114  
 Nicotinic receptor antagonists, 50–51, 142  
   ganglion blocking agents, 50–51  
   neuromuscular blocking drugs. *See* CNS pharmacology  
 Nicotinic receptors, 23, 39  
   activation, 47, 142–143  
   mechanism of action, 47  
   subunits, 142  
 Nifedipine, as calcium-channel blocker, 95  
 Nifurtimox, as antiprotozoal agent, 203  
 Nitrates, in antianginal drugs, 103f, 103–104  
 Nitric oxide signaling, 25, 94–95, 103–104  
 Nitroglycerine  
   as antianginal, 104  
   first-pass effect, 7  
 Nitroprusside, as direct-acting vasodilator, 95  
 Nitrosoureas, 295f  
 Nitrous oxide, characteristics, 139  
 NNRTIs (nonnucleoside reverse transcriptase inhibitors), 197  
 Noncompetitive antagonists, 21, 22  
 Nonmicrosomal metabolism, 11  
 Nonnucleoside reverse transcriptase inhibitors (NNRTIs), 197  
 Nonsteroidal anti  
 Nonsteroidal antiinflammatory drugs (NSAIDs). *See* NSAIDs  
   (nonsteroidal antiinflammatory drugs)  
 Nordiazepam, 9  
 Norepinephrine (NE), 40  
   clinical uses, 58  
   cocaine in blocking uptake, 141  
   dopamine as precursor, 149  
   drugs of abuse and, 159  
   as indirect-acting adrenergic receptor agonist, 58  
   as mixed-acting adrenergic agonist, 56, 56f  
   nicotinic receptor activation, 47  
   receptors for, 24  
   reuptake inhibitors, 25  
 Norethindrone, 277  
 Norgestrel, 277  
 NRTIs (nucleoside reverse transcriptase inhibitors), 197, 198  
 NSAIDs (nonsteroidal antiinflammatory drugs), 229–231  
   drug list, 243  
   for migraine headaches, 226  
   for rheumatoid arthritis, 234  
 Nucleic acid synthesis inhibitors, 184–186  
   direct, 185–186  
   folic acid synthesis inhibitors, 175, 184–185  
   mechanism of action, 175  
 Nucleoside reverse transcriptase inhibitors (NRTIs), 197  
   side effects, 198  
 Nystatin, characteristics/uses, 191–192
- O**
- Olanzapine, characteristic properties, 153, 225  
 Omeprazole  
   cytochrome P450 isozymes, 10  
   for treating *H. pylori* ulcers, 186, 222  
 Ondansetron, 225  
 Ophthalmology. *See also* Glaucoma, drugs for treating  
   effect of ANS drugs on eye, 43f  
   M blockers, 50  
 Opioid analgesics, 145–147  
   characteristics/uses, 145–147  
   common toxic syndromes, 309  
   contraindications/cautions, 145  
   drug list, 163  
   as drugs of abuse, 146–147, 160  
   duality of action, 223  
   for migraine headaches, 226  
   naloxone as antidote, 311  
 Oral administration, doubling oral doses, 15  
 Oral contraceptives, characteristics/uses, 277–278  
 Organophosphates  
   characteristics/uses, 48  
   toxicology, 48  
 Orlistat, as antihyperlipidemic, 115, 117  
 Oseltamivir  
   characteristics/uses, 199  
   mechanism of action, 195, 199  
 Osmotic diuretics, 108  
 Oxacillin, 177  
 Oxazepam, characteristics/uses, 130  
 Oxygen, as antidote, 311
- P**
- Paclitaxel, 295f  
 Palivizumab, clinical uses, 304  
 Paraoxon, 48  
 Parasympathetic nervous system (PANS), 39, 40  
   blood pressure control mechanisms, 40–41  
   characteristics of ganglion blockers, 51  
   nicotinic receptor activation, 47  
 Parathion, 46, 48  
 Parenteral formulations, 153  
 Parmlintide, 274  
 Paroxetine, 156  
 Partial agonists, 20, 20f, 21, 21f  
 PCP, as drug of abuse, 160  
 Penicillamine  
   as antidote, 311  
   characteristics/uses, 233  
   for heavy metal poisoning, 310  
 Penicillins  
   characteristics/uses, 176f, 177  
   drug list, 205

- mechanism of action, 175, 177
- mechanism of resistance, 176, 177
- Pentazocine, characteristics/uses, 146
- Peptic ulcer disease (PUD), drugs for treating, 186, 221–224, 243
- Permeation, in pharmacokinetics, 3f, 4–5
  - factors dependent on, 4
  - ionization, 4f, 4–5, 5f
  - modes of drug transport across a membrane, 5
- pH levels, 4f, 4–5
  - urinary, 5
- Pharmacodynamics, 19–27
  - definitions, 19
  - drug development and testing, 26
  - graded (quantitative) dose-response (D-R) curves, 19–22
  - quantal (cumulative) dose-response (D-R) curves, 22–23
  - signaling mechanisms, 23–25
- Pharmacokinetics, 3–18
  - absorption, 3f, 6–7, 89, 223
  - biotransformation, 3f, 9f, 9–11, 133f
  - calculations, 17
  - distribution, 3f, 8–9
  - drug biodisposition, 3f
  - elimination, 3f, 12–14
  - metabolism in, 3f, 9–11, 133f
  - permeation, 3f, 4–5
  - redistribution, 9, 9f
  - steady state, 3f, 14–16
- Phenelzine, 155
- Phenobarbital, 130
- Phenoxybenzamine, as alpha blocker, 58
- Phentolamine, as alpha blocker, 58
- Phenylephrine, as  $\alpha_1$  adrenergic agonist, 55, 55f
- Phenytoin
  - characteristics/uses, 135–136
  - cytochrome P450 isozymes, 10
  - seizure states/drugs of choice, 135
- Philocarpine, muscarinic agonist, 47
- Phosphatidylinositol biphosphate (PIP<sub>2</sub>), 24
- Phosphodiesterase (PDE) inhibitors
  - as antianginal, 105f
  - for heart failure, 101
  - for hypertension, 97
- Phospholipase, 24
- Phototoxicity, 182
- Physostigmine
  - as antidote, 50, 311
  - characteristics/uses, 48
- Pilocarpine, mechanism of action, 62
- Pindolol, characteristics, 59
- Pioglitazone, characteristics/uses, 273
- Piperacillin, 177
- Pituitary disorders, drugs for treating
  - characteristics/uses, 287
  - drug list, 287
- pKA levels, 4f, 4–5
- Placental barrier to distribution, 8
  - Food and Drug Administration (FDA) pregnancy risk categories, 26
- Plasma level curves, 6, 6f
  - bioavailability, 7, 7f
- Plasma-protein-binding sites, 8
- Plateau principle, 14, 14f
- Platelet-derived growth factor (PDGF), 25
- Polyenes, 191–192
- Potassium
  - as antiarrhythmic, 91
  - direct-acting vasodilators, 95
  - hypokalemia, 107
  - K<sup>+</sup>-sparing agents as diuretics, 111–112, 117
- Potency
  - antagonism and potentiation, 21f, 21–22, 22
  - defined, 19
  - of full and partial agonists, 20f
- Pralidoxine, as antidote, 311
- Praziquantel, 204
- Prazosin, as alpha blocker, 59, 94
- Prednisone, 237
- Pregnancy risk
  - alcohols, 133
  - anti-HIV agents, 199
  - antibacterial agents, 183
  - Food and Drug Administration (FDA) categories, 26
  - prostaglandins (PGs), 228
- Premarin (conjugated estrogens), 275
- Presynaptic membrane depolarization, 45
- Primaquine, 204
- Probenecid, 186
  - characteristics/uses, 235–236
- Procainamide
  - characteristics/uses, 89
  - metabolism of, 11
  - as Na<sup>+</sup> channel blocker, 89
- Procaine, as local anesthetic, 141
- Procarbazine
  - characteristics/uses, 296
  - toxicity, 297
- Prochlorperazine, 223
- Prodrugs, 9, 235
- Progesterin
  - characteristics/uses, 277
  - drug list, 277, 287
  - in oral contraceptives, 277–278
- Promethazine
  - as antiemetic, 223
  - properties, 220
- Propofol, characteristics/uses, 140
- Propranolol
  - as beta-blocker, 90
  - characteristics/uses, 59, 60, 90
  - for migraine headaches, 226
- Propylthiouracil (PTU), placental barrier to distribution, 8
- Prostacyclin
  - for pulmonary hypertension, 97, 229
  - receptors for, 24
- Prostaglandins (PGs), characteristics/uses, 227, 228–229
- Protamine, as antidote, 311

Protease inhibitors (PI)  
 characteristics/uses, 198–199  
 mechanism of action, 198

Protein-binding, 8

Protein C deficiency, 259f

Protein kinase, 23

Protein kinase C, 24

Proton pump inhibitors, 222

Pulmonary hypertension, drugs for treating, 97

Pupillary size and accommodation mechanisms, 43f

Purine, resistance to androgen receptors, 298

Pyrantel pamoate, 204

Pyrazinamide, 186, 187

Pyridostigmine, characteristics/uses, 48

Pyrimethamine  
 as antiprotozoal agent, 203  
 characteristics/uses, 185  
 folic acid synthesis inhibitors, 184  
 mechanism of action, 175

Pyrimidine antimetabolites, resistance to androgen receptors, 298

## Q

Quantal (cumulative) dose-response (D-R) curves, 22–23  
 therapeutic index (TI), 22f, 22–23

Quantitative D-R curves. *See* Graded (quantitative) dose-response (D-R) curves

Quinidine  
 apparent volume of distribution ( $V_d$ ), 8  
 characteristics/uses, 89  
 cytochrome P450 isozymes, 10  
 for heart failure, 101  
 as M blocker, 50, 89  
 as  $\text{Na}^+$  channel blocker, 89

Quinolones  
 characteristics/uses, 182, 185  
 cytochrome P450 isozymes, 10

Quinupristin-dalfopristin  
 characteristics/uses, 183–184  
 mechanism summary, 181

## R

Raloxifene, 276–277

Ranitidine, characteristics/uses, 221–222

Recovery  
 $\text{Na}^+$  channels, 86  
 rates of onset/recovery for inhaled anesthetics, 140

Redistribution, of lipid-soluble drugs, 9, 9f

Refractoriness, cardiac action potentials, 85

Relative refractory period (RRP), 85

Renal clearance of drugs, 4f, 5, 5f

Renal elimination, 14

Renal failure, diuretics for treating, 109

Repaglinide  
 characteristics/uses, 272  
 modes of action, 273f

Reserpine, 155  
 characteristics/uses, 93

Responsiveness, cardiac action potentials, 84

Resting potential (RP), 86

Reverse transcriptase, 25

Reverse transcriptase inhibitors (RTIs), 197–198

Rheumatoid arthritis, drugs for treating, 233–234, 244

Ribavirin  
 characteristics/uses, 199–200  
 mechanism of action, 195, 199

Rifampin, 186, 187  
 cytochrome P450 isozymes, 10  
 mechanism of action, 175

Risperidone, characteristic properties, 153

Ritodrine, as beta adrenergic agonist, 56

Ritonavir  
 cytochrome P450 isozymes, 10  
 mechanism of action, 195  
 side effects, 199

Rituximab, clinical uses, 304

Rosiglitazone, characteristics/uses, 273

## S

St. John's wort, characteristics/uses, 312

Salicylates  
 characteristics/uses, 229–230  
 common toxic syndromes, 309

Salmeterol, as beta adrenergic agonist, 56, 239

Saquinavir  
 characteristics/uses, 198  
 mechanism of action, 195

Sargramostim (GM-CSF), clinical uses, 297

Saw palmetto, characteristics/uses, 312

Saxitoxin, 141  
 mode of action, 141f

Schizophrenia  
 drugs for treating, 152  
 symptoms, 152

Scopolamine, characteristics/uses, 50, 223

Sedative-hypnotic-anxiolytic drugs, 129–133  
 alcohols. *See* Alcohols  
 characteristics/uses, 129–131  
 CNS effects based on dosage, 129f  
 common toxic syndromes, 309  
 drug list, 163  
 as drugs of abuse, 159–160  
 interaction with GABA complex, 129f

Selective COX 2 inhibitors, 109, 231

Selegiline, as antiparkinsonian drug, 150f, 151

Serotonin, reuptake inhibitors, 25

Serotonin receptors, 24, 223

Serotonin syndrome, 155

Sertraline, 156

Signal transducers and activators of transcription (STAT), 25

Signaling mechanisms, 23–25  
 cyclic GMP and nitric oxide signaling, 25, 94–95  
 intracellular receptors, 23  
 membrane receptors directly coupled to ion channels, 23  
 receptors for cytokines, 25  
 receptors linked via coupling proteins to intracellular effectors, 23–24

- receptors that function as enzymes or transporters, 25
  - receptors that function as transmembrane enzymes, 25
  - Sildenafil (Viagra)
    - phosphodiesterase 5 (PDE5) inhibition, 97, 104
    - for pulmonary hypertension, 97
    - toxicity, 104
  - Sitagliptin, 274
  - 6-Mercaptopurine
    - cell-cycle specificity, 295f, 296
    - characteristics/uses, 296
  - 6-Thioguanine, cell-cycle specificity, 295f
  - Skeletal muscle relaxants, 142–143
    - centrally acting, 143
    - characteristics/uses, 142–143
    - depolarizing (noncompetitive), 142–143
    - drug list, 163
    - nicotinic acid receptor, 142f
    - nondepolarizing (competitive), 142
  - Smooth muscle contraction/relaxation, 105f
  - Sodium nitrate, for cyanide poisoning, 95
  - Solubility, 4
  - Sotalol, as K<sup>+</sup> channel blocker, 60, 91
  - Spasm of accommodation, 43f
  - Spironolactone
    - as adrenal steroid antagonist, 275
    - as diuretic, 111
    - for heart failure, 100
    - modes of action/effects, 112
  - SSRIs (selective serotonin reuptake inhibitors), 156
    - characteristics/uses, 156
    - common toxic syndromes, 309
  - Statins, 113–114, 117
    - summary of antihyperlipidemic uses, 115
  - Stavudine
    - mechanism of action, 195
    - side effects, 198
  - Steady state, in pharmacokinetics, 3f, 14–16
    - infusion rate, 15, 15f, 17
    - loading dose (LD), 16, 16f, 17
    - plateau principle, 14, 14f
  - Steroids, 275–279
    - adrenosteroids, 275, 287
    - androgens, 278, 287
    - drug list, 287
    - estrogens, 275–277, 287
    - intracellular receptors, 23
    - minimizing toxicity, 237
    - oral contraceptives, 277–278
    - progestins, 277, 287
  - Stibogluconate, as antiprotozoal agent, 203
  - Streptogramins, mechanism of action, 175
  - Streptokinase
    - actions, 261f
    - characteristics/uses, 261
  - Streptomycin, 181, 186, 187
    - mechanism of resistance, 176
  - Succimer, for heavy metal poisoning, 310
  - Succinylcholine, 142
    - malignant hyperthermia as side effect, 143
  - Sucralfate, 222
  - Suicide inhibitors, 177
  - Sulbactam, 177
  - Sulfa antibiotics, 109
  - Sulfadiazine, as antiprotozoal agent, 203
  - Sulfasalazine
    - characteristics/uses, 184, 233
    - metabolism/uses, 184
  - Sulfonamides
    - allergic reactions, 109
    - characteristics/uses, 182, 184, 185
    - mechanism of action, 175
    - mechanism of resistance, 176
    - pregnancy risk, 183
    - protein-binding capacity, 8
  - Sulfonylureas
    - characteristics/uses, 272
    - drug list, 287
    - mode of action, 272f
  - Sulindac, 230
  - Sumatriptan, 225
  - Surface area, 4
  - Sympathetic autonomic nervous system (SANS), 39, 40
    - blood pressure control mechanisms, 40–41
    - characteristics of ganglion blockers, 51
    - drugs altering sympathetic activity, 93–94
    - nicotinic receptor activation, 47
  - Sympathomimetics, for heart failure, 101
  - Synapse, 54
  - Synaptobrevin, 45
  - Synthesis inhibitors, 275
  - Systolic pressure, 42f
- ## T
- Tacrine, characteristics/uses, 48
  - Tacrolimus, characteristics/uses, 303
  - Tamoxifen, 276–277
  - Tamsulosin, as alpha blocker, 59
  - Telithromycin, 183
  - Temazepam, characteristics/uses, 130
  - Teratogenicity, 26
  - Terazosin, as alpha blocker, 59, 94
  - Terbinafine, characteristics/uses, 193
  - Terbutaline, as beta adrenergic agonist, 56
  - Tetracyclines
    - characteristics/uses, 182
    - drug list, 205
    - mechanism of action, 175
    - mechanism of resistance, 176
    - mechanism summary, 182
    - pregnancy risk, 183
  - Tetrodotoxin, 141
    - mode of action, 141f
  - Theophylline
    - cytochrome P450 isozymes, 10
    - esmolol as antidote, 311
    - in treating asthma, 240
  - Therapeutic Index (TI), 22f, 22–23



- Thiazides, 109, 110–111, 117  
actions on distal convoluted tubule (DCT), 110f
- Thiazolidinediones, characteristics/uses, 273, 287
- Thiopental, 130  
characteristics/uses, 140
- Thioridazine, characteristic properties, 153
- Thrombolytics  
characteristics/uses, 261–262  
drug list, 265
- Thrombopoietin, clinical uses, 297
- Thromboxanes, 229
- Thyroid hormones, intracellular receptors, 23
- Thyrotoxicosis, drugs for treating, 60
- Ticarcillin, 177
- Ticlopidine  
characteristics/uses, 264  
platelet activation, 263f
- Tilolol, mechanism of action, 62
- Timolol, characteristics, 59, 60
- Tirofiban, characteristics/uses, 264
- Tobramycin, 181
- Tocainide, as Na<sup>+</sup> channel blocker, 90
- Tolbutamide, 272
- Tolcapone, as antiparkinsonian drug, 151
- Torsades, drugs for treating, 91
- Torseamide, modes of action/effects, 112
- Total peripheral resistance (TPR), 40, 41f
- Toxicology, 309–315  
acetaldehyde toxicity, 133f  
of AChE inhibitors, 48–49  
of anticancer drugs, 297  
antidotes, 310, 311  
of antimuscarinics, 50  
common toxic syndromes, 309  
cyanide poisoning, 95  
drug development and testing, 26  
of drugs of abuse, 159–160  
heavy metal poisoning, 310, 311  
minimizing steroid toxicity, 237  
Na<sup>+</sup> channel toxins, 141  
natural medicinals, 312–313  
nephrotoxicity, 133f  
phototoxicity, 182  
pregnancy and, 26, 133, 183, 199, 228  
teratogenicity, 26  
Therapeutic Index (TI), 22f, 22–23  
“tot” toxicity and, 231
- Toxoplasmosis, pyrimethamine + sulfadiazine for, 203
- Tranexamic acid, 261
- Transmembrane enzymes, receptors that function as, 25
- Transporters, receptors that function as, 25
- Tranylcypromine, 155
- Trastuzumab, clinical uses, 304
- Trazodone, 156
- Triamcinolone, 237
- Triamterene  
as diuretic, 111  
modes of action/effects, 112
- Trichomoniasis, metronidazole for, 203
- Tricyclic antidepressants  
characteristics/uses, 155  
common toxic syndromes, 309  
as indirect-acting adrenergic agonist, 58  
M blockers, 50
- Trihexyphenidyl  
as antiparkinsonian drug, 151  
characteristics/uses, 50
- Trimethoprim, mechanism of action, 175
- Trimethoprim-sulfamethoxazole, characteristics/uses, 184, 185
- Tropicamide, characteristics/uses, 50
- Trypanosomiasis  
arsenicals for, 203  
nifurtimox for, 203
- Tyramine, 155  
as indirect-acting adrenergic agonist, 58  
nonmicrosomal metabolism, 11
- Tyrosine, 53f
- Tyrosine hydroxylase, 53
- ## V
- Valacyclovir, 196
- Valproic acid  
characteristics/uses, 136–137  
for migraine headaches, 226  
seizure states/drugs of choice, 135
- Vancomycin  
characteristics/uses, 179–180  
mechanism of action, 175, 179
- Vancomycin-resistant enterococci (VRE), 184
- Vancomycin-resistant staphylococci (VRSA), 184
- Vascularity, 4
- Vasodilators  
acting through nitric oxide, 94–95  
acting to open potassium channels, 95
- Venlafaxine, 156
- Venodilators, for heart failure, 100
- Verapamil  
apparent volume of distribution ( $V_d$ ), 8  
as calcium-channel blocker, 91, 95  
for heart failure, 101  
for migraine headaches, 226
- Viagra (sildenafil)  
phosphodiesterase 5 (PDE5) inhibition, 97, 104  
for pulmonary hypertension, 97  
toxicity, 104
- Vinblastine  
cell-cycle specificity, 295f, 296  
characteristics/uses, 296
- Vinca alkaloids, resistance to, 298
- Vincristine  
cell-cycle specificity, 295f, 296  
characteristics/uses, 296  
toxicity, 297
- Vitamin D, intracellular receptors, 23
- Vitamin K, as antidote, 311
- Vitamin V<sub>3</sub>, as antihyperlipidemic, 114
- Volume of distribution ( $V_d$ ), 8, 17

Vomiting, drugs for treating, 220, 223, 243  
Voriconazole, characteristics/uses, 192

**W**

Warfarin (coumarin)  
  characteristics/uses, 258–259  
  comparison with heparin, 258  
  cytochrome P450 isozymes, 10  
  Vitamin K as antidote, 311  
Weak acids, 4–5  
Weak bases, 4–5  
Wolff-Parkinson-White syndrome, 101

**Y**

Yohimbine, as alpha blocker, 59

**Z**

Zafirlukast, 228, 240  
Zalcitabine  
  mechanism of action, 195  
  side effects, 198  
Zaleplon  
  characteristics/uses, 131  
  flumazenil as antidote, 311  
Zanamivir  
  characteristics/uses, 199  
  mechanism of action, 195, 199  
ZDV, 198, 199  
Zero-order elimination rate, 12, 12f, 13  
Zidovudine, 198  
  mechanism of action, 195  
  side effects, 198  
Zileuton, 228, 240  
Zolpidem  
  characteristics/uses, 131  
  flumazenil as antidote, 311



**KAPLAN**

MEDICAL

[kaplanmedical.com](http://kaplanmedical.com)



BK4032E

Printed in USA. ©2009 Kaplan, Inc.